

Diversity, pathogenicity and two new species of pestalotioid fungi (Amphisphaeriales) associated with Chinese Yew in Guangxi, China

Yifeng Wang¹, Kin-Ming Tsui^{2,3}, Shimei Chen¹, Chongjuan You¹

¹ Beijing Key Laboratory for Forest Pest Control, Beijing Forestry University, Beijing 100083, China

² National Centre for Infectious Diseases, Tan Tock Seng Hospital, Singapore 308433, Singapore

³ Faculty of Medicine, University of British Columbia, Vancouver, V6T 1Z3, Canada

Corresponding author: Chongjuan You (chongjuanyou@bjfu.edu.cn)

Abstract

Chinese yew, *Taxus chinensis* var. *mairei* is an endangered shrub native to south-eastern China and is widely known for its medicinal value. The increased cultivation of Chinese yew has increased the incidence of various fungal diseases. In this study, Pestalotioid fungi associated with needle spot of Chinese yew were isolated from Guangxi Province. Based on morphological examinations and multi-locus (ITS, *tub2*, *tef-1a*) phylogenies, these isolates were identified to five species, including two new species, *Pestalotiopsis taxicola* and *P. multicolor*, two potential novel *Neopestalotiopsis* species, *Neopestalotiopsis* sp. 3 and *Neopestalotiopsis* sp. 4, with a known *Pestalotiopsis* species (*Pestalotiopsis trachycarpicola*), firstly recorded from Chinese yew. These two new *Pestalotiopsis* species were morphologically and phylogenetically distinct from the extant Pestalotioid species in Chinese yew. Pathogenicity and culture characteristic tests of these five Pestalotioid species were also performed in this study. The pathogenicity test results revealed that *Neopestalotiopsis* sp. 3 can cause diseases in Chinese yew needles. These results have indicated that the diversity of Pestalotioid species associated with Chinese yew was greater than previously determined and provided helpful information for Chinese yew disease diagnosis and management.

Key words: *Neopestalotiopsis*, *Pestalotiopsis*, phylogeny, taxonomy, two new species



Academic editor: Nattawut Boonyuen

Received: 5 October 2023

Accepted: 27 January 2024

Published: 27 February 2024

Citation: Wang Y, Tsui K-M, Chen S, You C (2024) Diversity, pathogenicity and two new species of pestalotioid fungi (Amphisphaeriales) associated with Chinese Yew in Guangxi, China. MycoKeys 102: 201–224. <https://doi.org/10.3897/mycokeys.102.113696>

Copyright: © Yifeng Wang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The Chinese yew (*Taxus chinensis* var. *mairei* (Lemée et Lév.) Cheng et L.K. Fu) is an evergreen tall tree unique to south-eastern China and is widely known as a significant variety of medicinal plants because it contains taxol (paclitaxel) which is a natural antitumour drug with unique physiological functions in the bark, branches and leaves (Pyo et al. 2004; Li et al. 2008; Wu et al. 2013). Chinese yew has been listed on the IUCN Red List as an endangered species and has been under first-class protection in China, as it grows slowly and has poor regeneration ability, making it unable to meet strong medical demands (Ru 2006; Liu et al. 2019). Many natural reserves and conservation parks have been established to protect the Chinese yew. It has been widely cultivated in 17 provinces of China over the past few decades (Gao 2006; Zhang et al. 2018; Li et al. 2021a).

The expansion of Chinese yew cultivation in recent years has led to the emergence of various diseases that pose a significant threat to the Chinese yew industry (Zhang et al. 2018). Many fungal pathogens have been reported to cause Chinese yew diseases during their growth, such as needle spot caused by *Neopestalotiopsis clavispora* and *Pestalotiopsis affinis* (Chen et al. 2002; Wang et al. 2019), needle blight caused by *Pestalotiopsis microspora* (Li 2017), anthracnose disease caused by *Colletotrichum gloeosporioides* (Fan et al. 2006) and wilt disease caused by *Phoma* sp., while *Fusarium oxysporum* is responsible for root rot (Qian et al. 2015). Needle diseases caused by Pestalotioid fungal species (e.g. *N. clavispora*, *P. affinis* and *P. microspora*) are a serious threat to Chinese yew in Fujian and Yunnan Provinces, China (Chen et al. 2002; Li 2017; Wang et al. 2019). Up to 40% of the Chinese yew are severely affected in the field (Wang et al. 2019). The needles of Chinese yew became brown and reddish-brown from the leaf margin to the main vein, with irregular or round spots. Eventually, the spots coalesced and the needles withered and abscised (Li 2017; Wang et al. 2019). Jeon and Cheon (2014) reported that *P. microspora* also caused leaf blight in Japanese yew (*Taxus cuspidata* Sieb. & Zucc.) in South Korea. This disease developed on needles, particularly under high humidity, producing brown or tan spots that coalesce to form large lesions (Jeon and Cheon 2014).

Pestalotioid fungi (*Pestalotiopsis*-like fungi) represent an important fungal group that commonly occur as plant pathogens, endophytes and saprophytes in a wide range of hosts (Maharachchikumbura et al. 2014; Reddy et al. 2016; Ran et al. 2017; Freitas et al. 2019; Yang et al. 2021; Xiong et al. 2022). This group, which comprise *Neopestalotiopsis*, *Pestalotiopsis* and *Pseudopestalotiopsis*, belongs to the order Amphisphaeriales (Jiang et al. 2022; Zhang et al. 2022). The two genera *Neopestalotiopsis* and *Pseudopestalotiopsis* were segregated from *Pestalotiopsis* by Maharachchikumbura et al. (2014) based on multigene phylogenetic analysis and morphological differences (conidiogenous cells and colour intensities of the median conidial cell). *Pestalotiopsis* is characterised by lightly pigmented concolorous median cells, whereas *Neopestalotiopsis* species are characterised by versicolorous median cells and indistinct conidiophores and *Pseudopestalotiopsis* by darkly coloured concolorous median conidial cells (Maharachchikumbura et al. 2014). Many *Pestalotiopsis* species have been isolated from *Taxus* spp. as endophytes rich in secondary metabolites. For example, *Pestalotiopsis versicolor* has been isolated from healthy leaves of *Taxus cuspidata* and is an excellent alternative source of paclitaxel supply (Kumaran et al. 2010). Furthermore, *P. microspora*, which causes Chinese yew leaf blight, is an endophyte of the Himalayan yew (*Taxus wallichiana*), which produces taxol (Strobel et al. 1996).

During a survey of diseases in Chinese yew, moderate-to-severe incidences of needle spot and stem canker diseases were observed in some planting areas in Guangxi Province in 2020, and several Pestalotioid fungi were isolated from the diseased Chinese yew. The objectives of our study were to: (i) identify the Pestalotioid fungi recovered from symptomatic Chinese yew using morphological features and molecular data analyses; (ii) evaluate the pathogenicity of different fungal species on detached Chinese yew; and (iii) evaluate the effects of temperature, light duration and carbon source on the mycelial growth rate of different fungal species.

Materials and methods

Sample collections and Fungal Isolation

In May 2020, twenty symptomatic Chinese yew specimens were collected from the Guangxi Province. Small sections (5 × 5 mm) were cut from the margins of the infected needles and stems, surface-sterilised in 75% ethanol for 30 s, sterilised in 3% (vol/vol) sodium hypochlorite for 1 min, followed by three rinses in sterile distilled water and finally dried on sterilised filter paper (Rubini et al. 2005; Li et al. 2021b). The sections were plated on PDA plates and incubated at 25 °C. The isolates were aseptically transferred to fresh PDA and purified using a single-spore culture. The holotype specimens in this study are deposited in the Museum of Beijing Forestry University (Herbarium Code: BJFC). The pure fungal cultures in this study are deposited in China Forestry Culture Collection Center (Acronym: CFCC).

Morphological and culture characterisation

The colony characteristics (colour and texture) of each isolate on PDA were observed after 7 d incubation at 25 °C and the morphological characters of conidiophores, conidiogenous cells and conidia of each isolate on PDA were observed after 14 d incubation at 25 °C (Li et al. 2021b). The size was determined by measuring the length and width of 50 randomly chosen conidia using an Olympus SZX2-FOF Light microscope (Tokyo, Japan) according to the method described by Shu et al. (2020). Descriptions, nomenclature and illustrations of taxonomic novelties have been deposited in MycoBank (<https://www.mycobank.org/>).

DNA extraction and PCR amplification

Genomic DNA was extracted from 7 day-old colonies grown on PDA using the cetyltrimethylammonium bromide (CTAB) method (Doyle and Doyle 1990). Three loci were amplified: the internal transcribed spacer region of ribosomal DNA (ITS) with primers ITS1/ITS4 (White et al. 1990); beta-tubulin (*tub2*) with primers Bt2a/Bt2b (Glass and Donaldson 1995) and the translation elongation factor 1-alpha gene (*tef-1a*) with EF1-688F/EF1-1251R (Alves et al. 2008). The PCR mixture consisted of 10 µl TopTaq Master Mix, 7 µl nuclease-free H₂O, 1 µl of each primer and 1 µl DNA samples were made up to the final volume of 20 µl. Primer sequences and reaction conditions were shown in Suppl. material 1. The PCR products were sent for sequencing at Beijing Tsingke Biotech Company, Ltd., Beijing, China. The DNA sequences have been deposited in GenBank (Suppl. material 2).

Sequence alignment and phylogenetic analyses

The taxa used in the analyses were obtained from the sequence data of *Pestalotiopsis* and *Neopestalotiopsis* downloaded from GenBank (Suppl. material 2). Sequence alignments were performed using MAFFT v.7 (mafft.cbrc.jp/alignment/server) (Kato and Standley 2013). Phylogenetic analyses, based on a combined dataset of ITS, *tub2* and *tef-1a* sequence data, were performed with Maximum Parsimony (MP), Maximum Likelihood (ML) and Bayesian Inference (BI). MP analysis was performed with PAUP 4.0b using a heuristic search algorithm (1,000 random-addition sequences) with tree bisection and reconnection

(TBR) branch swapping. MaxTrees were set to 5,000, branches of zero length were collapsed and all equally parsimonious trees were saved (Swofford 2003). ML analysis was performed using RAxML-HPC BlackBox v. 8.2.12, with 1000 rapid bootstrap replicate runs using the GTRGAMMA model of nucleotide evolution (Stamatakis 2014). For BI analysis, the best-fit evolutionary models for each partitioned locus were deduced on the AIC (ITS: GTR+I+G, *tub2*: HKY+I+G and *tef-1 α* : TrN+I+G) which were estimated in MrModelTest v. 2.3. The BI analysis was performed using MrBayes v.3.1.2, with a Markov Chain Monte Carlo (MCMC) algorithm running from random trees for 1,000,000 generations. The resulting trees were edited using FigTree v.1.4.2 (Rambaut 2016) and Adobe Illustrator CS5.

Culture characteristics

Five Pestalotioid species were selected to evaluate the effects of temperature, light duration and carbon source on mycelial growth. Mycelial plugs of each species were taken from the colony margins of 3-day-old cultures and transferred to fresh plates (20 ml of culture medium, pH 6). The effects of temperature were tested in the range 19 to 31 °C with a 3 °C gradient on PDA and incubated in dark. To monitor the effects of light duration, the isolates were tested under three diel light cycles (24 h dark, 12 h light/12 h dark and 24 h light per day) on PDA and incubated at 25 °C. The effects of the carbon source were tested using sucrose, maltose and dextrose with the same carbon content. Three replicates were used for the temperature and light duration tests and two replicates were used for the carbon test. The colony diameters were measured daily.

Pathogenicity testing

Healthy needles were collected from five-year old and healthy Chinese yew grown in a greenhouse. The needles were washed with tap water, submerged in 70% ethanol for 2 min and rinsed twice with sterile water (Li et al. 2021b). Clean dry needles were placed on Petri dishes with moist filter paper underneath. Puncture wounds were made in the middle of each needle using a sterilised needle. The wounded needles were inoculated with mycelial PDA plugs (3 mm in diameter) and spore suspensions (18 μ l, 10^6 conidia/ml) of five Pestalotioid isolates, respectively. Three replicates were used for the mycelial PDA plug test and 10 replicates were used for the spore suspension test. Lesion length was recorded daily.

Results

Field observations and fungal isolation

A field survey was conducted from April to October 2020 in a forest farm in northern Guangxi, China, where more than 500 Chinese yew trees were maintained. Needle spot disease in Chinese yew causes serious damage, with an incidence of 58–70% in plants. The spots on the needles were initially small, brown to black and oval to irregular. Subsequently, they gradually expand and finally coalesce, forming large black spots. In severe cases, lesions can develop in large portions of a single needle. Over time, the heavily infected leaves dried and died. Lesions were also visible on the stems and the spots on the stems darkened and became necrotic (Fig. 1).

Thirty-six isolates were obtained from the diseased needles and stems of Chinese yew and 18 isolates of Pestalotioid fungi were identified based on their culture characteristics and conidial morphology. Eight isolates were grouped into *Neopestalotiopsis* species with versicolorous median cells and ten isolates were identified as *Pestalotiopsis* with lightly pigmented concolorous median cells. Colonies of most Pestalotioid stains were initially whitish and later greyish or yellow on PDA. Conidia varied from 14.5 to 25.0 μm mean length and 3.5 to 8.0 μm mean width. The apical appendages showed the largest variation in size, with a mean length of 5.0 to 23.0 μm . The number of apical appendages varied between two and four, with three being the most common. Basal appendages were hyaline, straight or slightly curved and varied from 1.0 to 5.0 μm mean length.

Phylogenetic analyses

The phylogenetic tree (*Pestalotiopsis*), based on the concatenated sequences of ITS, *tub2* and *tef-1a*, comprised 121 ingroups and one outgroup, *Pseudopestalotiopsis cocos* (CBS 272.29). A total of 1,496 characters including gaps (401 for ITS, 313 for *tub2* and 860 for *tef-1a*) were included in the phylogenetic analysis. Similar tree topologies were obtained using the MP, ML and BI methods and the best-scoring MP tree obtained from a heuristic search with 1000 random taxon additions is presented (Fig. 2). The phylogenetic tree placed the six *Pestalotiopsis* isolates into three well-supported monophyletic clades, representing two novel and one known species. Two of our isolates (CFCC59981 and CFCC59982), described as *Pestalotiopsis multicolor*, formed monophyletic groups with high support values (100% MP and ML, 1.00 BI). Two isolates (CFCC59976 and CFCC59978), identified as *Pestalotiopsis taxicola*, constituted an independent and strongly-supported subclade (100% MP and ML, 1.00 BI), sharing close affinity with *Pestalotiopsis unicolor* and *Pestalotiopsis jiangxiensis*. Two strains (BJFUCC42 and BJFUCC42-2) clustered with *Pestalotiopsis trachycarpicola* and were, therefore, described as known species.

The phylogenetic tree (*Neopestalotiopsis*), based on the concatenated gene sequences of ITS, *tub2* and *tef-1a*, comprised 95 ingroups and one outgroup, *Pestalotiopsis diversiseta* (MFLUCC 12-0287). A total of 1,386 characters including gaps (411 for ITS, 393 for *tub2* and 582 for *tef-1a*) were included in




Figure 1. a healthy Chinese yew needles b, c chinese yew needles with spots.



Figure 2. A phylogenetic tree of *Pestalotiopsis* generated from MP analysis, based on combined ITS, *tub2* and *tef1a* sequence data. Maximum Parsimony and Maximum Likelihood bootstrap values $\geq 65\%$ are given at the nodes. The branches with Bayesian Inference posterior probabilities ≥ 0.90 are bold. Strains from this study are marked in blue. Ex-type strains are labelled with *.

the phylogenetic analysis. Similar tree topologies were obtained using the MP, ML and BI methods and the best-scoring MP tree obtained from a heuristic search with 1000 random taxon additions is presented (Fig. 3). Two of our isolates (CFCC59989 and CFCC59990) clustered together and nested in a clade containing two different *Neopestalotiopsis* sp. 2 isolates (CFCC 54340 and ZX22B). In contrast, two isolates (CFCC59985 and CFCC59986), grouped *Neopestalotiopsis rhapsidis* (Fig. 3).



Figure 3. A phylogenetic tree of *Neopestalotiopsis* generated from MP analysis, based on combined ITS, *tub2* and *tef1a* sequence data. Maximum Parsimony and Maximum Likelihood bootstrap values $\geq 65\%$ are given at the nodes. The branches with Bayesian Inference posterior probabilities ≥ 0.90 are bold. Strains from this study are marked in blue. Ex-type strains are labelled with *.

Taxonomy

Pestalotiopsis trachycarpicola Yan M. Zhang & K.D. Hyde, 2012

Fig. 4

Conidiogenesis. Conidiophores reduced to conidiogenous cells, indistinct. Conidiogenous cells were discrete, ampulliform, thin-walled, hyaline, smooth. Conidia fusiform to clavate, straight or slightly curved, olivaceous to brown, 4-septate, $18.5\text{--}25 \times 4\text{--}6 \mu\text{m}$, with apical and basal appendages. Basal cell obconic, hyaline, thin-walled, smooth, $3\text{--}5 \mu\text{m}$; the three median cells dolioform, versicolor, pale brown to brown with septa darker than the rest of the cells, $11.5\text{--}13.5 \mu\text{m}$, the second cell from base $3.5\text{--}6 \mu\text{m}$; the third cell $3.5\text{--}4.5 \mu\text{m}$; the fourth cell $3.5\text{--}5 \mu\text{m}$; apical cell $2.5\text{--}4 \mu\text{m}$, cylindrical, hyaline; 2–4 tubular apical appendages, arising from the apex of the apical cell each at different point, filiform, $5\text{--}15 \mu\text{m}$; basal appendage present most of the time, single, tubular, unbranched, $3.5\text{--}4.5 \mu\text{m}$ (Fig. 4c–e). Sexual morph not observed.

Culture characteristics. Colonies on PDA reaching 90 mm diameter after seven days at $25 \text{ }^\circ\text{C}$, with an undulate and radial edge, with dense aerial mycelium on surface, white to faint yellow on front, pale honey-coloured on the reverse side (Fig. 4a). *Conidiomata acervular* in culture on PDA, globose, $100\text{--}500 \mu\text{m}$ in diameter, solitary or aggregated in clusters, exuding black conidial masses (Fig. 4b).




Figure 4. *Pestalotiopsis trachycarpicola* (BJFUCC42) **a** culture on PDA **b** conidiomata formed on PDA **c–e** conidia. Scale bars: $500 \mu\text{m}$ (**b**); $10 \mu\text{m}$ (**c–e**).

Material examined. CHINA, Guangxi Province, from diseased needles of Chinese yew, May 2020, Y. F. Wang (BJFC-S1955); living cultures BJFUCC42, BJFUCC42-2.

Notes. *Pestalotiopsis trachycarpicola* was originally described from leaves of *Trachycarpus fortunei* in Kunming Botany Garden, Kunming, Yunnan Province, China (Zhang et al. 2012). In the present study, the two isolates clustered with *P. trachycarpicola* and *P. kenyana* with high support values (MP/ML = 86/100). *P. kenyana* (CBS 442.67, ex-type) and *P. trachycarpicola* (OP068, ex-type MFLUCC 12-0263) were demonstrated to be the same species, as there was no genetic distance between the two samples (Suppl. material 3). Morphologically, our new collections resembled *P. trachycarpicola* in colour and size of the conidiogenous cells, conidia and appendages (Zhang et al. 2012). Therefore, we reported the two isolates as a new host record of *P. trachycarpicola* from yews.

***Pestalotiopsis taxicola* Y. F. Wang & C. J. You, sp. nov.**

MycoBank No: 847791

Fig. 5

Etymology. Named after the host species, *Taxus chinensis*.

Holotype. BJFC-S1954.

Conidiogenesis. Conidiophores reduced to conidiogenous cells, indistinct. Conidiogenous cells were discrete, ampulliform, thin-walled, hyaline, smooth. Conidia fusiform to clavate, straight or slightly curved, olivaceous to brown, 4-septate, 16.5–21 × 4–6 µm, with apical and basal appendages. Basal cell obconic, hyaline, thin-walled, smooth, 2.5–4 µm; the three median cells dolioform, versicolor, pale brown to brown with septa darker than the rest of the cells, 10.5–12 µm, the second cell from base 3–4 µm; the third cell 3.5–4 µm; the fourth cell 3.5–4 µm; apical cell 2.5–4 µm, cylindrical, hyaline; 3 tubular apical appendages, arising from the apex of the apical cell each at a different point, filiform, 9.5–15 µm; basal appendage present most of the time, single, tubular, unbranched, 2–5 µm (Fig. 5c–e). Sexual morph not observed.

Culture characteristics. Colonies on PDA reaching 90 mm diameter after seven days at 25 °C, with an undulate and radial edge, with dense aerial mycelium on surface, initially yellow in the centre, becoming white at the margin, with white appressed mycelia radiating outwards (Fig. 5a). *Conidiomata acervular* in culture on PDA, globose, 200–700 µm in diameter, solitary or aggregated in clusters, exuding black conidial masses (Fig. 5b).

Material examined. CHINA, Guangxi Province, from diseased needles of Chinese yew, May 2020, Y. F. Wang (BJFC-S1954, holotype); ex-type living culture CFCC59976, living cultures CFCC59978, CFCC59979 and CFCC59980.

Notes. *Pestalotiopsis taxicola* was phylogenetically placed in a clade encompassing *P. unicolor*, but as a unique lineage with high support (MP/ML = 100/100) (Fig. 2). Compared with *P. unicolor* (MFLUCC 12-0276, ex-type) and *P. taxicola* (CFCC59976, ex-type), there were one nucleotide difference in the ITS region and nine nucleotide differences in the *tef-1a* region. Morphologically, *P. taxicola* had smaller conidia (16.5–21 × 4–6 µm) than *P. unicolor* (20–24.5 × 4–6 µm). In addition, *P. taxicola* had only one basal appendage, whereas *P. unicolor* had 1–2 basal appendages (Maharachchikumbura et al. 2012). Therefore, the four isolates were designated as a new species.




Figure 5. *Pestalotiopsis taxicola* (CFCC59976) **a** culture on PDA **b** conidiomata formed on PDA **c** conidiogenous cells **d-f** conidia. Scale bars: 500 µm (**b**); 10 µm (**c-f**).

***Pestalotiopsis multicolor* Y. F. Wang & C. J. You, sp. nov.**

MycoBank No: 847792

Fig. 6

Etymology. In reference to the multicoloured median cells.

Holotype. BJFC-S1956.

Conidiogenesis. Conidiophores reduced to conidiogenous cells, short, subcylindrical and hyaline. Conidiogenous cells were discrete, ampulliform, thin-walled, hyaline, smooth. Conidia fusiform to clavate, straight or slightly curved, olivaceous to brown, 4-septate, $14.5\text{--}20 \times 3.5\text{--}5.5 \mu\text{m}$, with apical and basal appendages. Basal cell obconic, hyaline, thin-walled, smooth, $2\text{--}4.5 \mu\text{m}$; the three median cells dolioform, versicolor, pale brown to brown with septa darker than the rest of the cells, $9.5\text{--}13 \mu\text{m}$, the second cell from base $2.5\text{--}4.5 \mu\text{m}$; the third cell $3\text{--}4.5 \mu\text{m}$; the fourth cell $3\text{--}4 \mu\text{m}$; apical cell $2.5\text{--}3.5 \mu\text{m}$, cylindrical, hyaline; 2–3 (mostly 3) tubular apical appendages, arising from the apex of the apical cell each at a different point, filiform, $8\text{--}16 \mu\text{m}$; basal appendage present most of the time, single, tubular, unbranched, $2.5\text{--}4.5 \mu\text{m}$ (Fig. 6c–e). Sexual morph not observed.

Culture characteristics. Colonies on PDA reaching 90 mm diameter after seven days at 25 °C, with an undulate and radial edge, white aerial mycelium on surface flat or raised (Fig. 6a). *Conidiomata acervular* in culture on PDA, glo-




Figure 6. *Pestalotiopsis multicolor* (CFCC59981) **a** culture on PDA **b** conidiomata form on PDA **c** conidiogenous cells **d–f** conidia. Scale bars: 500 µm (**b**); 10 µm (**c–f**).

bose, 200–800 µm in diameter, solitary or aggregated in clusters, exuding black conidial masses (Fig. 5b).

Material examined. CHINA, Guangxi Province, from diseased needles of Chinese yew, May 2020, Y. F. Wang (BJFC-S1956, holotype); ex-type living culture CFCC59981, living cultures CFCC59982, CFCC59983, and CFCC59984.

Notes. The two isolates (CFCC59981 and CFCC59982) formed a distinct lineage with high support values (MP/ML = 100/100) in the phylogenetic tree. The morphology of the two isolates was distinctive within the *Pestalotiopsis* genus because the conidia were far smaller than those of any other species. In addition, the colour of median cells in our new collections changed from light concolorous to versicolorous. Therefore, four isolates in the present study were designated as a new species.

***Neopestalotiopsis* sp. 3**

Fig. 7

Conidiogenesis. Conidiophores reduced to conidiogenous cells, hyaline, smooth. Conidiogenous cells were discrete, ampulliform, thin-walled, hyaline to light brown, smooth. Conidia fusiform to clavate, straight or slightly curved, olivaceous to brown, 4-septate, 15.5–19 × 6–7.5 µm, with apical and basal appendages. Bas-




Figure 7. *Neopestalotiopsis* sp. 3 (CFCC59985) **a** culture on PDA **b** conidiomata form on PDA **c–e** conidia. Scale bars: 500 µm (**b**); 10 µm (**c–e**).

al cell obconic, hyaline, thin-walled, smooth, 2–4 µm; the three median cells 10.5–12 µm, dolioform, versicolor, pale brown to brown with septa darker than the rest of the cells, the second cell from base 2.5–4.5 µm; the third cell 3.5–4.5 µm; the fourth cell 3–4 µm; apical cell 1.5–3 µm, cylindrical, hyaline; 3–4 (mostly 3) tubular apical appendages, arising from the apex of the apical cell each at a different point, filiform, 16–23 µm; basal appendage present most of the time, single, tubular, unbranched, 1–2.5 µm (Fig. 7c–e). Sexual morph not observed.

Culture characteristics. Colonies on PDA reaching 90 mm diameter after seven days at 25 °C, white aerial mycelium on surface, flat or raised, radiating outwards with an undulate and radial edge (Fig. 7a). *Conidiomata acervular* in culture on PDA, globose, 80–800 µm diameter, solitary or aggregated in clusters, exuding black conidial masses (Fig. 7b).

Material examined. CHINA, Guangxi Province, from diseased shoots of Chinese yew, May 2020, Y. F. Wang (BJFC-S1957); living cultures CFCC59985, CFCC59986, CFCC59987, CFCC59988.

Notes. *Neopestalotiopsis* sp. 3 (CFCC59985 and CFCC59986) was phylogenetically close to *N. rhapsidis* (Yang et al. 2021) (Fig. 3) but differed in conidial size (15.5–19 × 6–7.5 µm in *Neopestalotiopsis* sp. 3 vs. 22–25.5 × 4–6 µm in *N. rhapsidis*). Furthermore, *Neopestalotiopsis* sp. 3 had longer apical appendages (16–23 µm vs. 11–16 µm) and shorter basal appendage (1–2.5 µm vs. 2–5.5 µm) than *N. rhapsidis*. Compared with *N. rhapsidis* (GUCC 21501, ex-type) and *Neopestalotiopsis* sp. 3, there were eleven nucleotide differences in the ITS region, four nucleotide differences in the *tef-1a* region and six nucleotide differences in the *tub2* region. In the genus *Neopestalotiopsis*, the interspe-

cific relationship remains unclear (Li et al. 2021b), as some clades were not unanimously strongly supported, meaning the genus will be largely revised in the near future. Many taxa are likely to be synonymised in future publications. Therefore, CFCC59985 and CFCC59986 were not proposed as a new species.

***Neopestalotiopsis* sp. 4**

Fig. 8

Conidiogenesis. Conidiophores reduced to conidiogenous cells, hyaline, smooth. Conidiogenous cells were discrete, ampulliform, thin-walled, hyaline to light brown, smooth. Conidia fusiform to clavate, straight or slightly curved, olivaceous to brown, 4-septate, $15.5\text{--}18.5 \times 6\text{--}8 \mu\text{m}$, with apical and basal appendages. Basal cell obconic, hyaline, thin-walled, smooth, $2.5\text{--}4 \mu\text{m}$; the three median cells $10.5\text{--}12 \mu\text{m}$, dolioform, versicolor, pale brown to brown with septa darker than the rest of the cells, the second cell from base $3\text{--}3.5 \mu\text{m}$; the third cell $3\text{--}4.5 \mu\text{m}$; the fourth cell $3\text{--}5.5 \mu\text{m}$; apical cell $1\text{--}2.5 \mu\text{m}$, cylindrical, hyaline; 3–4 (mostly 3) tubular apical appendages, arising from the apex of the apical cell each at a different point, filiform, $9.5\text{--}15.5 \mu\text{m}$; basal appendage present most of the time, single, tubular, unbranched, $1.5\text{--}4.5 \mu\text{m}$ (Fig. 8c–e). Sexual morph not observed.

Culture characteristics. Colonies on PDA reaching 90 mm diameter after seven days at $25 \text{ }^\circ\text{C}$, radiating outwards with an undulate and radial edge, white aerial mycelium was flat on the centres of the colony, while raised sharply in the outer ring (Fig. 8a). *Conidiomata acervular* in culture on PDA, globose, $100\text{--}900 \mu\text{m}$ diameter, solitary or aggregated in clusters, exuding black conidial masses (Fig. 8b).




Figure 8. *Neopestalotiopsis* sp. 4 (CFCC59989) **a** culture on PDA **b** conidiomata form on PDA **c–e** conidia. Scale bars: $500 \mu\text{m}$ (**b**); $10 \mu\text{m}$ (**c–e**).

Material examined. CHINA, Gungxi Province, from diseased branches of Chinese yew, May 2020, Y. F. Wang (BJFC-S1958); living cultures CFCC59989, CFCC59990, CFCC59991 and CFCC59992.

Notes. *Neopestalotiopsis* sp. 4 (CFCC59989 and CFCC59990) was phylogenetically placed in a clade encompassing two *Neopestalotiopsis* sp. 2 isolates (CFCC54340 and ZX22B) from Yaan City, Sichuan Province (Jiang et al. 2021) (Fig. 3). Morphologically, *Neopestalotiopsis* sp. 4 had shorter conidia and longer apical appendages ($15.5\text{--}18.5 \times 6\text{--}8 \mu\text{m}$, $9.5\text{--}15.5 \mu\text{m}$) than CFCC54340 and ZX22B ($22\text{--}25.2 \times 6.2\text{--}7.7 \mu\text{m}$, $5\text{--}10 \mu\text{m}$). In addition, *Neopestalotiopsis* sp. 4 had more tubular apical appendages (3–4, mostly 3) than *Neopestalotiopsis* sp. 2 (2, seldom 3). Compared with *Neopestalotiopsis* sp. 2 (ZX22B) and *Neopestalotiopsis* sp. 4, there were six nucleotide differences in the ITS region, five nucleotide differences in the *tef-1a* region and three nucleotide differences in the *tub2* region. The taxonomy of the genera and species in this group remains unclear (Li et al. 2021b), as some clades were not unanimously strongly supported, meaning the genus will be largely revised in near future. Many taxa are likely to be synonymised in future publications. Therefore, CFCC59989 and CFCC59990 were not proposed as a new species.

Culture characteristics

The effects of temperature on the growth of the five Pestalotioid species were shown in Fig. 9. When grown on PDA in the dark, all isolates grew at a temperature range of 19 to 31 °C, with optimum growth between 22 °C and 28 °C. The growth rate was drastically reduced below 22 °C and started to decline above 28 °C. The regression analysis showed the optimum temperatures for *P. trachycarpicola*, *P. taxicola*, *P. multicolor*, *Neopestalotiopsis* sp. 3 and *Neopestalotiopsis* sp. 4 were 25 °C, 24–26 °C, 22–25 °C, 25–28 °C and 25 °C, respectively. Within four days, the diameter of the majority of the isolates was approximately 90 mm, which is the diameter of the PDA plate.

Light duration had no effect on the mycelial growth of the two Pestalotioid species (*P. trachycarpicola* and *P. multicolor*) and the best growth of the three Pestalotioid fungi (*P. taxicola*, *P. multicolor* and *Neopestalotiopsis* sp. 4) was observed under continuous light (Suppl. material 4). The growth rate was highest for *P. taxicola* in 24 h light, which reached 66.4 mm diameter after four days. The growth rate was lowest for *P. trachycarpicola* in 12 h light/12 h dark, which reached only 55.5 mm in diameter after four days.

Five Pestalotioid species grew on all three tested carbon sources (Suppl. material 5). Maltose utilisation was the least efficient and the medium with dextrose showed comparatively high growth of the three tested fungi (*P. trachycarpicola*, *P. multicolor* and *Neopestalotiopsis* sp. 3). For *P. taxicola*, the utilisation of dextrose appeared to be higher than that of sucrose.

Pathogenicity assay

The pathogenicity of two *Neopestalotiopsis* isolates (*Neopestalotiopsis* sp. 3 and *Neopestalotiopsis* sp. 4) and three *Pestalotiopsis* isolates (*P. trachycarpicola*, *P. taxicola* and *P. multicolor*) was tested by inoculating detached healthy needles according to Keith et al. (2003). Dark brown lesions and a




Figure 9. Effects of temperature on growth of isolates.

necrotic zone, which resembled the symptoms that occurred in the field, were observed on the needles 20 days after inoculation with mycelium plugs of *Neopestalotiopsis* sp. 3, which did not occur in the control inoculated with agar media or with the other four *Pestalotioid* species (Fig. 10). *Neopestalotiopsis* sp. 3 was re-isolated from the infected needles and was confirmed to be the same as the inoculated pathogen through morphological and phylogenetic analyses. However, none of the five isolates produced typical symptoms in the wounded needles inoculated with the spore suspension (Fig. 10). The results revealed that only *Neopestalotiopsis* sp. 3 is pathogenic to Chinese yew needles.

Discussion

Cultivation of Chinese yew in some regions (e.g. Guangxi) has resulted in an increase in yew diseases, including new needle spot diseases, which have caused great losses in production. However, very little is known about the pathogens and the presence and prevalence of needle diseases in Chinese yew (Wang et al. 2019). During the investigation of Pestalotioid fungi isolated from diseased Chinese yew in the Guangxi Province of China, *Neopestalotiopsis* sp. 3 was found to cause needle spots on Chinese yew. Two new species of *Pestalotiopsis* were identified in association with symptomatic Chinese yew, namely, *P. taxicola* and *P. multicolor* and one known species, *P. trachycarpicola*, was identified, based on morphology and multi-locus phylogeny. The results indicated that the diversity of Pestalotioid species associated with Chinese yew was greater than previously determined.




Figure 10. Lesions of the Chinese yew needles after 20 days. CK is the blank control. The black spots in the mid-region of the needles are puncture wounds made by sterilised needles.

In the present study, the two species were distinct from other Pestalotioid species found in Chinese yew (Chen et al. 2002; Ding et al. 2009; Liu et al. 2010; Ismail et al. 2013; Li 2017; Subban et al. 2019; Qi et al. 2022; Sun et al. 2023). Morphologically, the conidia of *P. taxicola* and *P. multicolor* are smaller ($16.5\text{--}21 \times 4\text{--}6 \mu\text{m}$ and $14.5\text{--}20 \times 3.5\text{--}5.5 \mu\text{m}$, respectively). Phylogenetically, the two novel species in the present study were distantly related to four known Pestalotioid species, with the exception of *P. affinis*, which was known only from its morphological descriptions; there were no DNA-based sequence data to compare the phylogenetic relationship with our novel species (Chaiwan et al. 2020).

Pestalotioid fungi represent a diverse group of more than 593 taxa (<http://www.indexfungorum.org/>). However, the taxonomy of the genera and species in this group remains unclear (Maharachchikumbura et al. 2014). In recent years, multi-gene phylogenetic analyses (Combined ITS, *tub2* and *tef-1a* sequence data) have been increasingly utilised to distinguish taxa in the group (Maharachchikumbura et al. 2012). In this study, the low statistical support for some species in the phylogenetic tree of *Neopestalotiopsis* may be due to the absence of *tub2* and *tef-1a* data in some taxa (Yang et al. 2021). The interspecific relationships in the genus *Neopestalotiopsis* are ambiguous (Li et al. 2021b), as some clades are not unambiguously strongly supported, indicating that the relationships amongst different species should be revised and re-evaluated. We decided against describing *Neopestalotiopsis* sp. 3 and sp. 4 as novel species in the genus. These two potentially novel species may be synonymised with existing species during revision and/or monography. Additional examination of archived specimens and sequence data is required for a better resolution and circumscription of *Neopestalotiopsis*.

In the present study, *P. trachycarpicola* was isolated from diseased needles. This is the first report of *P. trachycarpicola* isolated from yews in China. Under the trial conditions, no symptom development occurred in any of the inoculated needles, suggesting that *P. trachycarpicola* may behave as an endophyte in Chinese yew. *P. trachycarpicola* was first described in *Trachycarpus fortunei* in the Yunnan Province of China, causing leaf spot (Zhang et al. 2012) and has already been isolated from a wide range of hosts and locations worldwide (Fan et al. 2021; Khalilabad and Fotouhifar 2022; Araujo et al. 2023; Lan et al. 2023). For example, this fungus has been identified as the causal agent of leaf spots on *Sorghum bicolor* and *Panax notoginseng* in China (Fan et al. 2021; Lan et al. 2023), *Codiaeum variegatum* in Iran (Khalilabad and Fotouhifar 2022) and blueberry in Brazil (Araujo et al. 2023).

In the present study, various *Neopestalotiopsis* and *Pestalotiopsis* species were isolated from infected Chinese yew samples with similar symptoms. Moreover, pathogenicity tests showed that only *Neopestalotiopsis* sp. 3 was pathogenic to Chinese yew, causing dark brown lesions on wounded needles, suggesting that *Neopestalotiopsis* sp. 3 is responsible for needle-spot disease in Chinese yew. The four other Pestalotioid fungi that do not produce typical disease symptoms may be due to their relatively low virulence or endophytes (Reddy et al. 2016; Ma et al. 2019). Alternatively, the development of needle spots disease is the result of more than one biotic or abiotic factor. Different studies have indicated that some *Neopestalotiopsis* and *Pestalotiopsis* species have been isolated from healthy and dead tissues and reported as endophytic fungi. Several *Pestalotiopsis* species have been isolated as endophytes, but have been reported as pathogens in various hosts and in different regions (Moslemi and Taylor 2015; Nozawa et al. 2019; Wang et al. 2019; Chen et al.

2021; Jiang et al. 2021). This suggests that some species may have interchangeable lifestyles such as endophytes, pathogens and saprobes within the same host. Understanding the external factors that influence fungal lifestyles can have major implications for plant health (Sun et al. 2023). Furthermore, the inoculation of wounded needles with mycelial plugs and spore suspensions of *Neopestalotiopsis* sp. 3 was found to be ineffective against disease infection and development. In future studies, the inoculum form and field conditions, such as relative humidity and temperature, should be considered. Furthermore, *Neopestalotiopsis* sp. 3 grew faster than other Pestalotioid species on all three carbon sources tested ($p < 0.01$), possibly contributing to its greater virulence than the others, but its growth on dextrose was not significantly different from that on the other two carbon sources (sucrose and maltose). In the present study, our results showed that temperature affected mycelial growth and *Neopestalotiopsis* sp. 3 grew well at 25–28 °C. The average monthly temperature between May and September is 25.6 °C and 26.8 °C, respectively, at our study site (<http://www.weather.com.cn/>). This is consistent with the most severe epidemic incidence and events observed during field surveillance and observation. Therefore, the culture characteristics of pathogenic fungi can be utilised to forecast and model disease spread.

In conclusion, the present study illustrates the diversity of Pestalotioid species associated with needle spot disease in Chinese yew. Understanding the taxonomy, biology and pathogenicity of Pestalotioid species associated with the Chinese yew will provide a foundation for monitoring disease development and provide information for management strategies for these pathogens.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by National Key R&D Program of China (No. 2021YFC2600400), The Fundamental Research Funds for the Central Universities, China (No. 2021ZY12) and National Natural Science Foundation of China (No. 31770683).

Author contributions

All authors contributed extensively to the work presented in this paper. Conceptualisation: You CJ; Investigation: Wang YF; Data analysis: Chen SM; Writing-original draft: Wang YF; Writing & editing: You CJ & Tsui KM.

Author ORCIDs

Yifeng Wang  <https://orcid.org/0009-0006-6533-6615>

Kin-Ming Tsui  <https://orcid.org/0000-0001-5129-1037>

Shimei Chen  <https://orcid.org/0009-0000-9862-9453>

Chongjuan You  <https://orcid.org/0000-0001-6130-5703>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Alves A, Crous PW, Correia A, Phillips AJL (2008) Morphological and molecular data reveal cryptic speciation in *Lasiodiplodia theobromae*. *Fungal Diversity* 28: 1–13. <https://doi.org/10.1002/yea.1554>
- Araujo L, Ferreira Pinto FAM, de Andrade CCL, Mituti T, Falkenbach BR, Gomes LB, Duarte V (2023) *Pestalotiopsis trachycarpicola* causes leaf spot disease on blueberry in Santa Catarina, Brazil. *Australasian Plant Disease Notes, Australasian Plant Pathology Society* 18(1): 14. <https://doi.org/10.1007/s13314-023-00500-7>
- Chaiwan N, Wanasinghe DN, Mapook A, Jayawardena RS, Norphanphoun C, Hyde KD (2020) Novel species of *Pestalotiopsis* fungi on *Dracaena* from Thailand. *Mycology* 11(04): 306–315. <https://doi.org/10.1080/21501203.2020.1801873>
- Chen Y, Wei G, Chen W (2002) New species of *Pestalotiopsis*. *Mycosystema* 21(03): 316–323.
- Chen Y, Wan Y, Zeng L, Meng Q, Yuan L, Tong H (2021) Characterization of *Pestalotiopsis chamaeropsis* causing gray blight disease on tea leaves (*Camellia sinensis*) in Chongqing, China. *Canadian Journal of Plant Pathology* 43(3): 413–420. <https://doi.org/10.1080/07060661.2020.1816582>
- Ding G, Zheng Z, Liu S, Zhang H, Guo L, Che Y (2009) Photinides A-F, cytotoxic benzofuranone-derived γ -Lactones from the plant endophytic fungus *Pestalotiopsis photiniae*. *Journal of Natural Products* 72(5): 942–945. <https://doi.org/10.1021/np900084d>
- Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus (San Francisco, Calif.)* 12: 13–15.
- Fan X, Zhu J, Zhou X, Meng X, Lin Z, Guo W (2006) The identification and biological characteristics of the anthracnose pathogen of Southern Chinese yew. *Journal-Fujian College Forestry* 26(2): 117–122.
- Fan J, Qiu HB, Long HJ, Zhao W, Xiang XL, Hu AL (2021) First report of leaf spot on sorghum caused by *Pestalotiopsis trachycarpicola* in China. *Journal of Plant Pathology* 103(3): 1043–1044. <https://doi.org/10.1007/s42161-021-00853-x>
- Freitas EFS, Silva MD, Barros MVP, Kasuya MCM (2019) *Neopestalotiopsis hadrolaeliae* sp. nov., a new endophytic species from the roots of the endangered orchid *Hadrolaelia jongheana* in Brazil. *Phytotaxa* 416(3): 211–220. <https://doi.org/10.11646/phytotaxa.416.3.2>
- Gao ZW (2006) Research on *Taxus chinensis* var. *Mairei*. China Forestry Publishing House, China, 143 pp.
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61(4): 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Ismail AM, Cirvilleri G, Polizzi G (2013) Characterisation and pathogenicity of *Pestalotiopsis uvicola* and *Pestalotiopsis clavispora* causing grey leaf spot of mango (*Mangifera indica* L.) in Italy. *European Journal of Plant Pathology* 135(4): 619–625. <https://doi.org/10.1007/s10658-012-0117-z>
- Jeon YH, Cheon W (2014) First report of leaf blight of Japanese yew caused by *Pestalotiopsis microspora* in Korea. *Plant Disease* 98(05): 691–691. <https://doi.org/10.1094/PDIS-08-13-0821-PDN>

- Jiang N, Fan X, Tian C (2021) Identification and characterization of leaf-inhabiting fungi from *Castanea* plantations in China. *Journal of Fungi* 7(1): 64. <https://doi.org/10.3390/jof7010064>
- Jiang N, Voglmayr H, Xue H, Piao CG, Li Y (2022) Morphology and phylogeny of *Pestalotiopsis* (Sporocadaceae, Amphisphaeriales) from Fagaceae leaves in China. *Microbiology Spectrum* 10(6): e03272–e22. <https://doi.org/10.1128/spectrum.03272-22>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Keith RC, Keith LM, Hernandez-Guzman G, Uppalapati SR, Bender CL (2003) Alginate gene expression by *Pseudomonas syringae* pv. tomato DC3000 in host and non-host plants. *Microbiology (Reading, England)* 149(5): 1127–1138. <https://doi.org/10.1099/mic.0.26109-0>
- Khalilabad AA, Fotouhifar KB (2022) Morphological and molecular characterization of a novel *Pestalotiopsis trachycarpicola*, causing Garden Croton leaf spot in Iran. *Mycologia Iranica* 9(1): 59–66.
- Kumaran RS, Kim HJ, Hur BK (2010) Taxol promising fungal endophyte, *Pestalotiopsis* species isolated from *Taxus cuspidata*. *Journal of Bioscience and Bioengineering* 110(5): 541–546. <https://doi.org/10.1016/j.jbiosc.2010.06.007>
- Lan CZ, Lin X, Dai YL, Gan L, Liu XF, Ruan HC, Yang XJ (2023) First report of leaf blight on *Panax notoginseng* caused by *Pestalotiopsis trachycarpicola* in China. *Journal of Plant Pathology* 105(1): 321–321. <https://doi.org/10.1007/s42161-022-01236-6>
- Li Y (2017) Identification of the leaf spot pathogen on Chinese yew and screening of fungicides to *Pestalotiopsis microspora*. PhD Thesis, Fujian Agriculture and Forestry University, Fujian, China.
- Li C, Huo C, Zhang M, Shi Q (2008) Chemistry of Chinese yew, *Taxus chinensis* var. *mairei*. *Biochemical Systematics and Ecology* 36(4): 266–282. <https://doi.org/10.1016/j.bse.2007.08.002>
- Li K, Zhang G, Zhang Y, Griffith MP (2021a) A noteworthy case of rewilding Chinese yew from a garden population in eastern China. *PeerJ* 9: e12341. <https://doi.org/10.7717/peerj.12341>
- Li L, Yang Q, Li H (2021b) Morphology, phylogeny, and pathogenicity of Pestalotioid species on *Camellia oleifera* in China. *Journal of Fungi (Basel, Switzerland)* 7(12): 1080. <https://doi.org/10.3390/jof7121080>
- Liu AR, Chen SC, Wu SY, Xu T, Guo LD, Jeewon R, Wei JG (2010) Cultural studies coupled with DNA based sequence analyses and its implication on pigmentation as a phylogenetic marker in *Pestalotiopsis* taxonomy. *Molecular Phylogenetics and Evolution* 57(2): 528–535. <https://doi.org/10.1016/j.ympev.2010.07.017>
- Liu L, Wang Z, Huang L, Wang T, Su Y (2019) Chloroplast population genetics reveals low levels of genetic variation and conformation to the central-marginal hypothesis in *Taxus wallichiana* var. *mairei*, an endangered conifer endemic to China. *Ecology and Evolution* 9(20): 11944–11956. <https://doi.org/10.1002/ece3.5703>
- Ma XY, Maharachchikumbura SSN, Chen BW, Hyde KD, McKenzie EHC, Chomnunti P, Kang JC (2019) Endophytic pestalotioid taxa in *Dendrobium* orchids. *Phytotaxa* 419(3): 268–286. <https://doi.org/10.11646/phytotaxa.419.3.2>
- Maharachchikumbura SSN, Guo LD, Cai L, Chukeatirote E, Wu WP, Sun X, Crous PW, Bhat DJ, McKenzie EHC, Bahkali A, Hyde KD (2012) A multi-locus backbone tree for *Pestalotiopsis*, with a polyphasic characterization of 14 new species. *Fungal Diversity* 56(1): 95–129. <https://doi.org/10.1007/s13225-012-0198-1>

- Maharachchikumbura SSN, Hyde KD, Groenewald JZ, Xu J, Crous PW (2014) *Pestalotiopsis* revisited. *Studies in Mycology* 79(1): 121–186. <https://doi.org/10.1016/j.simyco.2014.09.005>
- Moslemi A, Taylor PWJ (2015) *Pestalotiopsis chamaeropsis* causing leaf spot disease of round leaf mint-bush (*Prostanthera rotundifolia*) in Australia. *Australasian Plant Disease Notes, Australasian Plant Pathology Society* 10(1): 1–5. <https://doi.org/10.1007/s13314-015-0179-9>
- Nozawa S, Seto Y, Watanabe K (2019) First report of leaf blight caused by *Pestalotiopsis chamaeropsis* and *Neopestalotiopsis* sp. in Japanese andromeda. *Journal of General Plant Pathology* 85(6): 449–452. <https://doi.org/10.1007/s10327-019-00868-4>
- Pyo SH, Park HB, Song BK, Han BH, Kim JH (2004) A large-scale purification of paclitaxel from cell cultures of *Taxus chinensis*. *Process Biochemistry (Barking, London, England)* 39(12): 1985–1991. <https://doi.org/10.1016/j.procbio.2003.09.028>
- Qi XL, He J, Ju Y, Huang L (2022) First report of leaf spot on *Elaeagnus pungens* caused by *Neopestalotiopsis clavisporea* in China. *Plant Disease* 107(7): 2251. <https://doi.org/10.1094/PDIS-10-22-2457-PDN>
- Qian TM, Rong L, Zhu TH (2015) Identification of the pathogen isolated from root rot of *Taxus chinensis* var. *mairei* and screening of its antagonistic *Bacillus*. *Plant Protection* 41(06): 60–66.
- Rambaut A (2016) FigTree v1.4.3. Institute of Evolutionary Biology, University of Edinburgh, UK. <http://tree.bio.ed.ac.uk/software/figtree>
- Ran SF, Maharachchikumbura SSN, Ren YL, Liu H, Chen KR, Wang YX, Wang Y (2017) Two new records in Pestalotiopsidaceae associated with Orchidaceae disease in Guangxi Province, China. *Mycosphere* 8(01): 121–130. <https://doi.org/10.5943/mycosphere/8/1/11>
- Reddy MS, Murali TS, Suryanarayanan TS, Rajulu MG, Thirunavukkarasu N (2016) *Pestalotiopsis* species occur as generalist endophytes in trees of Western Ghats forests of southern India. *Fungal Ecology* 24: 70–75. <https://doi.org/10.1016/j.funeco.2016.09.002>
- Ru WM (2006) Study on the ecology of endangering plant *Taxus chinensis* var. *mairei*. PhD Thesis, Shanxi University, Shanxi, China.
- Rubini MR, Silva-Ribeiro RT, Pomella AW, Maki CS, Araújo WL, Dos Santos DR, Azevedo JL (2005) Diversity of endophytic fungal community of cacao (*Theobroma cacao* L.) and biological control of *Crinipellis pernicioso*, causal agent of Witches' Broom Disease. *International Journal of Biological Sciences* 1: 24–33. <https://doi.org/10.7150/ijbs.1.24>
- Shu J, Yu Z, Sun W, Zhao J, Li Q, Tang L, Guo T, Huang S, Mo J, Hsiang T, Luo S (2020) Identification and characterization of pestalotioid fungi causing leaf spots on mango in southern China. *Plant Disease* 104(4): 1207–1213. <https://doi.org/10.1094/PDIS-03-19-0438-RE>
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Strobel G, Yang X, Sears J, Kramer R, Sidhu RS, Hess WM (1996) Taxol from *Pestalotiopsis microspora*, an endophytic fungus of *Taxus wallichiana*. *Microbiology (Reading, England)* 142(2): 435–440. <https://doi.org/10.1099/13500872-142-2-435>
- Subban K, Subramani R, Srinivasan VPM, Johnpaul M, Chelliah J (2019) Salicylic acid as an effective elicitor for improved taxol production in endophytic fungus *Pestalotiopsis microspora*. *PLOS ONE* 14(2): e0212736. <https://doi.org/10.1371/journal.pone.0212736>
- Sun YR, Jayawardena RS, Sun JE, Wang Y (2023) Pestalotioid species associated with medicinal plants in southwest China and Thailand. *Microbiology Spectrum* 11(1): e03987–e22. <https://doi.org/10.1128/spectrum.03987-22>

- Swofford D (2003) PAUP* - Phylogenetic analysis using parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, Massachusetts, USA.
- Wang R, Chen S, Zheng B, Liu P, Li B, Weng Q, Chen Q (2019) Occurrence of leaf spot disease caused by *Neopestalotiopsis clavispora* on *Taxus chinensis* in China. *Forest Pathology* 49(5): e12540. <https://doi.org/10.1111/efp.12540>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR-Protocols: A guide to methods and applications* 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wu L, Han T, Li W, Jia M, Xue L, Rahman K, Qin L (2013) Geographic and tissue influences on endophytic fungal communities of *Taxus chinensis* var. *mairei* in China. *Current Microbiology* 66(1): 40–48. <https://doi.org/10.1007/s00284-012-0235-z>
- Xiong Y, Manawasinghe I, Maharachchikumbura SSN, Dong ZY, Xiang MM, Li L (2022) Pestalotioid species associated with palm species from Southern China. *Current Research in Environmental & Applied Mycology* 12(1): 285–321. <https://doi.org/10.5943/cream/12/1/18>
- Yang Q, Zeng XY, Yuan J, Zhang Q, He YK, Wang Y (2021) Two new species of *Neopestalotiopsis* from southern China. *Biodiversity Data Journal* 9: e70446. <https://doi.org/10.3897/BDJ.9.e70446>
- Zhang Y, Maharachchikumbura SS, Mckenzie EH, Hyde KD (2012) A novel species of *Pestalotiopsis* causing leaf spots of *Trachycarpus fortunei*. *Cryptogamie. Mycologie* 33(3): 311–318. <https://doi.org/10.7872/crym.v33.iss3.2012.311>
- Zhang GF, Li Q, Hou XR (2018) Structural diversity of naturally regenerating Chinese yew (*Taxus wallichiana* var. *mairei*) populations in ex situ conservation. *Nordic Journal of Botany* 36(4): njb–01717. <https://doi.org/10.1111/njb.01717>
- Zhang Z, Liu R, Liu S, Mu T, Zhang X, Xia J (2022) Morphological and phylogenetic analyses reveal two new species of Sporocadaceae from Hainan, China. *MycKeys* 88: 171–192. <https://doi.org/10.3897/mycokeys.88.82229>

Supplementary material 1

PCR primer and PCR reaction conditions

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: xlsx

Explanation note: **table S1**. PCR primer and PCR reaction conditions.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl1>

Supplementary material 2

Specimen information and GenBank accession numbers

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: xlsx

Explanation note: **table S2**. Isolates and GenBank accession numbers of sequences used in this study (the isolates from this study are marked in red).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl2>

Supplementary material 3

Evolutionary divergence between sequences

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: xls

Explanation note: **table S3**. Estimates of Evolutionary Divergence between Sequences.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl3>

Supplementary material 4

Effects of light on growth of isolates

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: jpg

Explanation note: **fig. S1**. Effects of light on growth of isolates.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl4>

Supplementary material 5

Effects of carbon sources on growth of isolates

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: jpg

Explanation note: **fig. S2**. Effects of carbon sources on growth of isolates.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl5>

Supplementary material 6

Nucleotide differences between the isolates in this study

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: pdf

Explanation note: documentation S1: Nucleotide differences between the isolates in this study.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl6>

Supplementary material 7

Sequence data of the isolates in this study

Authors: Yifeng Wang, Kin-Ming Tsui, Shimei Chen, Chongjuan You

Data type: zip

Explanation note: zip package S1: Sequence data of the isolates in this study.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.102.113696.suppl7>