

More pieces to a huge puzzle: Two new *Escovopsis* species from fungus gardens of attine ants

Quimi Vidaurre Montoya¹, Maria Jesus Sutta Martiarena¹,
Danilo Augusto Polezel¹, Sérgio Kakazu², Andre Rodrigues^{1,2}

1 Department of Biochemistry and Microbiology, UNESP – São Paulo State University, Rio Claro, SP, Brazil

2 Center for the Studies of Social Insects, UNESP – São Paulo State University, Rio Claro, SP, Brazil

Corresponding author: Andre Rodrigues (andrer@rc.unesp.br)

Academic editor: T. Lumbsch | Received 31 October 2018 | Accepted 16 January 2019 | Published 18 February 2019

Citation: Montoya QV, Martiarena MJS, Polezel DA, Kakazu S, Rodrigues A (2019) More pieces to a huge puzzle: Two new *Escovopsis* species from fungus gardens of attine ants. MycoKeys 46: 97–118. <https://doi.org/10.3897/mycokeys.46.30951>

Abstract

Escovopsis (Ascomycota: Hypocreales, Hypocreaceae) is the only known parasite of the mutualistic fungi cultivated by fungus-growing ants (Formicidae: Myrmicinae: Attini: Attina, the “attines”). Despite its ecological role, the taxonomy and systematics of *Escovopsis* have been poorly addressed. Here, based on morphological and phylogenetic analyses with three molecular markers (internal transcribed spacer, large subunit ribosomal RNA and the translation elongation factor 1-alpha), we describe *Escovopsis clavatus* and *E. multiformis* as new species isolated from fungus gardens of *Apterostigma* ant species. Our analysis shows that *E. clavatus* and *E. multiformis* belong to the most derived *Escovopsis* clade, whose main character is the presence of conidiophores with vesicles. Nevertheless, the most outstanding feature of both new species is the presence of a swollen region in the central hypha of the conidiophore named swollen cell, which is absent in all previously described *Escovopsis* species. The less derived *Escovopsis* clades lack vesicles and their phylogenetic position within the Hypocreaceae still remains unclear. Considering the high genetic diversity in *Escovopsis*, the description of these new species adds barely two pieces to a huge taxonomic puzzle; however, this discovery is an important piece for building the systematics of this group of fungi.

Keywords

Hypocreales, Taxonomy, Phylogeny, Parasitic fungi, Symbiosis

Introduction

Microorganisms play important roles in the stability of social insect colonies (Hughes et al. 2008, Joop and Vilcinskis 2016, Vanderpool et al. 2018). The environment of these insects has a high potential to harbour unique fungal species (Attili-Angelis et al. 2014, Harrington et al. 2014, Menezes et al. 2015, Montoya et al. 2016). The evolutionary success of the fungus garden of the fungus-farming ants (Formicidae: Myrmicinae: Attini: *Attina*, the “attines”) depends on complex symbiotic interactions amongst bacteria, fungi and the ants (Currie et al. 2003, Gerardo et al. 2006a, Kost et al. 2007). The association between attine ants and their mutualistic fungi (Basidiomycota: Agaricales) is the core of the attine colonies; however, *Escovopsis* (Ascomycota: Hypocreales: Hypocreaceae) can exploit this association. Although no specialised parasitic structures were found, studies showed that this parasite is able to kill the fungal cultivar as well as the ants and their mutualistic bacteria by chemical compounds (Currie 2001, Varanda-Haifig et al. 2017, Dhodary et al. 2018, Heine et al. 2018, Custodio and Rodrigues 2019). Despite the ecological relevance of *Escovopsis* as parasites of attine ant colonies, the taxonomy of this genus has been neglected.

Attine ants are classified in two sister clades: the Palaeoattina and Neoattina (Branstetter et al. 2017). Leafcutter ants (*Atta* and *Acromyrmex*) are considered the most derived attines within the Neoattina. Their behaviour is characterised by collecting fresh leaves and flowers to feed several cultivars from two clades of fungi in the Agaricaceae (Mueller et al. 2017, 2018). On the other hand, non-leafcutter ants also occur in both the Neoattina and Palaeoattina clades. Distinct from *Atta* and *Acromyrmex*, non-leafcutter ants collect seeds, insect frass and dry leaves to nourish a wide range of fungal cultivars in the Agaricaceae and Pterulaceae (Villesen et al. 2004, Schultz and Brady 2008).

The attine ant-fungus cultivar-*Escovopsis* symbiosis has been widely studied in leafcutter ants (Mueller and Gerardo 2002, Currie et al. 2003, Gerardo et al. 2004, 2006a,b). In addition to their contributions on the biology of *Escovopsis*, these studies also revealed considerable diversity of the parasite. Considering the variety of mutualistic fungi that non-leafcutter ants may cultivate, as well as the different substrates used for that purpose, a high diversity of *Escovopsis* species is unsurprising. This is especially true for *Apterostigma* (Gerardo et al. 2006b), a genus of non-leafcutter attine with species that cultivate different cultivars including *Leucoagaricus gongylophorus*, the domesticated fungus cultivated by many higher attine ant species, mostly leafcutter ants (Sosa-Calvo and Schultz 2010, Schultz et al. 2015, Ješovnik et al. 2016, Sosa-Calvo et al. 2017, Mueller et al. 2017, 2018).

While *Escovopsis* species exploiting gardens of *Atta*, *Acromyrmex*, *Trachymyrmex* and *Mycetophylax* were formally described, the morphological characters of the species associated with *Apterostigma* are unknown. A previous study associated clades of the parasite with the colour pattern of *Escovopsis* colonies (brown, yellow, white and pink; Gerardo et al. 2006b). However, no taxonomic studies were undertaken to formally describe these clades. Here, we describe *Escovopsis clavatus* and *E. multiformis* as new species isolated from the fungus garden of *Apterostigma*. The distinctive feature of

these lineages is the presence of swollen cells at the base of the conidiophore branches. This phenotype differentiates these two new species from previously described *Escovopsis*. Considering that previous studies showed a high genetic diversity within *Escovopsis*, the description of these species adds two pieces to the enormous taxonomic puzzle which is *Escovopsis*.

Material and methods

Sampling sites and *Escovopsis* isolation

Five *Escovopsis* isolates were obtained from fungus gardens of five different colonies of *Apterostigma* spp. (Suppl. material 1: Table S1). The isolates LESF 847, LESF 853, LESF 854 and LESF 855 were obtained from colonies found in the Atlantic Rain Forest in Florianópolis, State of Santa Catarina, Brazil (October 2015). The isolate LESF 1136 was obtained from a colony found in the Amazon Forest in Cotriguaçu, State of Mato Grosso, Brazil (October 2017). The nests were found inside or under rotten logs. Fungus gardens, along with tending workers and brood, were collected in UV-sterilised plastic containers using sterilised spoon and forceps. Samples were taken to the Laboratory of Fungal Ecology and Systematics (LESF) at the UNESP – São Paulo State University, Rio Claro, Brazil.

For fungal isolation, seven garden fragments (0.5–1 mm³) were inoculated on plates (three plates per colony) containing potato dextrose agar (PDA, Neogen Culture Media, Neogen) supplemented with chloramphenicol (150 µg mL⁻¹, Sigma) and incubated at 25 °C in darkness. Plates were monitored daily for fungal growth and, when *Escovopsis* mycelia sprouted, they were transferred to new PDA plates. All isolates were prepared as axenic (monosporic) cultures and stored under sterile distilled water kept at 8 °C (Castellani 1963) and at –80 °C (as conidia suspensions in 10% glycerol).

Morphological analysis

The morphological characters of the five isolates (LESF 847, LESF 853, LESF 854, LESF 855 and LESF 1136) were examined. Due to the lack of standardisation of culture conditions for *Escovopsis*, the macroscopic characters of the colonies, i.e. radial growth, mycelium colour, morphology and presence of soluble pigments, were evaluated on eight different media: PDA, malt agar 2% [MA2%: 20 g L⁻¹ of malt extract (Neogen Culture Media) and 15 g L⁻¹ of agar (Neogen Culture Media)], cornmeal agar (CMD, Neogen Culture Media), synthetic nutrient agar [SNA: 1 g L⁻¹ of KH₂PO₄ (Labsynth), 1 g L⁻¹ of KNO₃ (Labsynth), 0.5 g L⁻¹ of MgSO₄(7H₂O) (Labsynth), 0.5 g L⁻¹ of KCl (Labsynth), 0.2 g L⁻¹ of Glucose (Labsynth), 0.2 g L⁻¹ of Sucrose (Labsynth) and 15 g L⁻¹ of Agar (Neogen Culture Media)], oatmeal agar (OA), potato carrot agar (PCA, HiMedia), malt extract agar 2% [MEA: 30 g L⁻¹ of malt extract (Neogen Cul-

ture Media), 5 g L⁻¹ of bacteriological peptone (Neogen Culture Media), 20 g L⁻¹ of glucose (Labsynth) and 15 g L⁻¹ of Agar (Neogen Culture Media)] and Czapek yeast extract agar [CYA; 30 g L⁻¹ of Sucrose (Labsynth), 5 g L⁻¹ of Yeast extract (Neogen Culture Media), 1 g L⁻¹ of KH₂PO₄ (Labsynth), 0.3 g L⁻¹ of NaNO₃ (Synth), 0.05 g L⁻¹ of KCl (Labsynth), 0.05 g L⁻¹ of MgSO₄(7H₂O) (Labsynth), 0.001 g L⁻¹ of FeSO₄ (Labsynth), 0.001 g L⁻¹ of ZnSO₄ (Labsynth), 0.0005 g L⁻¹ of CuSO₄ (Labsynth), 15 g L⁻¹ of Agar (Neogen Culture Media)] at five temperatures (10 °C, 20 °C, 25 °C, 30 °C and 35 °C). These temperatures correspond to the conditions used in previous studies that described *Escovopsis* species (Seifert et al. 1995, Augustin et al. 2013, Masiulionis et al. 2015, Meirelles et al. 2015a). For this purpose, 200 µl of conidia were spread on plates with water-agar (WA) and incubated for seven days at 25 °C in darkness. Then, mycelium fragments of 0.5 cm diameter were cut from the WA plates and inoculated in the centre of the plates (90 × 15 mm) containing the eight culture media. All the strains examined showed better development in the dark and with unsealed Petri dishes to allow air passage; therefore, incubation was carried out in the darkness and without sealing the plates, for 14 days. Three replicate plates were inoculated for each media and for each incubation temperature.

To examine the microscopic characters, i.e. the morphology, size, branching patterns, vesicles and swollen cells of the conidiophores, as well as phialides and conidia, slide cultures on PDA and MEA were performed. Briefly, we placed a 5 mm² fragment of culture medium on a microscopic slide and then we inoculated the fungus at the centre of the fragment. Then, the inoculated medium was covered with a coverslip and incubated at 25 °C for 4–7 days in the dark. After that, the coverslips, where the fungus grew, were removed and placed in new slides with a drop of lactophenol. Finally, the slides were examined under a light microscope (DM750, Leica, Germany). Fungal microscopic structures were photographed and measured (with 30 measurements per structure) in LAS EZ v.4.0 (Leica Application Suite).

Microscopic structures were also examined under scanning electron microscopy (SEM). Fungal samples (five days old cultures on PDA) were fixed in osmium tetroxide vapour for 72 h. Then, samples were dehydrated using a series of acetone concentrations (50, 75, 90, 95 and 100%) and dried to critical point using liquid CO₂ (Balzers CPD030). The dried material was sputtered with gold (Balzers SCD050) and examined under the scanning electron microscope (TM3000, Hitachi).

DNA extraction, PCR and sequencing

DNA extraction of the five strains was performed, following the steps published in Meirelles et al. (2015a). Three molecular markers were amplified: the internal transcribed spacer (ITS) region (White et al. 1990, Schoch et al. 2012); translation elongation factor 1-alpha (*tef1*) (Taerum et al. 2007); and the large subunit ribosomal RNA (LSU) (White et al. 1990, Haugland and Heckman 1998, Currie et al. 2003) (Suppl. material 1: Table S2).

PCR and sequence reaction conditions followed the steps published in Meirelles et al. (2015b) for the ITS region, Meirelles et al. (2015a) for *tefl* and Augustin et al. (2013) for LSU. The final amplicons were cleaned up with Wizard SV Gel and PCR Clean-up System kit (Promega), following the manufacturer's protocol. Sequences (forward and reverse) were generated in ABI3500 (Life Technologies). The LSU of 29 strains previously used in Meirelles et al. (2015b) was also amplified and sequenced for this study (Suppl. material 1: Table S1). The sequences were assembled in contigs in BioEdit v. 7.1.3 (Hall 1999) and deposited in GenBank (Suppl. material 1: Table S1 for accession numbers).

Phylogenetic analyses

To infer the phylogenetic position of the new species in the *Escovopsis* clade, sequences from previous studies were retrieved from the GenBank and aligned with our new sequences in a dataset for each marker (Chaverri et al. 2003, Spatafora et al. 2007, Jaklitsch et al. 2011, Pöldmaa 2011, Meirelles et al. 2015b). This data included sequences from the seven *Escovopsis* ex-type strains, from *Escovopsioides nivea* and some species from *Hypomyces* and *Trichoderma*, as the phylogenetic closest relatives of *Escovopsis*. First, the three datasets [46 sequences of ITS (619 bp), LSU (594 bp) and *tefl* (758 bp)] were aligned separately in MAFFT v.7 (Kato and Standley 2013). The end parts of each alignment were removed manually by considering a point where the sequences presented greater homogeneity (all alignments are deposited in Treebase: <http://purl.org/phylo/treebase/phyloids/study/TB2:S23689>). Then, a phylogenetic tree was inferred using each dataset separately. The nucleotide substitution model was selected by independent runs in jModelTest 2 (Darriba et al. 2012) using the Akaike Information Criterion with a 95% confidence interval. Second, the three datasets were concatenated using Winclada v.1.00.08 (Nixon 2002). The final file comprised 46 sequences totalling 1971 bp. All phylogenetic trees were reconstructed using maximum likelihood (ML) in RAxML v.8 (Stamatakis 2014) with 1000 independent trees and 1000 bootstrap replicates (MLB) and Bayesian Inference (BI) in MrBayes v.3.2.2. (Ronquist et al. 2012). The ML phylogenetic trees were reconstructed using the GTR + G substitution model and the BI phylogenetic trees were performed with the GTR + I + G substitution model. In the case of BI, two separate runs were carried out, each consisting of three hot chains and one cold chain and a Markov Chain Monte Carlo (MCMC) sampling for two million generations to obtain Bayesian posterior probability (PP) values for the clades. Convergence occurred when the standard deviation of split frequencies fell below 0.01 and the first 25% of the generations of MCMC sampling were discarded as burn-in. The final phylogenetic trees were edited in FigTree v.1.4 and in Adobe Illustrator CC v.17.1. *Lecanicillium antillanum* CBS 350.85 was used as the outgroup in all trees, because it belongs to a family phylogenetically close to Hypocreaceae (Spatafora et al. 2007).

Results

Taxonomy

Escovopsis clavatus Q.V. Montoya, M.J.S. Martiarena, D.A. Polezel, S. Kakazu & A. Rodrigues, sp. nov.

MycoBank: MB828328

Figs 1–3

Etymology. “*clavatus*” in reference to the predominantly clavate shape of vesicles.

Typification. BRAZIL. Santa Catarina, Florianópolis, (27°44'39.6"S, 48°31'10.14"W), elev. 46 m, fungus garden, 08, 2015. *A. Rodrigues*. Holotype: CBS H-23845 (dried culture on PDA). Ex-type strain LESF 853 (= CBS 145326).

Sequences. ITS (MH715096), *tef1* (MH724270) and LSU (MH715110).

Description. Colonies grow only at 20 and 25 °C (Fig. 1). At both temperatures, growth starts on the third day on CMD, CYA, MA2%, MEA, OA, PCA, PDA; and on the sixth day on SNA. Colonies have floccose aerial mycelia with a pale-brown colour after seven days. Faster growth was observed on MA2% and heavy sporulation was identified on MA2%, PDA and OA. At 20 °C, colonies reached 0.5–0.7 cm, 1.5–2.5 cm and 0.5–1 cm on CMD, CYA and SNA, respectively. At this temperature, colonies reached the edge of the plate after 10 days on MA2% and PCA; after 12 days on OA and MEA; and after 14 days on PDA and CYA. At 25 °C, colonies reach 2 cm, 3–3.2 cm and 2 cm on CMD, CYA and SNA, respectively, after 14 days. At this temperature, colonies reached the plate edge after seven days on OA and PCA; and after 10 days on MA2%, MEA and PDA. Concentric rings were observed only on PCA at 20 °C (Fig. 1). No pustule-like structures were observed.

Conidiophores arising from aerial hypha alternated or opposite (Fig. 2A), with the main axis of 50–780 µm in length, some without branching and often with 1–2 levels of branching (Figs 2A, E, 3A, E). Branches arise from the main axis of the conidiophore in an alternated or opposite pattern, with a septum near to the central axis and before the vesicle, usually with 1–2 branches at each branching point (16–138 µm long) or 2–4 branches arising from swollen cells (28–35 µm long), mostly forming angles less than 90° and less frequently right angles, usually straight and sometimes slightly curved up or down. Each branch terminates in a vesicle, with 1–8 fertile heads per conidiophore. Swollen cells are present in 15% of the total of conidiophores examined (Figs 2C, D, 3E) and can measure 10–18 µm long × 7–9 µm wide. *Vesicles* with only a septum at the base, in various shapes: globose (8%), subglobose (24%), broadly ellipsoidal / clavate (33%), ellipsoidal (27%), cylindrical (8%) (Figs 2E–G and 3F–G); and reaching 9–27 µm long × 7–20 µm wide. *Phialides* lageniform formed on vesicles (Fig. 3H), with 5–8 µm in total length, elongated base (0.5–1.5 µm × 0.5–1 µm), followed by a swollen section (1.5–2.5 µm × 1–3 µm) and a thin neck (1.5–4 µm × 0.5 µm). *Conidia* with 1.5 µm–2.5 µm long × 0.5 µm–1.5 µm wide, in various shapes: broadly ellipsoidal (5%), ellipsoidal (43.3%), cylindrical (51.7%); brown, with smooth and slightly thickened walls and in chains (Figs 2H, 3I).




Figure 1. Colony macroscopic characters of *Escovopsis clavatus* and *Escovopsis multiformis* on CMD, CYA, MA2%, MEA, OA, PCA, PDA and SNA media after 14 days at 10, 20, 25 and 30 °C.

Habitat. Isolated from fungus gardens of *Apterostigma* sp.

Additional specimens examined. BRAZIL. Santa Catarina, Florianópolis, (27°44'38.94"S, 48°31'9.3"W), elev. 32 m, fungus garden, 08, 2015. *A. Rodrigues*. LESF 854 (ITS – MH715097, *tef1* – MH724271 and LSU – MH715111). Santa Catarina, Florianópolis, (27°44'39.49"S, 48°31'9.72"W), elev. 38 m, fungus garden, 08, 2015. *A. Rodrigues*. LESF 855 (ITS – MH71509, *tef1* – MH724272 and LSU – MH715112).




Figure 2. *Escovopsis clavatus*. **A, B** Conidiophores without “swollen cells” **C, D** Conidiophores with “swollen cells” (red arrows) **E–G** Vesicles in various shapes with phialides pattern **G** Conidia.




Figure 3. *Escovopsis clavatus*. SEM images **A–D** Conidiophores without “swollen cells” **E** Conidiophore with “swollen cells” (red arrows) **F, G** Vesicles **H** Phialides **G** Conidia.

Notes. *Escovopsis clavatus* is phylogenetically closely related to *E. multiformis* and its most distinctive characters are its growth temperatures, the conidiophore branching and the swollen cells. It grows at 20 and 25 °C; nevertheless, *E. multiformis* grows at 10, 20, 25 and 30 °C. The conidiophore of *E. clavatus* is larger and more branched than the conidiophore of *E. multiformis*. In addition, the swollen cells of *E. clavatus* are less frequent and shorter than in *E. multiformis*. The character distinguishing *E. clavatus* from other species of *Escovopsis* is the swollen cell on the conidiophores and because it is phylogenetically placed in a distinct clade.

***Escovopsis multiformis* Q.V. Montoya, M.J.S. Martiarena, D.A. Polezel, S. Kakazu & A. Rodrigues, sp. nov.**

Mycobank: MB828329

Figs 1, 4, 5

Etymology. “*multiformis*” in relation to the different vesicle shapes found in the same isolate.

Typification. BRAZIL. Santa Catarina, Florianópolis, (27°28'11.28"S, 48°22'39.48"W), elev. 119 m, fungus garden, 08, 2015. *A. Rodrigues*. Holotype: CBS H-23846 (dried culture on PDA). Ex-type strain LESF 847 (= CBS 145327).

Sequences. ITS (MH715091), *tefl* (MH724265) and LSU (MH715105).

Description. Colonies grow at 10, 20, 25 and 30 °C (Fig. 1). The best growth temperature was 30 °C. At this temperature, colonies reached 1.2–1.4 cm, 2.7–3 cm, 2.6–3 cm, 3.3–3.5 cm, 2.5–2.8 cm, 2.7–2.9 cm and 1.9–2.5 cm in radius on CMD, CYA, MA2%, MEA, OA, PCA and PDA, after 14 days, respectively. Colonies exhibit light-brown floccose mycelia (colony edge usually lighter or white). The colour shades and the character of the aerial mycelium vary on each culture medium (Fig. 1). Colonies present concentric rings with a hardened ring similar to a crust in the centre on CYA (Fig. 1) and the sporulation is more abundant on PCA and PDA. At 20 °C, on CMD, CYA, MA2%, MEA, OA, PCA, PDA and SNA, colonies attained 0.5–0.8 cm, 1.1–2.2 cm, 2–2.5 cm, 2.1–2.3 cm, 2–2.5 cm, 2.8 cm, 1.9–2.4 cm and 0–0.1 cm in radius, respectively. At 25 °C, colonies reached 1 cm, 2.1–2.3 cm, 2–2.4 cm, 2.5–2.6 cm, 2.2–2.7 cm, 2.8–3 cm, 1.8–2 cm and 0.1–0.2 cm in radius on CMD, CYA, MA2%, MEA, OA, PCA, PDA and SNA, respectively. Pustule-like structures were observed on OA and CMD at 20, 25 and 30 °C. At 10 °C, the colony growth was inconspicuous, reaching 0.2–0.3 cm, 0.2–0.4 cm, 0.3 cm, 0.6–0.8 cm, 0.8 cm and 0.3–0.5 cm in radius on CYA, MA2%, MEA, OA, PCA and PDA, respectively, after 14 days. At this temperature, growth started in these culture media after seven days and sporulation occurred only after the 12th day. No growth was observed at 35 °C.

Conidiophores arising from aerial hypha alternated or opposite (Fig. 3A), with the main axis of 41–293 µm in length, some without branching and most of them with one level of branching. Rarely, branches form two levels branching (Figs 4A–C, 5A, B). Branches arise from the main axis of the conidiophore alternated, with a septum near




Figure 4. *Escovopsis multififormis*. **A–C** Conidiophores mono- and polycephalous without “swollen cells” **D–G** Conidiophores mono and polycephalous with “swollen cells” (red arrows) **H, I** Vesicles in various shapes **J** Conidia.




Figure 5. *Escovopsis multiformis*. SEM images **A, B** Conidiophores mono- and polycephalous without “swollen cells” **C–F** Conidiophores mono- and polycephalous with “swollen cells” (red arrows) **G, H** Vesicles **I** Phialides **J** Conidia.

the central axis and before the vesicle, usually with one branch at each branching point (32–84 μm long) or 2–4 branches arising from swollen cells (17–86 μm long), mostly forming right angles, usually slightly curved up. Each branch terminates in a vesicle, with 1–4 fertile heads per conidiophore. Swollen cells are present in 27% of the total of con-

idiophores examined (Figs 4D–G, 5C–F) and can measure 16–34 μm long \times 9–20 μm wide. Sometimes, one swollen cells' branch gives rise to another swollen cell with more branches (Figs 2F, 3C). *Vesicles* with only a septum at the base, in various shapes: globose (22%), subglobose (37%), broadly ellipsoidal (26%), ellipsoidal (10%), cylindrical (5%) (Figs 4H, I, 5G, H); and reaching 12–27 μm \times 9–17 μm wide. *Phialides* lageniform formed on vesicles (Fig. 5I), with 6–10 μm in total length, elongated base (1–2.5 μm \times 0.5–1 μm), followed by a swollen section (2.5–4.5 μm \times 2–3.5 μm) and a thin neck (1–4.5 μm \times 0.5–1 μm). *Conidia* are 2.5–3.5 μm long \times 1.5–2.5 μm wide, in various shapes: globose (2%), subglobose (3%), broadly ellipsoidal (33%), ellipsoidal (47%), cylindrical (15%); brown, with smooth and slightly thickened walls and in chains (Figs 4, 5J).

Habitat. Isolated from fungus garden of *Apterostigma* sp.

Additional specimens examined. BRAZIL. Mato Grosso, Cotriguaçu, (09°49'22.74"S, 58°15'32.04"W), elev. 252 m, fungus garden, 10, 2017. *Q. V. Montoya*. LESF 1136 (ITS – MH715092, *tef1* – MH724266 and LSU – MH715106).

Notes. *Escovopsis multififormis* is closely related to *E. clavatus*. Different from *E. clavatus* that grow at 20 and 25 °C, *E. multififormis* grow at 10, 20, 25 and 30 °C. The optimum growth temperature of *E. multififormis* is 30 °C and that of *E. clavatus* is 25 °C. The conidiophores of *E. multififormis* are smaller and less branched than *E. clavatus* and the swollen cells are more frequent and larger than those found in *E. clavatus*. *E. multififormis* differs from other described species by the presence of conidiophores with a swollen cell, the presence of different vesicles shapes and because it is phylogenetically placed in a distinct clade.

Morphological analyses

The isolates LESF 853 (*Escovopsis clavatus*, Figs 1–3) and LESF 847 (*Escovopsis multififormis*, Figs 1, 4, 5) differed from the seven previously described *Escovopsis* species, mainly in micro-morphological structures. All isolates had white colonies with a floccose appearance on all culture media, but *E. clavatus* had the most floccose colonies. After 5–7 days incubation, the centre of the colonies turned pale brown and, after 7 days, the entire colony gradually turned from white to light brown (not always from the middle to the edge in *E. multififormis*).

Escovopsis multififormis showed growth at wide ranges of temperature (from 10–30 °C); nonetheless, *E. clavatus* showed growth only at 20 and 25 °C (Fig. 1). None of the isolates grew at 35 °C. On all culture media, the best growth was obtained at 25 °C for *E. clavatus* and at 30 °C for *E. multififormis*. In all cases where growth was observed, it started between 24 to 36 hours and sporulation started on the third day.

All strains of both species have a unique type of conidiophore with a swollen cell, from which branches emerge (Figs 2C, D, 3E, 4D–F, 5C–F). These conidiophores were more frequent in *E. multififormis* than *E. clavatus* (27% and 15%, respectively). Mono or polycephalous conidiophores, without the swollen cells, that were described in the other *Escovopsis* species, were also present but with some differences in the size and branching pattern (Figs 2A, B, 3A–D, 4A–C, 5A, B). Conidiophores

with cruciform or opposed branches were rarely observed. On the other hand, the two new species had basipetal and smooth-walled conidia with slightly thickened walls, formed from phialides. No chlamydospores were observed in the aerial or submersed mycelia of any of the strains.

Phylogenetic analyses

Separate phylogenetic analyses with the three molecular markers showed topological differences because of the incongruity placement of the formal described *Escovopsis* species and some strains that form new phylogenetic clades within the genus (Fig. 6). The phylogenetic placement of *E. multiformis* and *E. clavatus* also presented conflicts amongst the three molecular markers; however, the position of each strain that made up both new species was concordant through the three genealogies (PP= 1; MBL= 100%, Fig. 6).

The combined analysis also confirmed *E. multiformis* and *E. clavatus* as two new phylogenetic species in *Escovopsis* (PP= 1; MLB= 100%, Fig. 7) and showed the strain LESF 018 (a vesiculated *Escovopsis* species) as the closest relative of both. Nevertheless, the concatenated BI and ML trees also presented few differences between them with respect to the position of the *E. aspergilloides* and *E. lentescens*. The BI analysis placed *E. aspergill-*




Figure 6. Phylogenetic position of *Escovopsis clavatus* and *Escovopsis multiformis* considering each molecular marker separately (ITS, LSU and *tef1*). The trees were reconstructed under Bayesian and Maximum Likelihood inferences. The numbers on branches indicate the posterior probabilities and the bootstrap support values, respectively. The seven *Escovopsis* ex-type strains are denoted in bold and the new species are highlighted in green (*E. clavatus*) and light brown (*E. multiformis*). The trees include a total of 46 *Escovopsis* sequences of each marker (ITS – 619 bp, LSU – 594 bp and *tef1* – 758 bp) and *Escovopsioides*, *Hypomyces*, *Sphaerostilbella*, *Trichoderma* and *Protocrea* were included as the closest phylogenetic relatives of *Escovopsis*. *Lecanicillium antillarum* CBS 350.85 was used as the outgroup. ET: ex-type.




Figure 7. Phylogenetic position of *Escovopsis clavatus* and *Escovopsis multififormis*. The phylogenetic analysis is based on the concatenated sequences of ITS, LSU and *tefl*; and the tree was reconstructed using Bayesian and Maximum Likelihood inferences. Numbers on branches indicate the posterior probabilities and the bootstrap support values, respectively. All *Escovopsis* species previously described are denoted in bold and the new species are highlighted in green (*E. clavatus*) and light brown (*E. multififormis*). The tree includes a total of 40 *Escovopsis* sequences with 1971 bp (ITS – 619 bp, LSU – 594 bp and *tefl* – 758 bp). The data also included sequences from *Escovopsioides*, *Hypomyces*, *Sphaerostilbella*, *Trichoderma* and *Protocrea* as the closest phylogenetic relatives of the parasite. *Lecanicillium antillarum* CBS 350.85 was used as the outgroup. ET: ex-type strains. Bar: 0.04 substitutions per nucleotide position.

loides and *E. lentescens* separate from *E. multififormis* and *E. clavatus* (Fig. 7); however, the ML analysis showed the former species as sister clades of *E. multififormis* and *E. clavatus*.

It is important to highlight that the concatenated analysis, as well as the trees inferred with ITS and LSU, showed the vesiculated *Escovopsis* (*E. aspergilloides*, *E. clavatus*, *E. lentescens*, *E. microspora*, *E. moelleri*, *E. multififormis*, *E. weberi*) as the most derived group, separated from the non-vesiculated *Escovopsis* (*E. kreiselii* and *E. trichodermoides*). In addition, both the combined and the analysis performed with ITS and *tefl* showed some *Escovopsis* species (*E. aspergilloides*, *E. kreiselii*, *E. lentescens* and *E. trichodermoides*) often clustering with other Hypocreaceae genera or falling outside the *Escovopsis* clade, which reveals that *Escovopsis* is apparently paraphyletic (Figs 6, 7).

Discussion

The attine ants have persisted for millions of years because of the biological relationships that these insects maintain with the beneficial microorganisms that inhabit their colonies. Several studies tried to understand how these biological relationships sculptured the evolutionary history of the attines (Mueller et al. 1998, Currie et al. 2003, Gerardo et al. 2006ab, Nygaard et al. 2016, Mueller et al. 2018). Nevertheless, the taxonomy of *Escovopsis*, the only known parasite in the attine's colony, has been poorly addressed. Considering that *Escovopsis* co-evolved with the attine ants' cultivar, improved knowledge about the taxonomy and systematics of this genus could shed light on the evolutionary success of these insects. Therefore, the discovery and description of new *Escovopsis* species is an important advance in understanding this system.

Subsequent to the formal description of *Escovopsis* (Muchovej and Della Lucia 1990), several studies showed a high genetic diversity of this genus in the colonies of both leafcutter and non-leafcutter attine ants (Gerardo et al. 2006a, Meirelles et al. 2015b). However, only seven species of the parasite have been described so far (Seifert et al. 1995, Augustin et al. 2013, Masiulionis et al. 2015, Meirelles et al. 2015a) and the morphological diversity and physiology of the parasite remain unknown. In addition, a lack of standardised conditions for describing the morphology of *Escovopsis* hinders researchers from identifying morphological characters that might help to distinguish *Escovopsis* species from one another and from the other related genera from the Hypocreaceae. Unfortunately, this fact made it difficult to describe new species of the parasite. Studies showed that the expressed phenotypic characters (phenotypic plasticity) of fungi are directly influenced by growth conditions (Slepecky and Starmer 2009, Sharma and Pandey 2010, Wrzosek et al. 2017, Kim et al. 2017). As the morphological plasticity of *Escovopsis* species is still poorly understood, the standardisation of cultivation conditions is imperative. The strains described as new species here were evaluated on eight different culture media (those used in the description of the seven previous species) and at five temperatures (to establish cardinal growth temperatures). Due to the lack of standard culture conditions, the comparison with each species previously described was only partial. Nonetheless, we are providing characters of these two new species in all the conditions previously used, to help future researchers to standardise the taxonomy of the genus.

Recent attempts to expand the morphological concept of *Escovopsis* generated inconsistencies in the taxonomy and systematics of this genus (Masiulionis et al. 2015, Meirelles et al. 2015a). The morphological characters that initially gave rise to the concept of *Escovopsis* (presence of terminal vesicles and phialidic conidiogenesis, see Muchovej and Della Lucia 1990) are distinctive to delineate *Escovopsis*, because no other genus in the Hypocreaceae family has such combined characters. However, some *Escovopsis* species described recently, namely *E. trichodermoides* and *E. kreiselii*, lack vesicles and each has a different kind of conidiogenesis (synchronous and sympodial, respectively). Besides, the results of the phylogenetic analysis performed in previous studies (Meirelles et al. 2015b, Masiulionis et al. 2015, Augustin et al. 2013), as well as the results from our analysis, reveal that *Escovopsis* is paraphyletic (Figs 6, 7). Therefore, future studies will have to reconsider if both species indeed belong to *Escovopsis*. For

this purpose, the taxonomic conditions need to be delimited and additional molecular markers will have to be included to help resolve those phylogenetic incongruities. Then the generic concept of *Escovopsis* species should be revisited.

Our study shows that the ex-type strains LESF 853 (*E. clavatus*) and LESF 847 (*E. multiformis*) form a monophyletic clade within most derived *Escovopsis* (vesiculated *Escovopsis*, PP = 1, BML = 100%). Most interesting, unlike the other *Escovopsis* species, the two new species present a unique type of conidiophore with a swollen cell, from which one to four branches arise. The newly described species also possess smooth conidia with slightly thickened walls. A recent study suggests the possibility that conidia ornamentation could be associated with the mechanism for horizontal transmission of *Escovopsis* between ant colonies and with the latency of the parasite conidia. This hypothesis was based on observations of some conidia adhering to the ant legs and in spore dormancy in vitro bioassays (Augustin et al. 2017). The same authors also argued that such character could be used as morphological markers for the taxonomy of the genus. However, because of scarce knowledge of the morphological features of the *Escovopsis* species, it is difficult to decipher which phenotypic character could be considered diagnostic for this genus. Therefore, future researchers need to carefully evaluate the phenotypic characters of each *Escovopsis* clade to determine which of characters are homologous versus those that are homoplasious to build taxonomic keys.

Considering the high genetic diversity of *Escovopsis* and the poor knowledge of its taxonomy, our study suggests that the fungus gardens of attine ants host a great diversity of *Escovopsis* that has yet to be discovered. Thus, the description of these new species are merely two small pieces of a complex puzzle. Nonetheless, our work should help future researchers to build the framework for the systematics of this parasitic fungus.

Acknowledgements

We are grateful to “Fundação de Amparo à Pesquisa do Estado de São Paulo” (FAPESP) for financial support (Grants # 2014/24298-1 and #2017/12689-4) conceded to AR and for the scholarship (# 2016/04955-3) to QVM. We also would like to thank Antônio Teruyoshi Yabuki (UNESP, Rio Claro) for SEM assistance, Lia Costa Pinto Wentzel, Rodolfo Bizarria Jr and Caitlin Conn for valuable comments on this manuscript. In addition, we are grateful to Dr. Huzefa Raja (reviewer #1), Keith Seifert (reviewer #2) and the editor Thorsten Lumbsch for providing constructive comments on this study.

References

Atili-Angelis D, Duarte APM, Pagnocca FC, Nagamoto NS, de Vries M, Stielow JB, de Hoog GS (2014) Novel *Phialophora* species from leaf-cutting ants (tribe Attini). Fungal Diversity 65(1): 65–75. <https://doi.org/10.1007/s13225-013-0275-0>

- Augustin JO, Groenewald JZ, Nascimento RJ, Mizubuti ESG, Barreto RW, Elliot SL, Evans HC (2013) Yet more “weeds” in the garden: Fungal novelties from nests of leaf-cutting ants. *PLoS One* 8(12): e82265. <https://doi.org/10.1371/journal.pone.0082265>
- Augustin JO, Simões TG, Dijksterhuis J, Elliot SL, Evans HC (2017) Putting the waste out: a proposed mechanism for transmission of the mycoparasite *Escovopsis* between leafcutter ant colonies. *Royal Society Open Science* 4(5): 161013. <https://doi.org/10.1098/rsos.161013>
- Branstetter MG, Ješovnik A, Sosa-Calvo J, Lloyd MW, Faircloth BC, Brady SG, Schultz TR (2017) Dry habitats were crucibles of domestication in the evolution of agriculture in ants. *Proceedings of the Royal Society B Biological Sciences* 284(1852): 20170095. <https://doi.org/10.1098/rspb.2017.0095>
- Castellani AMD (1963) Further researches on the long viability and growth of many pathogenic fungi and some bacteria in sterile distilled water. *Mycopathologia et Mycologia Applicata* 20: 1–6. <https://doi.org/10.1007/BF02054872>
- Chaverri P, Castlebury LA, Samuels GJ, Geiser DM (2003) Multilocus phylogenetic structure within the *Trichoderma harzianum*/*Hypocrea lixii* complex. *Molecular Phylogenetics and Evolution* 27(2): 302–313. [https://doi.org/10.1016/S1055-7903\(02\)00400-1](https://doi.org/10.1016/S1055-7903(02)00400-1)
- Currie CR, Wong B, Stuart AE, Schultz TR, Rehner SA, Mueller UG, Sung GH, Spatafora JW, Straus NA (2003) Ancient tripartite coevolution in the attine ant-microbe symbiosis. *Science* 299(5605): 386–388. <https://doi.org/10.1126/science.1078155>
- Currie CR (2001) Prevalence and impact of a virulent parasite on a tripartite mutualism. *Oecologia* 128(1): 99–106. <https://doi.org/10.1007/s004420100630>
- Custodio BC, Rodrigues A (2019) *Escovopsis kreiselii* specialization to its native hosts in the fungiculture of the lower attine ant *Mycetophylax morschi*. *Antonie van Leeuwenhoek* 112(2): 305–317. <https://doi.org/10.1007/s10482-018-1158-x>
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772. <https://doi.org/10.1038/nmeth.2109>
- Dhodary B, Schilg M, Wirth R, Spiteller D (2018) Secondary metabolites from *Escovopsis weberi* and their role in attacking the garden fungus of leaf-cutting ants. *Chemistry – A European Journal* 24(17): 4445–4452. <https://doi.org/10.1002/chem.201706071>
- Gerardo NM, Jacobs SR, Currie CR, Mueller UG (2006a) Ancient host-pathogen associations maintained by specificity of chemotaxis and antibiosis. *PLoS Biology* 4(8): e235. <https://doi.org/10.1371/journal.pbio.0040235>
- Gerardo NM, Mueller UG, Currie CR (2006b) Complex host-pathogen coevolution in the *Apterostigma* fungus-growing ant-microbe symbiosis. *BMC Evolutionary Biology* 6: 88. <https://doi.org/10.1186/1471-2148-6-88>
- Gerardo NM, Mueller UG, Price SL, Currie CR (2004) Exploiting a mutualism: parasite specialization on cultivars within the fungus-growing ant symbiosis. *Proceedings of the Royal Society B Biological Sciences* 271(1550): 1791–1798. <https://doi.org/10.1098/rspb.2004.2792>
- Harrington TC, McNew D, Mayers C, Fraedrich SW, Reed SE (2014) *Ambrosiella roeperi* sp. nov. is the mycangial symbiont of the granulate ambrosia beetle, *Xylosandrus crassiusculus*. *Mycologia* 106(4): 835–845. <https://doi.org/10.3852/13-354>

- Haugland RL, Heckman JL (1998) Identification of putative sequence specific PCR primers for detection of the toxigenic fungal species *Stachybotrys chartarum*. *Molecular and Cellular Probes* 12(6): 387–396. <https://doi.org/10.1006/mcpr.1998.0197>
- Heine D, Holmes NA, Worsley SF, Santos ACA, Innocent TM, Scherlach K, Patrick EH, Yu DW, Murrell JC, Viera PC, Boomsma JJ, Hertweck C, Hutchings M, Wilkinson B (2018) Chemical warfare between leafcutter ant symbionts and a co-evolved pathogen. *Nature Communications* 9(1): 2208. <https://doi.org/10.1038/s41467-018-04520-1>
- Hughes DP, Pierce NE, Boomsma JJ (2008) Social insect symbionts: evolution in homeostatic fortresses. *Trends in Ecology and Evolution* 23(12): 672–677. <https://doi.org/10.1016/j.tree.2008.07.011>
- Jaklitsch WM, Pöldmaa K, Samuels GJ (2011) Reconsideration of *Protocrea* (Hypocreales, Hypocreaceae). *Mycologia* 100(6): 962–984. <https://doi.org/10.3852/08-101>
- Ješovnik A, González VL, Schultz TR (2016) Phylogenomics and divergence dating of fungus-farming ants (Hymenoptera: Formicidae) of the genera *Sericomyrmex* and *Apterostigma*. *PLoS One* 11(7): e0151059. <https://doi.org/10.1371/journal.pone.0151059>
- Joop G, Vilcinskis A (2016) Coevolution of parasitic fungi and insect hosts. *Zoology* 119(4): 350–358. <https://doi.org/10.1016/j.zool.2016.06.005>
- Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kim YK, Xiao CL, Rogers JD (2005) Influence of culture media and environmental factors on mycelial growth and pycnidial production of *Sphaeropsis pyripitrescens*. *Mycologia* 97(1): 25–32. <https://doi.org/10.1080/15572536.2006.11832835>
- Kost C, Lakatos T, Böttcher I, Arendholz WR, Redenbach M, Wirth R (2007) Non-specific association between filamentous bacteria and fungus-growing ants. *Naturwissenschaften* 94(10): 821–828. <https://doi.org/10.1007/s00114-007-0262-y>
- Kwong WK, Medina LA, Koch H, Sing KW, Soh EJY, Ascher JS, Jaffé R, Moran NA (2017) Dynamic microbiome evolution in social bees. *Science Advances* 3(3): e1600513. <https://doi.org/10.1126/sciadv.1600513>
- Masiulionis VE, Cabello MN, Seifert KA, Rodrigues A, Pagnocca FC (2015) *Escovopsis trichodermoides* sp. nov., isolated from a nest of the lower attine ant *Mycocepurus goeldii*. *Antonie van Leeuwenhoek* 107(3): 731–40. <https://doi.org/10.1007/s10482-014-0367-1>
- Meirelles LA, Montoya QV, Solomon SE, Rodrigues A (2015a) New light on the systematics of fungi associated with attine ant gardens and the description of *Escovopsis kreiselii* sp. nov. *PLoS One* 10(1): e0112067. <https://doi.org/10.1371/journal.pone.0112067>
- Meirelles LA, Solomon SE, Bacci M, Wright AM, Mueller UG, Rodrigues A (2015b) Shared *Escovopsis* parasites between leaf-cutting and non-leaf-cutting ants in the higher attine fungus-growing ant symbiosis. *Royal Society Open Science* 2(9): 150257. <https://doi.org/10.1098/rsos.150257>
- Menezes C, Vollet-Neto A, Marsaioli AJ, Zampieri D, Fontoura IC, Luchessi AD, Imperatriz-Fonseca VL (2015) A Brazilian social bee must cultivate fungus to survive. *Current Biology* 25(21): 2851–2855. <https://doi.org/10.1016/j.cub.2015.09.028>

- Montoya QV, Meirelles LA, Chaverri P, Rodrigues A (2016) Unraveling *Trichoderma* species in the attine ant environment: description of three new taxa. *Antonie van Leeuwenhoek* 109(5): 633–651. <https://doi.org/10.1007/s10482-016-0666-9>
- Moriya S, Inoue T, Ohkuma M, Yaovapa T, Johjima T, Suwanarit P, Sangwanit U, Vongkaluang C, Noparatnaraporn N, Kudo T (2005) Fungal community analysis of fungus gardens in termite nests. *Microbes and Environments* 20(4): 243–252. <https://doi.org/10.1264/jsme2.20.243>
- Muchovej JJ, Della Lucia TC (1990) *Escovopsis*, a new genus from leaf-cutting ant nests to replace *Phialocladus* nomen invalidum. *Mycotaxon* 37: 191–195.
- Mueller UG, Rehner SA, Schultz TR (1998) The evolution of agriculture in ants. *Science* 281(5385): 2034–2038. <https://doi.org/10.1126/science.281.5385.2034>
- Mueller UG, Gerardo N (2002) Fungus-farming insects: multiple origins and diverse evolutionary histories. *Proceedings of the National Academy Sciences of the United States of America* 99(24): 15247–15249. <https://doi.org/10.1073/pnas.242594799>
- Mueller UG, Ishak HD, Bruschi SM, Smith CC, Herman JJ, Solomon SE, Mikheyev AS, Rabeling C, Scott JJ, Cooper M, Rodrigues A, Ortiz A, Brandão CRF, Lattke JE, Pagnocca FC, Rehner SA, Schultz TR, Vasconcelos HL, Adams RMM, Bollazzi M, Clark RM, Himler AG, LaPolla JS, Leal IR, Johnson RA, Roces F, Sosa-Calvo J, Wirth R, Bacci Jr M (2017) Biogeography of mutualistic fungi cultivated by leafcutter ants. *Molecular Ecology* 26(24): 6921–6937. <https://doi.org/10.1111/mec.14431>
- Mueller UG, Kardish MR, Ishak HD, Wright AM, Solomon SE, Bruschi SM, Carlson AL, Bacci Jr M (2018) Phylogenetic patterns of ant-fungus associations indicate that farming strategies, not only a superior fungal cultivar, explain the ecological success of leafcutter ants. *Molecular Ecology* 27(10): 2414–2434. <https://doi.org/10.1111/mec.14588>
- Nixon KC (2002) WinClada ver. 1.0000, Ithaca. [Published by the author]
- Nygaard S, Hu H, Li C, Schiøtt M, Chen Z, Yang Z, Xie Q, Ma C, Deng Y, Dikow RB, Rabeling C, Nash DR, Wcislo WT, Brady SG, Schultz TR, Zhang G, Boomsma JJ (2016) Reciprocal genomic evolution in the ant-fungus agricultural symbiosis. *Nature Communications* 7: 12233. <https://doi.org/10.1038/ncomms12233>
- Pöldmaa K (2011) Tropical species of *Cladobotryum* and *Hypomyces* producing red pigments. *Studies in Mycology* 68: 1–34. <https://doi.org/10.3114/sim.2011.68.01>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Sharma G, Pandey RR (2010) Influence of culture media on growth, colony character and sporulation of fungi isolated from decaying vegetable wastes. *Journal of Yeast and Fungal Research* 1(8): 157–164.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Fungal Barcoding Consortium (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy Sciences of the United States of America* 109(16): 6241–6246. <https://doi.org/10.1073/pnas.1117018109>

- Schultz TR, Brady SG (2008) Major evolutionary transitions in ant agriculture. *Proceedings of the National Academy Sciences of the United States of America* 105(14):5435–5440. <https://doi.org/10.1073/pnas.0711024105>
- Schultz TR, Sosa-Calvo J, Brady SG, Lopes CT, Mueller UG, Bacci Jr M, Vasconcelos HL (2015) The most relictual fungus-farming ant species cultivates the most recently evolved and highly domesticated fungal symbiont species. *The American Naturalist* 185(5): 693–703. <https://doi.org/10.1086/680501>
- Seifert KA, Samson RA, Chapela IH (1995) *Escovopsis aspergilloides*, a rediscovered hyphomycete from leaf-cutting ant nests. *Mycologia* 87(3): 407–413. <https://doi.org/10.2307/3760838>
- Slepecky RA, Starmer WT (2009) Phenotypic plasticity in fungi: a review with observations on *Aureobasidium pullulans*. *Mycologia* 101(6): 823–832. <https://doi.org/10.3852/08-197>
- Sosa-Calvo J, Ješovnik A, Lopes CT, Rodrigues A, Rabeling C, Bacci Jr M, Vasconcelos HL, Schultz T (2017) Biology of the relict fungus-farming ant *Apterostigma megacephala* Latke, including descriptions of the male, gyne, and larva. *Insectes Sociaux* 64(3): 329–346. <https://doi.org/10.1007/s00040-017-0550-2>
- Sosa-Calvo J, Schultz TR (2010) Three remarkable new fungus-growing ant species of the genus *Myrmicocrypta* (Hymenoptera: Formicidae), with a reassessment of the characters that define the genus and its position within the attini. *Annals of the Entomological Society of America* 103(2): 181–195. <https://doi.org/10.1603/AN09108>
- Spatafora JW, Sung GH, Sung JM, Hywel-Jones NL, White Jr JF (2007) Phylogenetic evidence for an animal pathogen origin of ergot and the grass endophytes. *Molecular Ecology* 16(8): 1701–11. <https://doi.org/10.1111/j.1365-294X.2007.03225.x>
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Taerum SJ, Cafaro MJ, Little AE, Schultz TR, Currie CR (2007) Low host-pathogen specificity in the leaf-cutting ant-microbe symbiosis. *Proceedings of the Royal Society B Biological Sciences* 274(1621): 1971–1978. <https://doi.org/10.1098/rspb.2007.0431>
- Hall TA (1999) Bioedit a user-friendly biological sequence alignment editor an analysis program for windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98
- Vanderpool D, Bracewell RR, McCutcheon JP (2018) Know your farmer: ancient origins and multiple independent domestications of ambrosia beetle fungal cultivars. *Molecular Ecology* 27(8): 2077–2094. <https://doi.org/10.1111/mec.14394>
- Varanda-Haifig SS, Albarici TR, Nunes PH, Haifig I, Vieira PC, Rodrigues A (2017) Nature of the interactions between hypocrealean fungi and the mutualistic fungus of leaf-cutter ants. *Antonie van Leeuwenhoek* 110(4): 593–605. <https://doi.org/10.1007/s10482-016-0826-y>
- Villesen P, Mueller UG, Schultz TR, Adams RMM, Bouck AC (2004) Evolution of ant-cultivar specialization and cultivar switching in *Apterostigma* fungus-growing ants. *Evolution* 58(10): 2252–2265. <https://doi.org/10.1111/j.0014-3820.2004.tb01601.x>
- White TJ, Bruns T, Lee SH, Taylor JW (1990) PCR Protocols: A Guide to Methods and Application. Academic Press, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wrzosek M, Ruskiewicz-Michalska M, Sikora K, Damszel M, Sierota Z (2017) The plasticity of fungal interactions. *Mycological Progress* 16(2): 101–108. <https://doi.org/10.1007/s11557-016-1257-x>

Supplementary material I

Supplementary tables

Authors: Quimi Vidaurre Montoya, Maria Jesus Sutta Martiarena, Danilo Augusto Polezel, Sérgio Kakazu, Andre Rodrigues

Data type: phylogenetic data

Explanation note: Table S1. *Escovopsis* strains used in the phylogenetic analyses and their associated metadata. Table S2. Molecular markers, primers and PCR conditions used in this study.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.46.30951.suppl1>