

New contributions to Diatrypaceae from karst areas in China

Sihan Long^{1,2}, Lili Liu³, Yinhui Pi¹, Youpeng Wu¹, Yan Lin¹, Xu Zhang¹,
Qingde Long¹, Yingqian Kang⁴, Jichuan Kang⁵, Nalin N. Wijayawardene^{1,6},
Feng Wang⁷, Xiangchun Shen^{1,2}, Qirui Li^{1,2}

1 State Key Laboratory of Functions and Applications of Medicinal Plants, Guizhou Medical University, Guiyang 550004, China **2** The High Efficacy Application of Natural Medicinal Resources Engineering Center of Guizhou Province (The Key Laboratory of Optimal Utilization of Natural Medicine Resources), School of Pharmaceutical Sciences, Guizhou Medical University, University Town, Gui'an New District, Guizhou 550025, China **3** Immune Cells and Antibody Engineering Research Center of Guizhou Province/ Key Laboratory of Biology and Medical Engineering, Guizhou Medical University, Guiyang 550004, China **4** Key Laboratory of Environmental Pollution Monitoring and Disease Control, Ministry of Education of Guizhou and Guizhou Talent Base for Microbiology and Human Health, School of Basic Medical Sciences, Guizhou Medical University, Guiyang, China **5** Engineering and Research Center for Southwest Bio-Pharmaceutical Resources of National Education Ministry of China, Guizhou University, Guiyang, Guizhou 550025, China **6** Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, Yunnan 655011, China **7** Guizhou Provincial Academician Workstation of Microbiology and Health, Guizhou Academy of Tobacco Science, Guiyang, Guizhou, 550000, China

Corresponding author: Qirui Li (lqrnd2008@163.com)

Academic editor: Andrew Miller | Received 20 May 2021 | Accepted 20 July 2021 | Published 20 August 2021

Citation: Long S-H, Liu L-L, Pi Y-H, Wu Y-P, Lin Y, Zhang X, Long Q-D, Kang Y-Q, Kang J-C, Wijayawardene NN, Wang F, Shen X-C, Li Q-R (2021) New contributions to Diatrypaceae from karst areas in China. MycoKeys 83: 1–37. <https://doi.org/10.3897/mycokeys.83.68926>

Abstract

In this study, fungal specimens of the family Diatrypaceae were collected from karst areas in Guizhou, Hainan and Yunnan Provinces, China. Morpho-molecular analyses confirmed that these new collections comprise a new genus *Pseudodiatrype*, three new species (*Diatrype lancangensis*, *Diatrypella pseudooregonensis* and *Eutypa cerasi*), a new combination (*Diatrypella oregonensis*), two new records (*Allodiatriype thailandica* and *Diatrypella vulgaris*) from China and two other known species (*Neoeutypella baoshanensis* and *Paraeutypella citricola*). The new taxa are introduced, based on multi-gene phylogenetic analyses (ITS, β -tubulin), as well as morphological analyses. The new genus *Pseudodiatrype* is characterised by its wart-like stromata with 5–20 ascomata immersed in one stroma and the endostroma composed of thin black

outer and inner layers of large white cells with thin, powdery, yellowish cells. These characteristics separate this genus from two similar genera *Allodiatriype* and *Diatrype*. Based on morphological as well as phylogenetic analyses, *Diatrype lancangensis* is introduced as a new species of *Diatrype*. The stromata of *Diatrype lancangensis* are similar to those of *D. subundulata* and *D. undulate*, but the ascospores are larger. Based on phylogenetic analyses, *Diatrype oregonensis* is transferred to the genus *Diatrypella* as *Diatrypella oregonensis* while *Diatrypella pseudooregonensis* is introduced as a new species of *Diatrypella* with 8 spores in an ascus. In addition, multi-gene phylogenetic analyses show that *Eutypa cerasi* is closely related to *E. lata*, but the ascocata and asci of *Eutypa cerasi* are smaller. The polyphyletic nature of some genera of Diatrypaceae has led to confusion in the classification of the family, thus we discuss whether the number of ascospores per ascus can still be used as a basis for classification.

Keywords

Five novel taxa, phylogeny, systematics, taxonomy, Xylariales

Introduction

Diatrypaceae is an important family of higher ascomycetes, belonging to Xylariales (Maharachchikumbura et al. 2016). In the latest compilation, Hyde et al. (2020a) revised the family Diatrypaceae and included several new genera (i.e. *Allodiatriype* Konta & K.D. Hyde, *Halocryptovalsa* Dayar. & K.D. Hyde and *Neoeutypella* M. Raza et al.). This was followed by Wijayawardene et al. (2020) in which 20 genera were accepted into Diatrypaceae. The Diatrypaceae is characterised by perithecial ascocata embedded in a poor or well-developed, brown or black-coloured stroma, long-stalked and 8-spored or numerous-spored asci and allantoid, unicellular ascospores (Glawe and Rogers 1984; Rappaz 1987; Mehrabi et al. 2015; de Almeida et al. 2016).

Members of Diatrypaceae occur on a wide range of hosts in terrestrial and marine environments worldwide, some of which are important plant pathogens (Moyo et al. 2018a; Mehrabi et al. 2019; Dayarathne et al. 2020; Konta et al. 2020). For many decades, canker diseases on grapevine have been attributed to the species of Diatrypaceae worldwide, for example in China *Cryptovalsa* Ces. & De Not., *Cryptosphaeria* Ces. & De Not, *Diatrype* Fr., *Diatrypella* (Ces. & De Not.) De Not., *Eutypa* Tul. & C. Tul. And *Eutypella* (Nitschke) Sacc., are responsible for canker diseases in grapevine (Trouillas et al. 2011; Gao et al. 2013; Moyo et al. 2018b). Besides cankers of grapevine, some species have been reported as the causal pathogenic agents of fruit trees and woody plants in Europe and the USA (Trouillas et al. 2011; Gao et al. 2013).

Thirteen species of *Cryptosphaeria* and *Diatrype* were introduced by Vasiljeva and Ma (2014) from north-eastern China, which includes two new species and four new records. China has the largest range of karst distribution in the world. The landform of karst can be found in almost all Provinces of China, with the most extensive distribution in Guizhou and Yunnan Provinces (Miao et al. 2007). Karst virgin forest is a relatively stable ecosystem with rich biological resources, highly primitive and maintaining stable biological diversity (Dong et al. 2002). The special karst and ecological environment is home to a rich diversity of diatrypaceous fungi.

In this study, we revisit species of Diatrypaceae collected from karst areas in Guizhou, Hainan and Yunnan Provinces of China. Based on morpho-molecular analyses, one new genus and three new species are introduced; in addition, a new combination and two new records from China are reported. Descriptions and illustrations of new taxa and new records are provided.

Materials and Methods

Fungi collection, isolation and identification

Samples of decaying wood were collected from October 2019 to November 2020 in forests and nature reserves of Guizhou, Hainan and Yunnan Provinces in China. The specimens were observed with a stereomicroscope while microscopic images of the samples were taken using a Nikon ECLIPSE Ni compound microscope, with a Canon EOS 700D digital camera. Measurements were taken with Tarosoft (R) Image Frame Work (v.0.9.7). More than 30 ascospores were measured for each specimen examined. Photoplates were arranged and improved by using Adobe Photoshop CS6 software. Isolations of fungi were made by single spore isolation (Chomnunti et al. 2014) and germinated spores were transferred to potato dextrose agar (**PDA**) medium for purification. The specimens were deposited at the Herbarium of Cryptogams, Kunming Institute of Botany Academia Sinica (**KUN-HKAS**) and Herbarium of Guizhou Medical University (**GMB**). Strains of the new genus and new species are maintained in the Guizhou Medical University Collection Centre (**GMBC**).

DNA extraction, Polymerase Chain Reaction (PCR) and phylogenetic analyses

Genomic DNA was extracted from fungal mycelium following the manufacturer's protocol of the BIOMIGA Fungal gDNA isolation Kit (BIOMIGA, Hangzhou City, Zhejiang Province, China). Extracts of DNA were stored at -20 °C.

PCR was carried out in a volume of 25 µl containing 9.5 µl of ddH₂O, 12.5 µl of 2× Taq PCR Master Mix (2 × Taq Master Mix with dye, TIANGEN, China), 1 µl of DNA extracts and 1 µl of forward and reverse primers (10 µM each) in each reaction. Primers pairs, ITS4 and ITS5, fRPB2-7CR and fRPB2-5f, LROR and LR5, T1 and Bt2b, as well as Bt2a and Bt2b (Vilgalys and Hester 1990; White et al. 1990; Glass and Donaldson 1995; O'Donnell and Cigelnik 1997), were used to amplify internal transcribed spacer (ITS) sequences, RNA polymerase II second largest subunit (RPB2) sequences, large subunit ribosomal (LSU) sequences and β-tubulin (TUB2) sequences, respectively.

PCR profiles for the ITS and LSU are as follows: initially at 95 °C for 5 minutes, followed by 35 cycles of denaturation at 94 °C for 1 minute, annealing at 52 °C for 1 minute, elongation at 72 °C for 1.5 minutes and a final extension at 72 °C for 10 minutes. PCR profile for the RPB2 is as follows: initially at 95 °C for 5 minutes, followed by 35 cycles of denaturation at 95 °C for 1 minute, annealing at 54 °C for 2 minutes,

Table I. Taxa used in the phylogenetic analyses and their corresponding GenBank accession numbers.

Taxa	Strain number	GenBank Accession number		Reference
		ITS	β -tubulin	
<i>Allocryptotarsa elaeidis</i>	MFLUCC 15-0707	MN308410	MN340296	Konta et al. (2020)
<i>A. polyporat</i>	MFLUCC 17-0364	MF959500	MG334556	Senwanna et al. (2017)
<i>A. rabenhorstii</i>	WA08CB	HQ692619	HQ692523	Trouillas et al. (2011)
<i>Allodiatrype arencae^T</i>	MFLUCC 15-0713	MN308411	MN340297	Konta et al. (2020)
<i>A. elaeidicola</i>	MFLUCC 15-0737a	MN308415	MN340299	Konta et al. (2020)
<i>A. elaeidis</i>	MFLUCC 15-0708a	MN308412	MN340298	Konta et al. (2020)
<i>A. thailandica</i>	MFLUCC 15-3662	KU315392	NA	Li et al. (2016)
<i>A. thailandica</i>	MFLUCC 15-0711	MN308414	NA	Konta et al. (2020)
<i>A. thailandica</i>	GMB0050	MW797108	MW814880	This study
<i>Anthostoma decipiens^T</i>	IPV-FW349	AM399021	AM920693	Unpublished
<i>A. decipiens</i>	JL567	JN975370	JN975407	Luque et al. (2012)
<i>Cryptosphaeria ligniota</i>	CBS 273.87	KT425233	KT425168	Acero et al. (2004)
<i>C. pullmanensis</i>	ATCC 52655	KT425235	KT425170	Trouillas et al. (2015)
<i>C. subcutanea</i>	DSUB100A	KT425189	KT425124	Trouillas et al. (2015)
<i>C. subcutanea</i>	CBS 240.87	KT425232	KT425167	Trouillas et al. (2015)
<i>Cryptotarsa ampelina</i>	A001	GQ293901	GQ293972	Trouillas et al. (2010)
<i>C. ampelina</i>	DRO101	GQ293902	GQ293982	Trouillas et al. (2010)
<i>Diatrype bullata</i>	UCDDCh400	DQ006946	DQ007002	Rolshausen et al. (2006)
<i>D. disciformis^T</i>	GNA14	KR605644.1	KY352434.1	Senanayake et al. (2015)
<i>D. disciformis</i>	D21C, CBS 205.87	AJ302437	NA	Acero et al. (2004)
<i>D. enteroxantha</i>	HUEFS155114	KM396617	KT003700	de Almeida et al. (2016)
<i>D. enteroxantha</i>	HUEFS155116	KM396618	KT022236	de Almeida et al. (2016)
<i>D. lancangensis</i>	GMB0045	MW797113	MW814885	This study
<i>D. lancangensis</i>	GMB0046	MW797114	MW814886	This study
<i>D. lancangensis</i>	GMB0047	MW797116	MW814887	This study
<i>D. palmicola</i>	MFLUCC 11-0020	KP744438	NA	Liu et al. (2015)
<i>D. palmicola</i>	MFLUCC 11-0018	KP744439	NA	Liu et al. (2015)
<i>D. spilomea</i>	D17C	AJ302433	NA	Acero et al. (2004)
<i>D. stigma</i>	DCASH200	GQ293947	GQ294003	Trouillas et al. (2010)
<i>D. undulata</i>	D20C, CBS 271.87	AJ302436	NA	Acero et al. (2004)
<i>Diatrypella atlantica</i>	HUEFS 136873	KM396614	KR259647	de Almeida et al. (2016)
<i>D. banksiae</i>	CPC 29118	KY173402	NA	Crous et al. (2013)
<i>D. delonicis</i>	MFLUCC 15-1014	MH812994	MH847790	Hyde et al. (2019)
<i>D. delonicis</i>	MFLU 16-1032	MH812995	MH847791	Hyde et al. (2019)
<i>D. elaeidis</i>	MFLUCC 15-0279	MN308417	MN340300	Konta et al. (2020)
<i>D. favacea</i>	Islotate 380	KU320616	NA	de Almeida et al. (2016)
<i>D. favacea</i>	DL26C	AJ302440	NA	Unpublished
<i>D. frostii</i>	UFMGC B 1917	HQ377280	NA	Vieira et al. (2011)
<i>D. heveae</i>	MFLUCC 15-0274	MN308418	MN340301	Konta et al. (2020)
<i>D. heveae</i>	MFLUCC 17-0368	MF959501	MG334557	Senwanna et al. (2017)
<i>D. hubeiensis</i>	CFCC 52413	MW632937	NA	Zhu et al. (2021)
<i>D. iranensis</i>	KDQ18	KM245033	KY352429	Mehrabi et al. (2015)
<i>D. macrospora</i>	KDQ15	KR605648	KY352430	Mehrabi et al. (2016)
<i>D. oregonensis</i> (<i>Diatrype oregonensis</i>)	DPL200	GQ293940	GQ293999	Trouillas et al. (2010)
<i>D. oregonensis</i> (<i>Diatrype oregonensis</i>)	CA117	GQ293934	GQ293996	Trouillas et al. (2010)
<i>D. pseudooregonensis</i>	GMB0039	MW797115	MW814888	This study
<i>D. pseudooregonensis</i>	GMB0040	MW797117	MW814889	This study
<i>D. pseudooregonensis</i>	GMB0041	MW797118	MW814890	This study
<i>D. pseudooregonensis</i>	GMB0042	MW797119	MW814891	This study
<i>D. pseudooregonensis</i>	GMB0043	MW797120	MW814892	This study
<i>D. pseudooregonensis</i>	GMB0044	MW797110	MW814882	This study
<i>D. pulvinata</i>	H048	FR715523	FR715495	de Almeida et al. (2016)
<i>D. pulvinata</i>	DL29C	AJ302443	NA	Unpublished
<i>D. tectonae</i>	MFLUCC 12-0172a	KY283084	NA	Shang et al. (2017)
<i>D. tectonae</i>	MFLUCC 12-0172b	KY283085	KY421043	Shang et al. (2017)
<i>D. verruciformis^T</i>	UCROK1467	JX144793	JX174093	Lynch et al. (2013)

Taxa	Strain number	GenBank Accession number		Reference
		ITS	β -tubulin	
<i>D. verruciformis</i> ^T	UCROK754	JX144783	JX174083	Lynch et al. (2013)
<i>D. vulgaris</i>	HVFRA02	HQ692591	HQ692503	Trouillas et al. (2011)
<i>D. vulgaris</i>	HVGRF03	HQ692590	HQ692502	Trouillas et al. (2011)
<i>D. vulgaris</i>	GMB0051	MW797107	MW814879	This study
<i>D. yunnanensis</i>	VT01	MN653008	MN887112	Zhu et al. (2021)
<i>Eutypa armeniacae</i>	ATCC 28120	DQ006948	DQ006975	Rolshausen et al. (2006)
<i>E. astroidea</i>	E49C, CBS 292.87	AJ302458	DQ006966	Rolshausen et al. (2006)
<i>E. cerasi</i>	GMB0048	MW797104	MW814893	This study
<i>E. cerasi</i>	GMB0049	MW797105	MW814877	This study
<i>E. flavovirens</i>	E48C, CBS 272.87	AJ302457	DQ006959	Rolshausen et al. (2006)
<i>E. laevata</i>	E40C CBS 291.87	AJ302449	NA	Acero et al. (2004)
<i>E. lata</i> ^T	CBS290.87	HM164736	HM164770	Trouillas and Gubler (2010)
<i>E. lata</i> ^T	EP18	HQ692611	HQ692501	Trouillas et al. (2011)
<i>E. lata</i> ^T	RGA01	HQ692614	HQ692497	Trouillas et al. (2011)
<i>E. leioplaca</i>	CBS 248.87	DQ006922	DQ006974	Rolshausen et al. (2006)
<i>E. leptoplaca</i>	CBS 287.87	DQ006924	DQ006961	Rolshausen et al. (2006)
<i>E. maura</i>	CBS 219.87	DQ006926	DQ006967	Rolshausen et al. (2006)
<i>E. microasca</i>	BAFC 51550	KF964566	KF964572	Grassi et al. (2014)
<i>E. sparsa</i>	3802 3b	AY684220	AY684201	Trouillas and Gubler (2004)
<i>E. tetragona</i>	CBS284.87	DQ006923	DQ006960	Rolshausen et al. (2006)
<i>Eutypella caricae</i>	EL51C	AJ302460	NA	Acero (2000)
<i>E. cerviculata</i> ^T	M68	JF340269	NA	Arhipova et al. (2012)
<i>E. cerviculata</i> ^T	EL59C	AJ302468	NA	Acero et al. (2004)
<i>E. leprosa</i>	EL54C	AJ302463	NA	Acero et al. (2004)
<i>E. leprosa</i>	Isolate 60	KU320622	NA	de Almeida et al. (2016)
<i>E. microtheca</i>	BCMX01	KC405563	KC405560	Paolinelli-Alfonse et al. (2015)
<i>E. parasitica</i>	CBS 210.39	DQ118966	NA	Jurc et al. (2006)
<i>E. semicircularis</i>	MP4669	JQ517314	NA	Mehrabi et al. (2016)
<i>Halocryptovalsa salicorniae</i>	MFLUCC 15-0185	MH304410	MH370274	Dayaratne et al. (2020)
<i>Halodiatrype avicenniae</i>	MFLUCC 15-0953	KX573916	KX573931	Dayaratne et al. (2016)
<i>H. salinicolus</i> ^T	MFLUCC 15-1277	KX573915	KX573932	Dayaratne et al. (2016)
<i>Kretzschmaria deusta</i>	CBS 826.72	KU683767	KU684190	U'Ren et al. (2016)
<i>Monosporascus cannonballus</i> ^T	CMM3646	JX971617	NA	Unpublished
<i>M. cannonballus</i> ^T	ATCC 26931	FJ430598	NA	Unpublished
<i>Neoeutypella baoshanensis</i> ^T	GMB0052	MW797106	MW814878	This study
<i>N. baoshanensis</i> ^T	LC 12111	MH822887	MH822888	Hyde et al. (2019)
<i>N. baoshanensis</i> ^T	EL51C, CBS 274.87	AJ302460	NA	Acero et al. (2004)
<i>N. baoshanensis</i> ^T	MFLUCC 16-1002	MT310662	NA	Phukhamsakda et al. (2020)
<i>N. baoshanensis</i> ^T	GL08362	JX241652	NA	Gao et al. (2013)
<i>Paraeutypella citricola</i>	HVVIT07	HQ692579	HQ692512	Trouillas et al. (2011)
<i>Pa. citricola</i>	HVGFR01	HQ692589	HQ692521	Trouillas et al. (2011)
<i>Pa. citricola</i>	GMB0053	MW797109	MW814881	This study
<i>Pa. guizhouensis</i> ^T	KUMCC 20-0016	MW039349	MW239660	Dissanayake et al. (2021)
<i>Pa. guizhouensis</i> ^T	KUMCC 20-0017	MW036141	MW239661	Dissanayake et al. (2021)
<i>Pa. vitis</i>	UCD2291AR	HQ288224	HQ288303	Úrbez-Torres et al. (2012)
<i>Pa. vitis</i>	UCD2428TX	FJ790851	GU294726	Úrbez-Torres et al. (2009)
<i>Pedumispora rhizophorae</i> ^T	BCC44877	KJ888853	NA	Klaysuban et al. (2014)
<i>Pe. rhizophorae</i> ^T	BCC44878	KJ888854	NA	Klaysuban et al. (2014)
<i>Peroneutypa alsophila</i>	EL58C, CBS 250.87	AJ302467	NA	Acero et al. (2004)
<i>Pe. curvispora</i>	HUEFS 136877	KM396641	NA	de Almeida et al. (2016)
<i>Pe. diminutispora</i>	MFLUCC 17-2144	MG873479	NA	Shang et al. (2018)
<i>Pe. mackenziei</i>	MFLUCC 16-0072	KY283083	KY706363	Shang et al. (2017)
<i>Pe. mangrovei</i>	PUFD526	MG844286	MH094409	Phookamsak et al. (2019)
<i>Pseudodiatrype hainanensis</i> ^T	GMB0054	MW797111	MW814883	This study
<i>Ps. hainanensis</i> ^T	GMB0055	MW797112	MW814884	This study
<i>Quaternaria quaternata</i>	EL60C, CBS 278.87	AJ302469	NA	Acero et al. (2004)
<i>Q. quaternata</i>	GNF13	KR605645	NA	Mehrabi et al. (2016)
<i>Xylaria hypoxylon</i>	CBS 122620	AM993141	KX271279	Peršoh et al. (2009)

^T: Types species of the genus; NA: No sequence is available in GenBank; Newly generated sequences are indicated in **bold**.

elongation at 72 °C for 1.5 minutes and a final extension at 72 °C for 10 minutes (Konta et al. 2020). PCR profile for the TUB2 are as follows: initially at 95 °C for 5 minutes, followed by 35 cycles of denaturation at 94 °C for 1 minute, annealing at 52 °C for 1 minute, elongation at 72 °C for 1.5 minutes and a final extension at 72 °C for 10 minutes (de Almeida et al. 2016). PCR products were submitted to Sangon Biotech, Shanghai, China for purification and sequencing.

Phylogenetic analyses

Phylogenetic analyses were performed by searching homologous sequence data of the family Diatrypaceae in the GenBank database, selected from NCBI and recently published papers (Mehrabi et al. 2019; Dayarathne et al. 2020; Konta et al. 2020; Dissanayake et al. 2021; Zhu et al. 2021). After the preliminary identification results of the sequences, multiple sequence alignments (ITS and β -tubulin) were aligned using BioEdit v. 7.0 (Hall 1999). Alignments were converted from FASTA to PHYLIP format by using Alignment Transformation Environment online (<https://sing.ei.uvigo.es/ALTER/>, Glez-Peña et al. 2010). Maximum Likelihood (ML) analyses and Bayesian posterior probabilities (BYPP) were performed by using RAxML-HPC BlackBox (8.2.12) and MrBayes on XSEDE (3.2.7a) tools in the CIPRES Science Gateway platform, based on a combination of ITS and TUB2 sequence data (Miller et al. 2010). Both of the two methods use the GTR+I+G model of evolution (Nylander 2004). The Bootstrap supports of ML analyses were obtained by running 1,000 pseudo-replicates and BYPP using a simulation technique called Markov chain Monte Carlo (or MCMC) to approximate the posterior probabilities of trees. Six simultaneous Markov Chains were run for 3,000,000 generations and trees were sampled every 1,000th generation. Finally, the tree was visualised in FigTree v.1.4.4 (Rambaut 2012) and edited by using Adobe Photoshop CS6 software. The final alignment and phylogenetic trees were deposited in TreeBASE under the submission ID28176 (<http://www.treebase.org/>)

Result

Phylogenetic analyses

Based on RAxML and BYPP analyses, phylogenetic analyses were similar in overall tree topologies and did not differ significantly. The dataset consists of 105 taxa for representative strains of species in Diatrypaceae, including outgroup taxa with 1071 characters, including gaps (ITS: 1–486, β -tubulin: 486–1071). The RAxML analyses resulted in a best scoring likelihood tree selected with a final ML optimisation likelihood value of -15731.506304, which is shown in Fig. 1.

The phylogenetic tree, based on combining ITS and β -tubulin sequence data, is also shown in Fig. 1 and contains 17 clades within Diatrypaceae. Below, we list the placements of new taxa:




Figure 1. Phylogram generated from Maximum Likelihood (RAxML) analyses, based on ITS- β -tubulin matrix. ML bootstrap supports ($\geq 70\%$) and Bayesian posterior probability (≥ 0.90) are indicated as ML/BYPP. The tree is rooted to *Kretzschmaria deusta* (CBS 826.72) and *Xylaria hypoxylon* (CBS 122620). Ex-type strains are in red. Newly generated strains are in black bold.




Figure 1. Continued.

Clade 1: *Diatrypella pseudooregonensis* and *Diatrypella oregonensis* clustered with the species of *Diatrypella* in Clade 1 with high bootstrap support, *Diatrypella pseudooregonensis* is introduced as an 8-spored new species of *Diatrypella* and *Diatrypella oregonensis* is renamed as *Diatrypella oregonensis*.

Clade 4: *Pseudodiatrypella* formed a separate branch in a clade (Clade 4) basal to the genus *Allodiatripella*.

Clade 7: *Diatrypella lancangensis* clusters with the species of *Diatrypella* and *Diatrypella* in an unresolved clade. However, *Diatrypella* and *Diatrypella* have previously shown confused classification which is difficult to distinguish, based on phylogenetic aspects alone. Therefore, we introduce *Diatrypella lancangensis* as a new species of *Diatrypella*, based on phylogenetic analyses and morphological differences (Table 2).

Clade 8: *Eutypa cerasi* forms a distinct lineage which is sister to *Eutypa lata* (EP18, RGA01) (Fig. 1).

Taxonomy

Diatrypella Fr.

Notes. The genus *Diatrypella* was introduced by Fries (1849). The genus is characterised by stromata widely effuse or verrucose, flat or slightly convex, with discoid or sulcate ostioles at the surface, 8-spored and long-stalked ascii and hyaline or brownish, allantoid ascospores. In this study, we introduce a new species of *Diatrypella* from China.

***Diatrype lancangensis* S.H. Long & Q. R. Li, sp. nov.**

MycoBank No: 839655

Fig. 2

Holotype. GMB0045.

Etymology. Refers to the name of the location, where the type specimen was collected.

Description. Saprobic on decaying branches of an unidentified plant. **Sexual morph:** *Stromata* immersed in bark, aggregated, irregular in shape, widely effused, flat, margin diffuse, surface dark brown to black, with punctiform ostioles scattered at surface, with tissues soft, white between perithecia. *Entostroma* dark with embedded perithecia in one layer. *Perithecium* semi-immersed in stroma, globose to subglobose, glabrous, with cylindrical neck, brevicolous or longicolous 283.5–343.5 µm high, 207–290 µm broad (av. = 315.5 × 248.0 µm, n = 10), ovoid, obovoid to oblong, monostichous, aterrimus. *Ostiole* opening separately, papillate or apapillate, central. *Peridium* 30–50 µm thick, dark brown to hyaline with *textura angularis* cell layers. *Asci* 90.5–160.5 × 7.0–15.0 µm (av. = 129.5 × 10.5 µm n = 30) 8-spored clavate, unitunicate, with rounded apex, apical rings inamyloid. *Ascospores* 11–18.5 × 2–4 µm (av. = 14.9 × 2.8 µm, n = 30), irregularly arranged, allantoid, slightly curved, brown to dark brown, smooth, aseptate, usually with oil droplets. Asexual morph: undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became luteous, dense but, thinning towards edge, margin rough, white from above, reverse white at margin, pale yellow to luteous at centre, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.

Specimens examined. CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve (25°1'17.44"N, 99°35'10.05"E) on branches of an unidentified plant, 4 October 2019. Altitude: 2549 m., Y.H. Pi & Qiong Zhang, LC172 (GMB0045, *holotype*, KUN-HKAS 112664, *isotype*, ex-type living culture GMBC0045).

Additional specimens examined. CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve (25°1'17.44"N, 99°35'10.05"E) on branches of an unidentified plant, 4 October 2019. Altitude: 2549 m., Y.H. Pi and Qiong Zhang, LC173 (GMB0046, KUN-HKAS 112665, living culture GMBC0046); CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve (25°1'15.48"N, 99°35'24.08"E) on branches of an unidentified plant, 5 October 2019. Altitude: 2623 m., Y.H. Pi and Qiong Zhang, LC262 (GMB0047, KUN-HKAS 112672, living culture GMBC0047).

Additional sequences. GMB0045 (LSU: MW797057, RPB2: MW81490); GMB0046 (LSU: MW797058); GMB0047 (LSU: MW797060, RPB2: MW814903)

Note. Our new strain, GMBC0045 falls into the unresolved clade (Clade 7) which comprises five *Diatrypella* and one *Diatrype* species (Fig. 1), this clade is consistent with the study of Konta et al. (2020). The taxonomic confusion of Diatrypaceae has led to difficulties in separating the genera. We consider that the new species belongs to the genus *Diatrype*, based on the stromata features mentioned above which closely resemble descriptions of *Diatrype subundulata* Lar. N. Vassiljeva & Hai X. Ma and *Diatrype undu-*




Figure 2. *Diatrype lancangensis* (GMB0045, holotype) **A** stromata on host substrate **B, C** stromata on host **D** transverse sections through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** ostiolar canal **H** peridium **I–K** ascospores **L–N** asci. Scale bars: 10 μm (**G–N**).

lata (Pers.) Fr. (Vasilyeva et al. 2014). However, the ascospores of these species are larger than the ascospores of *D. subundulata* and *D. undulata* (Table 2). Phylogenetic analyses also showed that *D. lancangensis* falls on a separate branch that clustered with species of *Diatrypella* and *Diatrype* (Fig. 1). Hence, by combining morphological characteristics and phylogenetic analyses, it seems appropriate to categorise this species as *Diatrype*.

In the phylogenetic analyses, it can be seen that Clade 7 can be defined as a new genus, but it is difficult to find the common morphological similarities among these species. More specimens and sequence or chemical composition analysis are needed in the future to determine whether Clade 7 can be a new genus. The characteristics of the stromata of *Diatrypella* spp. in clade 7 are solitary and scattered, which is distinctly different from widely effuse, flat and slightly convex stromata of *Diatrype lancangensis* and *Diatrype palmicola* (Liu et al. 2015; Hyde et al. 2020b; Zhu et al. 2021). And in the recent study, Zhu et al. (2021) proposed that the species of *Diatrypella* in Clade 7 were isolated from *Betula* spp., it may have host specificity. Because of the above two reasons, we think it is better to classify our strains into *Diatrype*.

***Pseudodiatrype* S.H. Long & Q.R. Li, gen. nov.**

MycoBank No: 839658

Etymology. Refers to this genus resembling *Diatrype* in morphology, but it is phylogenetically distinct.

Type species. *Pseudodiatrype hainanensis* S. H. Long & Q.R. Li sp. nov.

Description. Saprobic on decaying branches of an unidentified plant. **Sexual morph:** Stromata scattered or aggregated on host, wart-like, pustulate, visible as black, rounded to irregular in shape on host surface, erumpent through host bark, 5–20 ascomata immersed in one stroma. Endostroma consists of outer layer of black, small, dense, thin parenchymal cells and inner layer of white, large, loose parenchymal cells, thin, pale yellow, powdery near margin of the black cells. Ostiole opening through host bark and appearing as black spots, separately, papillate or apapillate, central. Peritheциum immersed in stroma, globose to subglobose, glabrous, with cylindrical neck, brevicolous or longicolous. Peridium is composed of an outer layer of dark brown to black, thin-walled cells, arranged in *textura angularis*, the inner layer of hyaline thin-walled cells of *textura angularis*. Ascii 8-spored, unitunicate, clavate, long-stalked, apically rounded, apical rings inamyloid. Ascospores irregularly arranged, allantoid, slightly or moderately curved, smooth, subhyaline, aseptate, usually with two oil droplets. **Asexual morph:** undetermined.

Note. The genus *Pseudodiatrype* is introduced to accommodate the new collection made from Hainan Province of China and typified by *Pseudodiatrype hainanensis*. *Pseudodiatrype* is monotypic and, morphologically, resembles *Diatrype* and *Allodiatriype* Konta & K.D. Hyde. However, *Pseudodiatrype* can be distinguished from *Diatrype* by its 5–20 ascomata immersed in a stroma, while the stroma of species of *Diatrype* is

distributed over large areas, sometimes covering the surface of the host (Vasilyeva and Ma 2014; Konta et al. 2020). *Pseudodiatripe* differs from *Allodiatripe* by having its 5–20 ascocarps immersed in a stroma, whereas the stroma of *Allodiatripe* has only 1–10 ascocarps. Moreover, the endostroma of *Allodiatripe* is composed of dark brown outer layer cells and yellow inner layer cells (Konta et al. 2020), which are different from the endostroma of *Pseudodiatripe* having black outer and inner cells surrounded by powdery, pale yellow cells. In addition, the sizes of stroma and ascospores are different from species of *Diatrype* and *Allodiatripe* (Table 2). In the phylogenetic analyses, species of *Pseudodiatripe* appeared in a separate branch which is distinct from other genera within *Diatrypaceae* (Fig. 1), thus, justifying the erection of the new genus *Pseudodiatripe*.

***Pseudodiatripe hainanensis* S. H. Long & Q.R. Li, sp. nov.**

MycoBank No: 839659

Fig. 3

Holotype. GMB0054.

Eymology. Refers to the location of collections, Hainan Province.

Description. *Saprobic* on decaying branches of an unidentified plant. **Sexual morph:** *Stromata* wart-like, pustulate, 2–3.6 mm long and 1.6–3 mm broad (av. = 3.2 × 1.9 mm, n = 30), about 2 mm thick, 5–20 in single stroma, visible as black, rounded to irregular in shape on the host surface, erumpent through host bark, solitary to gregarious. *Endostroma* composed of an outer layer of dark brown to black, small, tightly packed, thin parenchymatous cells and an inner layer of white, large, loose parenchymal cells with powdery, thin, yellowish tissue. *Ostiole* opening separately, papillate or apapillate, central. *Perithecium* immersed in the stroma, globose to subglobose, glabrous, with cylindrical neck, brevicollous or longicollous, 193–347 µm high, 138–206 µm diam. (av. = 278 × 156 µm, n = 10). *Peridium* 30–50 µm thick, dark brown to hyaline with *textura angularis* cell layers. *Asci* 110–155.5 × 6–10 µm (av. = 132 × 8 µm, n = 30), 8-spored, unitunicate, clavate, long-stalked, apically rounded with inamyloid rings. *Ascospores* 8.5–13 × 1.5–2.5 µm (av. = 10.5 × 2 µm, n = 30), irregularly arranged, allantoid, slightly or moderately curved, smooth, subhyaline, aseptate, usually with two oil droplets. **Asexual morph:** undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became pale brown, dense, but thinning towards edge, fluffy to slightly fluffy, white from above, pale brown from below, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.

Specimens examined. CHINA, Hainan Province, Wuzhishan City, Wuzhishan Nature Reserve (18°54'21.81"N, 109°40'54.12"E) on branches of unidentified plant, 14 November 2020. Altitude: 775 m. Y.H. Pi & Q.R. Li, WZS59 (GMB0054, *holotype*, KUN-HKAS 112700, *isotype*, ex-type living culture GMBC0054).

Additional specimen examined. CHINA, Hainan Province, Wuzhishan City, Wuzhishan Nature Reserve (18°54'21.81"N, 109°40'54.12"E) on branches of an uni-




Figure 3. *Pseudodiatrype hainanensis* (GMB0054, holotype) **A** stromata on host substrate **B**, **C** stromata on host **D** transverse section through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H** ostiolar canal **I** peridium **J–M** ascospores **N–P** asci. Scale bars: 40 µm (**G**); 10 µm (**H–P**).

dentified plant, 14 November 2020. Altitude: 775 m, Y.H. Pi & Q.R. Li, WZS66 (GMB0055, living culture GMBC0055)

Additional sequences. GMB0054 (LSU: MW797055, RPB2: MW814900); GMB0055 (LSU: MW797056, RPB2 MW814901).

Note. A peculiar feature of *Pseudodiatrype hainanensis* is the composition of endostroma. There are black outer layer cells, white inner layer cells and powdery, yellowish cells that are smaller than the white cells at the edge of the endostroma near the black cells in endostroma.

Diatrypella (Ces. & De Not.) De Not.

Notes. The genus *Diatrypella* was introduced by Cesati & De Notaris (1863) and was typified with *Diatrypella verruciformis* (Ehrh.) Nitschke. This genus was characterized by pustule-like stromata erumpent through the host surface, polyporous ascii and allantoid ascospores and libertella-like asexual morphs (Senanayake et al. 2015; Hyde et al. 2017; Shang et al. 2017). In this study, we introduce a new species, a new combination and a new record of *Diatrypella vulgaris* from Guizhou Province for China.

Diatrypella pseudooregonensis S.H. Long & Q.R. Li, sp. nov.

MycoBank No: 839656

Fig. 4

Holotype. GMB0041

Etymology. Refers to its similar species of *Diatrype oregonensis*.

Description. *Saprobic* on decaying branches of unidentified plant. **Sexual morph:** *Stromata* pustulate, with groups of 3–16 perithecia, rugose, visible as black, erumpent, scattered, surrounded by a thin, black line in host tissue, solitary to gregarious, 1–3 mm long and 0.5–2 mm broad (av. = 2 × 1.5 mm, n = 30), about 1 mm thick. *Endostroma* white to light yellow. *Ostiole* opening separately, papillate or apapillate, central. *Perithecium* immersed in stroma, globose to subglobose, glabrous, with cylindrical neck, brevicollous or longicollous 218.5–465 µm high, 112–257 µm diam. (av. = 306 × 164 µm, n = 10), globose to subglobose, glabrous, ostioles individual. *Peridium*: 30–50 µm thick, dark brown to hyaline with *textura angularis* cell layers. *Asci* 95–149 × 6.5–11.5 µm (av. = 120 × 10.5 µm, n = 30), 8-spored, unitunicate, clavate or cylindrical, long-stalked, apically rounded, apical rings inamyloid. *Ascospores* 11–16 × 1.5–3.5 µm (av. = 14 × 2.5 µm, n = 30), irregularly arranged, allantoid, slightly or moderately curved, subhyaline to slightly brown, smooth, aseptate, usually with two oil droplets. **Asexual morph:** undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became pale brown, dense, but thinning towards the edge, margin rough, white from above, white at margin and light brown at centre from




Figure 4. *Diatrypella pseudooregonensis* (GMB0041, holotype) **A** stromata on host substrate **B, C** stromata on host substrate **D** transverse section through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H** ostiolar canal **I, J** asci **K-N** ascospores. Scale bars: 20 µm (**G**); 10 µm (**H-N**).

below, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.

Specimens examined. CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($25^{\circ}1'19.88''N$, $99^{\circ}35'30.68''E$) on branches of an unidentified plant, 5 October 2019. Altitude: 2677 m, Y.H. Pi & Qiong Zhang, LC323 (GMB0041, *holotype*, KUN-HKAS 112646, *isotype*, ex-type living culture GMBC0041)

Additional specimens examined. CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($25^{\circ}1'13.51''N$, $99^{\circ}35'25.59''E$) on branches of an unidentified plant, 6 October 2019. Altitude: 2630 m, Y.H. Pi & Qiong Zhang, LC384 (GMB0043, KUN-HKAS 112681, living culture GMBC0043); CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($25^{\circ}1'15.00''N$, $99^{\circ}35'39.73''E$) on branches of an unidentified plant, 5 October 2019. Altitude: 2698 m, Y.H. Pi & Qiong Zhang, LC312 (GMB0040, KUN-HKAS 112674, living culture GMBC0040); CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($25^{\circ}35'19.09''N$, $99^{\circ}35'19.09''E$) on branches of an unidentified plant, 5 October 2019. Altitude: 2569 m, Y.H. Pi & Qiong Zhang, LC193 (GMB0039, KUN-HKAS 112667, living culture GMBC0039); CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($25^{\circ}1'9.11''N$, $99^{\circ}35'24.80''E$) on branches of an unidentified plant, 5 October 2019. Altitude: 2649 m, Y.H. Pi & Qiong Zhang, LC335 (GMB0042, KUN-HKAS 112647, living culture GMBC0042); CHINA, Guizhou Province, Anshun City, Pingba District ($26^{\circ}25'9.65''N$, $106^{\circ}24'24.48''E$) on branches of an unidentified plant, 1 August 2020. Altitude: 1250 m, Y.H. Pi, PB51 (GMB0044, KUN-HKAS 112693, living culture GMBC0044).

Additional sequences. GMB0041 (LSU: MW797062, RPB2: MW814906); GMB0043 (LSU: MW797064, RPB2: MW814907); GMB0040 (LSU: MW797061, RPB2: MW814905); GMB0039 (LSU: MW797059, RPB2: MW814904); GMB0042 (LSU: MW797063); GMLB0044 (LSU: MW979054, RPB2: MW814899).

Note. Morphologically, *Diatrype* has 8 ascospores in a single ascus, while *Diatrypella* has more than eight ascospores in each ascus (Senanayake et al. 2015). However, previous research (e.g. Acero et al. 2004 and Trouillas et al. 2011) suggested that both *Diatrypella* and *Diatrype* are polyphyletic within the family. In the phylogenetic analyses, *Diatrypella pseudooregonensis* grouped closely to the *D. verruciformis* and thus, we consider this new species to belong in the genus *Diatrypella*, because it is doubtful whether the number of ascospores per ascus is useful as a basis for generic classification.

Diatrypella vulgaris Trouillas, W.M. Pitt & Gubler, Fungal Diversity 49: 212 (2011)

MycoBank No: 519404

Fig. 5

Description. Saprobiic on decaying branches of an unidentified plant. **Sexual morph:** Stromata scattered on the host, 0.8–1.5 mm long and 0.8–2 mm broad (av. = 1.2 ×




Figure 5. *Diatrypella vulgaris* (GMB0051, new record for China) **A** stromata on host substrate; **B, C** close-up of stroma **D** transverse sections through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H, I** ostiolar canal **J, K** ascospores **L–O** ascospores. Scale bars: 20 µm (**G**); 10 µm (**H–I**).

1.3 mm, $n = 30$) pustulate, visible as black, rounded to irregular in shape on host surface, semi-immersed, erumpent through host bark, with 2–8 ascomata immersed in one stroma. *Endostroma* consists of outer dark brown, small, dense, thin parenchymal cells and an inner layer of white, large, loose parenchymal cells. *Ostiole* opening separately, papillate or apapillate, central 710.7–787.2 μm high, 270.2–422 μm diam. (av. = 742 \times 363 μm , $n = 10$). *Perithecium* immersed in stroma, round to oblong, with cylindrical neck, brevicollous or longicollous. *Peridium* composed of outer layer of dark brown to black, thin-walled cells, arranged in *textura angularis*, inner layer of hyaline thin-walled cells of *textura angularis*. *Asci* 111.4–152.9 \times 10.6–17.5 μm (av. = 124.5 \times 15.5 μm , $n = 30$), polysporous, clavate, long-stalked, apically rounded. *Ascospores* 8–11 \times 1–2 μm (av. = 8.9 \times 1.7 μm , $n = 30$), overlapping, crowded, allantoid, slightly or moderately curved, smooth, subhyaline, yellowish in mass, aseptate, usually with two oil droplets. **Asexual morph:** undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became pale brown, dense, but thinning towards edge, medium dense, white from above, reverse side white at margin, flesh to pale brown at centre, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.

Specimens examined. CHINA, Guizhou Province, Guiyang City, Gaopo Township (26°29'72.02"N, 106°29'55.57"E), on branches of unidentified plant, 30 October 2020. Altitude: 1589 m, S.H. Long, GP02 (GMB0051, KUN-HKAS 112697, living culture GMBC0051).

Additional sequences. GMB0051 (LSU: MW797051, RPB2: MW814897).

Note. The comparison of ITS sequences in NCBI showed that this isolate is 100% similar to the strain of *Diatrypella vulgaris* (HVGRF03), isolated from holotype specimens. Morphologically, GMB0051 shows the same features as *Diatrypella vulgaris*. The stromata of these specimens are similar, but ascospores of GMB0051 are thinner than those of the HVGRF03 (8–10 \times 2–2.5 μm) and, when compared with the ascospores of strain MFLUCC 17-0128 (4.5–7.5 \times 1–2 μm), they are shorter than GMB0051 (Trouillas et al. 2011; Hyde et al. 2017). Here, we use the ITS sequence similarity between the new collection and the type strain of *Diatrypella vulgaris* as the identification tool. *Diatrypella vulgaris* has been reported in Austria and Thailand (Trouillas et al. 2011, Hyde et al. 2017). This is the first report of *Diatrypella vulgaris* from China.

Diatrypella oregonensis (Wehm.) S.H. Long & Q.R. Li, comb. nov.

MycoBank No: 839728

≡ *Eutypella oregonensis* Wehm. Pap. Mich. Acad. Sci. 11: 163 (1930)

≡ *Diatrype oregonensis* (Wehm.) Rappaz, Mycol. helv. 2(3): 420 (1987)

Description. See Trouillas et al. (2010).

Note. The strains of *Diatrype oregonensis* (DPL200, CA117) generated from Trouillas et al. (2010) grouped in *Diatrypella s. str.* *Diatrype oregonensis* was erected

in 1930 as *Eutypella oregonensis* (Kauffman 1930). No available sequences from type material were found. After re-examination of holotype specimen of *Diatrype oregonensis*, Trouillas et al. (2010) introduced two strains of *Diatrype oregonensis* (DPL200 and CA117). Although neither of these strains are ex-type, they are, the most authoritative strains. Here, we tentatively transfer *Diatrype oregonensis* to *Diatrypella* as *Diatrypella oregonensis*, based on the phylogenetic analyses (Fig. 1). *Diatrypella oregonensis* is similar to *D. pseudooregonensis* in having 8-spored ascospores (Rappaz 1987; Trouillas et al. 2011). Nevertheless, we consider that the number of ascospores as a basis for distinguishing *Diatrypella* from *Diatrype* is not useful.

***Allodiatrype* Konta & K.D. Hyde Mycosphere 11(1): 247 (2020)**

Notes. The genus *Allodiatrype* was introduced by Konta et al. (2020), which was characterised by regular or irregular-shaped stromata, erumpent through host surface, ascospores with 8 spores and aseptate, allantoid ascospores. In this study, we introduce a new record of *Allodiatrype thailandica* (R.H. Perera et al.) Konta & K.D. Hyde collected from Yunnan Province in China.

***Allodiatrype thailandica* (R.H. Perera et al.) Konta & K.D. Hyde, Mycosphere 11(1): 253 (2020)**

Mycobank No: 556932

Fig. 6

≡ *Diatrype thailandica* R.H. Perera et al., Fungal Diversity 78: 1–237, [105] (2016)

Description. Saprobic on decaying branches of unidentified plant. **Sexual morph:** *Stromata* wart-like, pustulate, 0.5–1.8 mm long and 0.8–2.2 mm broad (av. = 1.2 × 1.3 mm, n = 30), about 1 mm thick, 1–18 in a single stroma, visible as black, rounded to irregular in shape on the host surface, erumpent through host bark, solitary to gregarious. *Endostroma* composed of an outer layer of dark brown to black, small, tightly packed, thin parenchymatous cells and an inner layer of white to yellow, large, loose parenchymal cells. *Ostiole* opening separately, papillate or apapillate, central. *Perithecium* immersed in stroma, globose to subglobose, glabrous, with cylindrical short neck, 377–447 µm high, 191–264 µm diam. (av. = 406 × 221 µm, n = 10). *Peridium* hyaline to dark brown with *textura angularis* cell layers. *Asci* 80–113.5 × 6.9–10 µm (av. = 109.3 × 8.5 µm, n = 30), 8-spored, unitunicate, clavate, long-stalked, upper part inflated, apically rounded to truncate, apical rings inamyloid. *Ascospores* 6–11 × 2–2.5 µm (av. = 8.9 × 2.3 µm, n = 30), irregularly arranged, allantoid, slightly curved, smooth, subhyaline, aseptate, usually with two oil droplets. **Asexual morph:** undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became pale yellow, irregular in shape, medium dense, flat or effuse, slightly raised, with edge fimbriate, fluffy to fairly fluffy, white




Figure 6. *Allodiatripte thailandica* (GMB0050, new record for China) **A** stromata on host substrate **B, C** close-up of stromata **D** transverse section through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H** ostiolar canal **I–K** ascospores **L–N** asci. Scale bars: 20 μm (**G**); 10 μm (**H–N**).

from above, reverse side white at margin, pale brown at centre, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.

Specimens examined. CHINA, Yunnan Province, Baoshan City, Lancang River Nature Reserve ($24^{\circ}57'25.35''N$, $99^{\circ}44'22.82''E$), on branches of unidentified plant, 2 October 2019. Altitude: 1317 m, Y.H. Pi & Qiong. Zhang, LC103 (GMB0050, KUN-HKAS 112660, living culture GMBC0050).

Additional sequences. GMB0050 (LSU: MW797052).

Note. The ITS sequence data were subjected to BLAST in NCBI and the results showed that it is 100% similar to *Allodiatripe thailandica*. Additionally, based on morphological and phylogenetic analyses, this strain was identified as the *A. thailandica*. The stromata are similar, but the ascospores of GMB0050 are longer and wider than the ascospores of strain MFLUCC 15-3662 ($3.8\text{--}6.9 \times 1\text{--}1.4 \mu\text{m}$) isolated from the holotype specimen, but it is similar to the strain MFLU 17-0735 ($6.5\text{--}10.7 \times 1.6\text{--}2.7 \mu\text{m}$) (Perera et al. 2020). Here, we use the ITS sequence similarity between the new collection and the type strain of *Allodiatripe thailandica* as basis for identification. *A. thailandica* has been reported in Thailand in 2016 as *Diatrype thailandica* and recognised as *A. thailandica* by Konta et al. (2020). This is the first report of *Allodiatripe thailandica* from China.

***Neoeutypella* M. Raza, Q.J. Shang, Phookamsak & L. Cai, Fungal Diversity 95: 167 (2019)**

Note. The genus *Neoeutypella* was introduced by Phookamsak et al. (2019) and is characterised by carbonaceous stromata immersed or semi-immersed on the host, 8-spored asci and hyaline or pale reddish-brown to brown ascospores. In this study, we introduce a new collection of *N. baoshanensis*, isolated from Guizhou Province in China.

***Neoeutypella baoshanensis* M. Raza, Q.J. Shang, Phookamsak & L. Cai, Fungal Diversity 95: 168 (2019)**

Mycobank No: 555372

Fig. 7

Description. see Phookamsak et al. (2019).

Specimens examined. CHINA, Guizhou Province, Guiyang City, Gaopo Township ($26^{\circ}29'72.37''N$, $106^{\circ}29'59.33''E$), on branches of unidentified plant, 30 November 2020. Altitude: 1589 m, S.H. Long, GP01 (GMB0052, KUN-HKAS 112696, living culture GMBC0052).

Additional sequences. GMB0052 (LSU: MW797050, RPB2: MW814896).

Note. The morphological characteristics of this specimen are consistent with those of *N. baoshanensis* a species described by Phookamsak et al. (2019). Based on phylogenetic and morphological analyses, we consider that this specimen is *Neoeutypella baoshanensis*.



Figure 7. *Neoeutypella baoshanensis* (GMB0052) **A** stromata on host substrate **B** close-up of stromata **C** transverse section through ascostroma **D** vertical section through ascostroma **E** pigments in KOH **F** culture on PDA **G** section through the ascostroma **H** ostiolar canal **I,J** ascospores **K–M** asci. Scale bars: 20 µm (**G**); 10 µm (**H–M**).

Neoeutypella baoshanensis was described as the type species of *Neoeutypella* on dead wood of *Pinus armandii* Franch. from Yunnan Province in China (Phookamsak et al. 2019). This is the first record of *N. baoshanensis* from Guizhou Province, China.

Eutypa Tul. & C. Tul.

Notes. Tulasne & Tulasne (1863) introduced the genus *Eutypa* with *Eutypa lata* as the type species. This genus includes several phytopathogens, such as *E. lata* (Pers.) Tul. & C. Tul. and *E. leptoplaca* (Durieu & Mont.) Rappaz (Moyo et al. 2017). The morphological characteristics of this genus are black, rounded to irregular-shaped stromata on the host surface, erumpent through host epidermis, solitary to gregarious, entostromatic region, consisting of white pseudoparenchymatous cells and thin black pseudoparenchymatous tissue around the white entostroma, 8-spored, spindle-shaped ascospores (Rappaz 1987; Moyo et al. 2017). We introduce a new species of *Eutypa* collected from Guizhou Province in China.

Eutypa cerasi S.H. Long & Q.R. Li, sp. nov.

Mycobank No: 839657

Fig. 8

Holotype. GMB0048.

Etymology. Refers to its host, *Prunus cerasus*.

Description. Saprobic on decaying branches of *Prunus cerasus*. **Sexual morph:** Stromata immersed in bark, covering surface of host, irregular in shape, widely effused, flat, margin diffuse, surface dark brown to black, with punctiform ostioles scattered at surface. Endostroma consists of an outer layer of black, small, dense, thin parenchymal cells and an inner layer of white, large, loose parenchymal cells. Peritheciun semi-immersed in stroma, globose to subglobose, glabrous, with cylindrical neck, brevicollous 203–304 µm high, 346–477 µm diam. (av. = 408 × 250 µm, n = 10), ovoid, obovoid to oblong. Ostiole opening separately, papillate or apapillate, central. Peridium 30–50 µm thick, dark brown to hyaline with *textura angularis* cell layers. Ascospores 83.2–120 × 5.1–8.2 µm (av. = 104.4 × 6.3 µm n = 30) 8-spored clavate, unitunicate, rounded to truncate apex, apical rings inamyloid. Ascospores 7.3–9.9 × 1.4–2 µm (av. = 8.5 × 1.7 µm, n = 30), overlapping, allantoid, slightly curved, subhyaline, smooth, aseptate, usually with oil droplets. **Asexual morph:** undetermined.

Culture characteristics. Ascospores germinating on PDA within 24 hours. Colonies on PDA, white when young, became pale yellow, irregular in shape, medium dense, flat or effuse, white from above, reverse white at margin, pale yellow at centre, no pigmentation produced on PDA medium, no conidia observed on PDA or on OA media.




Figure 8. *Eutypa cerasi* (GMB0048, holotype) **A** stromata on host substrate **B, C** close-up of stroma **D** transverse section through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H** peridium **I–K** ascospores **L–N** ascii. Scale bars: 20 µm (**G**); 10 µm (**H–N**).

Specimens examined. CHINA, Guizhou Province, Guiyang City, Aha Lake National Wetland Park ($26^{\circ}32'50.21''N$, $106^{\circ}40'15.78''E$), on branches of *Prunus cerasus*, 12 August 2020. Altitude: 1089 m, S.H. Long, AH4 (GMB0048, *holotype*, KUN-HKAS 112685, *isotype*, ex-type living culture GMBC0048).

Additional specimens examined. CHINA, Guizhou Province, Guiyang City, Aha Lake National Wetland Park ($26^{\circ}32'47.79''N$, $106^{\circ}40'21.09''E$), on branches of *Cerasus* sp., 12 August 2020. Altitude: 1089 m, S.H. Long, AH40 (GMB0049, KUN-HKAS 112683, living culture GMBC0049).

Additional sequences. GMB0048 (LSU: MW797048, RPB2: MW814894); GMB0049 (LSU: MW797049, RPB2: MW814895).

Notes. *Eutypa lata* is an important pathogen that has a wide range of hosts. However, the classification of *E. lata* is confusing because there are many variants in previous studies; now all are classified as *E. lata* (Index Fungorum 2020). Morphologically, the new collection GMB0048 has similar stromata with *Eutypa lata*, but the ascocarps of the new collection are smaller than the ascocarps (400 μm diam.) of the original description of *E. lata* (Tulasne & Tulasne, 1863). The ascocarps and ascospores of the new collection are smaller than the ascocarps (400–600 μm diam.) and ascospores (110–180 \times 5–7 μm) of the description of *E. lata* (Rappaz 1987). Additionally, in the phylogenetic analyses, *E. cerasi* is located on a branch that forms a sister clade with EP18 and RGA01 and CBS 290.87 basal to *E. cerasi*. Therefore, combining phylogenetic and morphological analyses, we introduce *Eutypa cerasi* as a new species of *Eutypa*.

Paraeutypella L.S. Dissan., J.C. Kang, Wijayaw. & K.D. Hyde.

Notes. *Paraeutypella* was introduced by Dissanayake et al. (2021) to accommodate *Paraeutypella guizhouensis* and the genus currently comprises three species. The genus is characterised by poorly developed stromata erumpent through the bark, grouped and irregularly shaped, sometimes confluent, dark brown to black, spindle-shaped, 8-spored ascospores and allantoid, overlapping, subhyaline ascospores (Trouillas et al. 2011; de Almeida et al. 2016; Dissanayake et al. 2021). In this study, we illustrate *Paraeutypella citricola* collected from Guizhou Province in China.

Paraeutypella citricola (Speg.). L.S. Dissan., Wijayaw., J.C. Kang & K.D. Hyde, in Dissanayake, Wijayawardene, Dayaratne, Samarakoon & Dai, Biodiversity Data Journal 9: e63864, 14 (2021)

Mycobank No: 228646

Fig. 9

≡ *Eutypella citricola* Speg., Anal. Mus. nac. Hist. nat. B. Aires 6: 245 (1898)




Figure 9. *Paraeutypella citricola* (GMB0053) **A** stromata on host substrate **B, C** stromata on host **D** transverse section through ascostroma **E** vertical section through ascostroma **F** culture on PDA **G** section through the ascostroma **H** ostiolar canal **I** peridium **J–K** ascospores **L–O** asci. Scale bars: 40 µm (**G**); 10 µm (**H–O**).

Table 2. The dimensions of the present species and some related species of *Diatrype* and *Allodiatriype*.

Species name	Stromata		Asci		Ascospores		Reference
	Length (mm)	Wide (mm)	Length (μm)	Wide (μm)	Length (μm)	Wide (μm)	
<i>Allodiatriype arengae</i>	0.69–0.94	0.37–0.93	54–109	6–10	7–10	2–3	Konta et al. (2020)
<i>A. elaeidicola</i>	1.2–2.8	0.9–1.66	60–91	4–7	8–10	1.5–3	Konta et al. (2020)
<i>A. elaeidis</i>	0.47–0.86	0.44–0.71	56–95	9–11	8–10	1.5–3	Konta et al. (2020)
<i>A. thailandica</i>	NA	1–2	55–80	5–7	3.8–6.9	1–1.4	Li et al. (2016)
<i>Diatrype acericola</i>	1–2	1–1.5	23–27	5–7	7.5–9	0.9–1.1	Vasilyeva and Ma (2014)
<i>D. albopruinosa</i>	0.5–1 diam.	0.5–1 diam	40–60	10–15	12–15	3.5–4	Vasilyeva and Ma (2014)
<i>D. bullata</i>	2–7 diam.	2–7 diam	25–30	5–7	7.5–9	Very thin	Vasilyeva and Ma (2014)
<i>D. disciformis</i>	NA	NA	75–115	NA	5–9	1.5–2	Senanayake et al. (2015)
<i>D. enteroxantha</i>	NA	1–3.5	18–28.5	5–9	7–10	1.5–2.5	de Almeida et al. (2016)
<i>D. hypoxyloides</i>	NA	NA	20–25	4–6	4–6	Very thin	Vasilyeva and Ma (2014)
<i>D. lancangensis</i>	NA	NA	90.5–160.5	7–15	11–18.5	2–4	This study
<i>D. lijiangensis</i>	1 diam.	1 diam	50–90	6–9	6–8	1–2	Thiyagaraja et al. (2019)
<i>D. macounii</i>	1–1.8 diam.	1–1.8 diam	25–30	4–6	4–6	0.7–1	Vasilyeva and Ma (2014)
<i>D. stigma</i>	NA	NA	25–30	5–7	6–8	1.5–2	Vasilyeva and Ma (2014)
<i>D. subundulata</i>	NA	NA	35–40	5–7	7–9	1.7–1.9	Vasilyeva and Ma (2014)
<i>D. undulata</i>	NA	NA	25–30	3.5–4.5	5–7	0.9–1.3	Vasilyeva and Ma (2014)
<i>D. whitmanensis</i>	NA	NA	50–82	8–15	7.5–10	1–1.5	Trouilla et al. 2010
<i>Pseudodiatrype hainanensis</i>	2–3.6	1.6–3	110–155.5	6–10	8.5–13	1.5–2.5	This study

Newly identified taxa are indicated in bold, NA: No description available.

Description. For description, see Dissanayake et al. (2021)

Specimens examined. CHINA, Guizhou Province, Guiyang City: Aha Lake National Wetland Park (26°20'37.28"N, 108°21'4.34"E), on branches of unidentified plant, 30 August 2020. Altitude: 802 m, S.H. Long, LGS147 (GMB0053, KUN-HKAS 112704, living culture GMBC0053).

Additional sequences. GMB0053 (LSU: 797053, RPB2: MW814898).

Notes. The ITS sequence data were compared by using NCBI and the result showed that it is 100% similar to the ex-type strain (HVVIT07) of *P. citricola*. The morphological features of the new collection are consistent with those described by Dissanayake et al. (2021). This collection is identified as a *P. citricolca*, based on morphological and molecular data.

Discussion

In this study, one new genus, three new species, two new records from China, a novel combination and two known species were reported from karst areas of China. We used molecular data to delimit the species of Diatrypaceae. The new genus *Pseudodiatrype* is morphologically similar to *Allodiatriype* and *Diatrype*, but distinct in the size of stromata, number of ascocarps and colour of endostroma; it also formed a distinct branch in the phylogenetic analyses (Fig. 1). *Diatrype oregonensis* was transferred to *Diatrypella oregonensis* based on the phylogenetic analyses. Based on phylogenetic analyses, *Diatrypella pseudooregonensis* was introduced as an 8-spored species of *Diatrypella*.

Our phylogenetic analyses, based on ITS and β -tubulin, agree with the previous studies (Acero et al. 2004; Trouillas et al. 2011; Mehrabi et al. 2015, 2016; de Almeida et al. 2016; Shang et al. 2017; Dissanayake et al. 2021; Zhu et al. 2021). However, several genera are not monophyletic; for example, *Cryptosphaeria*, *Diatrype*, *Diatrypella*, and *Eutypa*. The identification of species of Diatrypaceae has been a problem due to the polyphyletic generic concepts based on the features of the stromata in early research (Fries 1823). Recently, new approaches have been proposed for classifying Diatrypaceae. Acero et al. (2004) proposed to classify them by ITS sequence-based phylogenetic analyses, while Carmarán et al. (2006) suggested that the identification should be based on the morphology of the asci. However, due to the lack of type specimens, the lack of β -tubulin sequence and polyphyletic origins have resulted in molecular data that correlate poorly with morphological criteria used to delineate genera and species within the Diatrypaceae (Acero et al. 2004). Moreover, Acero et al. (2004) has mentioned that *Diatrypella quercina* should be placed in the genus *Diatrype* despite its polysporous ascii since the molecular data placed *Diatrypella quercina* in the branch of the genus *Diatrype*.

Diatrype and *Diatrypella* have morphologically similar verruculose stromata and allantoid ascospores and the polysporous or 8-spored ascus serve as a basis for distinguishing the two genera. However, in phylogenetic analyses, species of these two genera overlap. In this study, we used the phylogenetic analyses as the main basis for classification following Vasilyeva and Stephenson (2005) and Liu et al. (2015). Clade 1 contains *Diatrypella verruciformis* which is the type species of *Diatrypella*, of which *Diatrypella pseudooregonensis*, *Diatrypella oregonensis* have 8-spored, and other species in clade 1 have polyspored ascus. Clade 12 contains the *Diatrype* type species *Diatrype disciformis*, of which *Diatrype iranensis* and *Diatrype macrospora* have polyspored ascus, and other species in clade 12 have 8-spored ascus. Hence, we concluded that the number of ascospores in each ascus cannot be used as a criterion for distinguishing *Diatrypella* from *Diatrype*.

The phylogenetic tree shows that the classification of Diatrypaceae is confusing. Members of *Diatrypella* (*D. favacea*, *D. hubeiensis*, *D. pulvinata* and *D. yunnanensis*) cluster with *Diatrype palmicola* and *Diatrype lancangensis*. Maybe this clade should be identified as a new genus. We will discuss its classification status after more strains, more gene sequences and new taxonomic features are collected. Some species of *Diatrypella* (*D. iranensis* and *D. macrospora*) which have polysporous ascus are placed between species of *Diatrype*, and they are transferred to *Diatrype iranensis* and *Diatrype macrospora* by Zhu et al. (Zhu et al. 2021). *Diatrype enteroxantha* is often derived from the sister clade of *Allodiatrype* rather than the *Diatrype* clade. Additionally, *Eutypa microasca* (BAFC51550) clusters with *Peroneutypa* species (Clade 17). The above-mentioned confusion also showed in the original publication and other recent studies (Grassi et al. 2014; Mehrabi et al. 2016; Shang et al. 2018; Hyde et al. 2019; Phookamsak et al. 2019; Konta et al. 2020). Therefore, addressing the taxonomic confusion of this family requires a re-examination of older taxa, based on morphological studies, epitypification and multi-gene phylogenetic analyses (Ariyawansa et al. 2014).

Acknowledgements

This research was supported by the National Natural Science Foundation of China (32000009 and 31960005); the Fund of the Science and Technology Foundation of Guizhou Province ([2020]1Y059); Guiyang Science and Technology Planning Project No. (2017)30-19; Guizhou Province Ordinary Colleges and Universities Youth Science and Technology Talent Growth Project [2021]154. Nalin N. Wijayawardene would like to thank the National Natural Science Foundation of China (No. NSFC 31950410558), the State Key Laboratory of Functions and Applications of Medicinal Plants, Guizhou Medicinal University (No. FAMP201906K) and High-Level Talent Recruitment Plan of Yunnan Provinces (“Young Talents” Program and “High-End Foreign Experts” Program); the Fund of High-Level Innovation Talents [No. 2015-4029], the Base of International Scientific and Technological Cooperation of Guizhou Province [No. [2017]5802].

Reference

- Acero FJ, González V, Ballesteros JS, Rubio V, Checa J, Bills GF, Salazar O, Platas G, Peláez F (2004) Molecular phylogenetic studies on the Diatrypaceae based on rDNA-ITS sequences. *Mycologia* 96: 249–259. <https://doi.org/10.1080/15572536.2005.11832975>
- Arhipova N, Gaitnieks T, Donis J, Stenlid J, Vasaitis R (2012) Heart-rot and associated fungi in *Alnus glutinosa* stands in Latvia. *Scandinavian Journal of Forest Research* 27: 327–336. <https://doi.org/10.1080/02827581.2012.670727>
- Ariyawansa HA, Hawksworth DL, Hyde KD, Jones EGB, Maharachchikumbura SSN, Manamgoda DS, Thambugala KM, Udayanga D, Camporesi E, Daranagama A, Jayawardena R, Liu JK, Mckenzie EHC, Phookamsak R, Senanayake IC, Shivas RG, Tian Q, Xu JC (2014) Epitypification and neotypification: guidelines with appropriate and inappropriate examples. *Fungal Diversity* 69: 57–91. <https://doi.org/10.1007/s13225-014-0315-4>
- Carmarán CC, Romero AI, Giussani LM (2006) An approach towards a new phylogenetic classification in Diatrypaceae. *Fungal Diversity* 23: 67–87.
- Chomnunti P, Hongsanan S, Hudson BA, Tian Q, Peršoh D, Dhami MK, Alias AS, Xu JC, Liu XZ, Stadler M, Hyde KD (2014) The sooty moulds. *Fungal Divers* 66: 1–36. <https://doi.org/10.1007/s13225-014-0278-5>
- Crous PW, Wingfield MJ, Guarro J, Cheewangkoon R, van der Bank M, Swart WJ, Stchigel AM, Cano-Lira JF, Roux J, Madrid H, Damm U, Wood AR, Shuttleworth LA, Hodges CS, Munster M, de Jesús Yáñez-Morales M, Zúñiga-Estrada L, Cruywagen EM, De Hoog GS, Silvera C, Najafzadeh J, Davison EM, Davison PJN, Barrett MD, Barrett RL, Manamgoda DS, Minnis AM, Kleczewski NM, Flory SL, Castlebury LA, Clay K, Hyde KD, Maúisse-Sitoe SND, Chen S, Lechat C, Hairaud M, Lesage-Meessen L, Pawłowska J, Wilk M, Śliwińska-Wyrzychowska A, Mętrak M, Wrzosek M, Pavlic-Zupanc D, Maleme HM, Slippers B, Mac Cormack WP, Archuby DI, Grünwald NJ, Tellería MT, Dueñas M, Martín MP, Marincowitz S, de Beer ZW, Perez CA, Gené J, Marin-Felix Y, Groenewald JZ

- (2013) Fungal Planet description sheets: 154–213. Persoonia: Molecular Phylogeny and Evolution of Fungi 31: 1–188. <https://doi.org/10.3767/003158513X675925>
- Dayarathne MC, Phookamsak R, Hyde KD, Manawasinghe IS, Toanun C, Jones EBG (2016) *Halodiatriype*, a novel diatrypaceous genus from mangroves with *H. salinicola* and *H. avicenniae* spp. nov. Mycosphere 7: 612–627. <https://doi.org/10.5943/mycosphere/7/5/7>
- Dayarathne MC, Wanasinghe DN, Devadatha B, Abeywickrama P, Jones EBG, Chomnunti P, Sarma VV, Hyde KD, Lumyong S, McKenzie EHC (2020) Modern taxonomic approaches to identifying diatrypaceous fungi from marine habitats, with a Novel Genus *Halocryptovalsa* Dayarathne & K.D.Hyde, Gen. Nov. Cryptogamie Mycologie, 41: 21–67. <https://doi.org/10.5252/cryptogamie-mycologie2020v41a3>
- de Almeida DAC, Gusmão LFP, Miller AN (2016) Taxonomy and molecular phylogeny of Diatrypaceae (Ascomycota, Xylariales) species from the Brazilian semi-arid region, including four new species. Mycological Progress 15: 1–27.
- Dissanayake LS, Wijayawardene NN, Dayarathne MC, Samarakoon MC, Dai DQ, Hyde KD, Kang JC (2021) *Paraeutypella guizhouensis* gen. et sp. nov. and *Diatrypella longiasca* sp. nov. (Diatrypaceae) from China. Biodiversity Data Journal 9: e63864. <https://doi.org/10.3897/BDJ.9.e63864>
- Dong D, Fang C, Zhao W, Xie Z (2002) Evaluation of geochemical quality control in determination of Mn in soils using a sequential chemical extraction. Chinese Geographical Science 12: 166–170. <https://doi.org/10.1007/s11769-002-0026-8>
- Fries EM (1823) Systema Mycologicum 3: 1–202.
- Gao LL, Zhang Q, Sun XY, Jiang L, Zhang R, Sun GY, Zha YL, Biggs AR (2013) Etiology of moldy core, core browning, and core rot of fuji apple in China. Plant Disease 97: 510–516. <https://doi.org/10.1094/PDIS-01-12-0024-RE>
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. Applied & Environmental Microbiology 61: 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Glawe DA, Rogers JD (1984) Diatrypaceae in the Pacific Northwest. Mycotaxon 20: 401–460.
- Glez-Peña D, Gómez-Blanco D, Reboiro-Jato M, Fdez-Riverola F, David P (2010) ALTER: programoriented conversion of DNA and protein alignments. Nucleic Acids Research 38: 14–18. <https://doi.org/10.1093/nar/gkq321>
- Grassi E, Belen Pildain M, Levin L, Carmaran C (2014) Studies in Diatrypaceae: the new species *Eutypa microasca* and investigation of ligninolytic enzyme production. Sydowia 66: 99–114.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.
- Hyde KD, Norphanphoun C, Maharanachchikumbura SSN, Bhat DJ, Jones EBG, Bundhun D, Chen YJ, Bao DF, Boonmee S, Calabon MS, Chaiwan N, Chethana KWT, Dai DQ, Dayarathne MC, Devadatha B, Dissanayake AJ, Dissanayake LS, Doilom M, Dong W, Fan XL, Goonasekara ID, Hongsanan S, Huang SK, Jayawardena RS, Jeewon R, Karunaratna A, Konta S, Kumar V, Lin CG, Liu JK, Liu NG, Luangsa-ard J, Lumyong S, Luo ZL, Marasinghe DS, McKenzie EHC, Niego AGT, Niranjan M, Perera RH, Phukhamsakda C, Rathnayaka AR, Samarakoon MC, Samarakoon SMBC, Sarma VV, Senanayake IC,

- Shang QJ, Stadler M, Tibpromma S, Wanasinghe DN, Wei DP, Wijayawardene NN, Xiao YP, Yang J, Zeng XY, Zhang SN, Xiang MM (2020a) Refined families of Sordariomycetes. *Mycosphere* 11: 305–1059. <https://doi.org/10.5943/mycosphere/11/1/7>
- Hyde KD, Dong Y, Phookamsak R, Jeewon R, Bhat DJ, Jones EBG, Liu NG, Abeywickrama PD, Mapook A, Wei D, Perera RH, Manawasinghe IS, Pem D, Bundhun D, Karunaratna A, Ekanayaka AH, Bao DF, Li JF, Samarakoon MC, Chaiwan N, Lin CG, Phutthacharoen K, Zhang SN, Senanayake IC, Goonasekara ID, Thambugala KM, Phukhamsakda C, Tennakoon DS, Jiang HB, Yang J, Zeng M, Huanraluek N, Liu JK, Wijesinghe SN, Tian Q, Tibpromma S, Brahmanage RS, Boonmee S, Huang SK, Thiyagaraja V, Lu YZ, Jayawardena RS, Dong W, Yang EF, Singh SK, Singh SM, Rana S, Lad SS, Anand G, Devadatha B, Nirajan M, Sarma VV, Liimatainen K, Hudson BA, Niskanen T, Overall A, Alvarenga RLM, Gibertoni TB, Pfleigler WP, Horváth E, Imre A, Alves AL, da Silva Santos AC, Tiago PV, Bulgakov TS, Wanasinghe DN, Bahkali AH, Doilom M, Elgorban AM, Maharachchikumbura SSN, Rajeshkumar KC, Haelewaters D, Mortimer PE, Zhao Q, Lumyong S, Xu JC, Sheng J (2020b) Fungal diversity notes 1151–1276: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal diversity* 100: 5–277. <https://doi.org/10.1007/s13225-020-00439-5>
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Thilini Chethana KW, Clericuzio M, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, He MQ, Hongsanan S, Huang SK, Jayasiri SC, Jayawardena RS, Karunaratna A, Konta S, Kušan I, Lee H, Li JF, Lin CG, Liu NG, Lu YZ, Luo ZL, Manawasinghe IS, Mapook A, Perera RH, Phookamsak R, Phukhamsakda C, Siedlecki I, Soares AM, Tennakoon DS, Tian Q, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Abdel-Aziz FA, Li WJ, Senanayake IC, Shang QJ, Daranagama DA, de Silva NI, Thambugala KM, Abdel-Wahab MA, Bahkali AH, Berbee ML, Boonmee S, Bhat DJ, Bulgakov TS, Buyck B, Camporesi E, Castañeda-Ruiz RF, Chomnunti P, Doilom M, Dovana F, Gibertoni TB, Jadan M, Jeewon R, Jones EBG, Kang JC, Karunaratna SC, Lim YW, Liu JK, Liu ZY, Plautz Jr HL, Lumyong S, Maharachchikumbura SSN, Matočec N, McKenzie EHC, Mešić A, Miller D, Pawłowska J, Pereira OL, Promputtha I, Romero AL, Ryvarden L, Su HY, Suetrong S, Tkalčec Z, Vizzini A, Wen TC, Wisittrassameewong K, Wrzosek M, Xu JC, Zhao Q, Zhao RL, Mortimer PE (2017) Fungal diversity notes 603–708: taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235. <https://doi.org/10.1007/s13225-017-0391-3>
- Hyde KD, Tennakoon DS, Jeewon R, Bhat DJ, Maharachchikumbura SSN, Rossi W, Leonardi M, Lee HB, Mun HY, Houbraken J, Nguyen TTT, Jeon SJ, Frisvad JC, Dhanushka N, Wanasinghe DN, Luücking R, Aptroot A, Cáceres MES, Karunaratna SC, Hongsanan S, Phookamsak R, de Silva NI, Thambugala KM, Jayawardena RS, Senanayake IC, Boonmee S, Chen J, Luo ZL, Phukhamsakda C, Pereira OL, Abreu VP, Rosado AWC, Bart B, Randlerianjohany E, Hofstetter V, Gibertoni TB, da Silva Soares AM, Plautz Jr HL, Sotão HMP, Xavier WKS, Bezerra JDP, de Oliveira TGL, de Souza-Motta CM, Magalhães OMC, Bundhun D, Harishchandra D, Manawasinghe IS, Dong W, Zhang SN, Bao DF, Samarakoon MC, Pem D, Karunaratna A, Lin CG, Yang J, Perera RH, Kumar V, Huang SK, Dayarathne MC, Ekanayaka AH, Jayasiri SC, Xiao YP, Konta S, Niskanen T, Liimatainen K, Dai YC, Ji XH, Tian XM, Mešić A, Singh SK, Phutthacharoen K, Cai L, Sorvongxay T,

- Thiyagaraja V, Norphanphoun C, Chaiwan N, Lu YZ, Jiang HB, Zhang JF, Abeywickrama PD, Aluthmuhandiram JVS, Brahmanage RS, Zeng M, Chethana T, Wei DP, Réblová M, Fournier J, Nekvindová J, do Nascimento Barbosa R, dos Santos JEF, de Oliveira NT, Li GJ, Ertz D, Shang QJ, Phillips AJL, Kuo CH, Camporesi E, Bulgakov TS, Lumyong S, Jones EBG, Chomnunti P, Gentekaki E, Bungartz F, Zeng XY, Fryar S, Tkalčec Z, Liang J, Li GS, Wen TC, Singh PN, Gafforov Y, Promputtha I, Yasanthika E, Goonasekara ID, Zhao RL, Zhao Q, Kirk PM, Liu JK, Yan JY, Mortimer PE, Xu JC (2019) Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 96: 1–242. <https://doi.org/10.1007/s13225-019-00429-2>
- Index Fungorum (2020) Index Fungorum <http://www.indexfungorum.org/Names/Names.asp>
- Jurc D, Ogris N, Slippers B, Stenlid J (2006) First report of *Eutypella* canker of *Acer pseudoplatanus* in Europe. *Plant Pathology* 55: 577–577. <https://doi.org/10.1111/j.1365-3059.2006.01426.x>
- Kauffman CH (1930) The fungous flora of Siskiyou Mountains in southern Oregon. *Papers of the Michigan Academy of Science, Arts, and Letters* 11: 151–210.
- Klaysuban A, Sakayaroj J, Jones EG (2014) An additional marine fungal lineage in the Diatrypaceae, Xylariales: *Pedumispora rhizophorae*. *Botanica marina* 57: 413–420. <https://doi.org/10.1515/bot-2014-0017>
- Konta S, Maharachchikumbura SSN, Senanayake IC, McKenzie EHC, Stadler M, Boonmee S, Phookamsak R, Jayawardena RS, Senwanna C, Hyde KD, Elgorban AM, Eungwanichayapant PD (2020) A new genus *Allodiatripe*, five new species and a new host record of diatrypaceous fungi from palms (Arecaceae). *Mycosphere* 11: 239–268. <https://doi.org/10.5943/mycosphere/11/1/7>
- Li GJ, Hyde KD, Zhao RN, Hongsanan S, Abdel-Aziz FA, Abdel-Wahab MA, Alvarado P, Alves-Silva G, Ammirati JF, Ariyawansa HA, Baghela A, Bahkali AH, Beug M, Bhat DJ, Bojantchev D, Boonpratuang T, Bulgakov TS, Camporesi E, Boro MC, Ceska O, Chakraborty D, Chen JJ, Chethana KWT, Chomnunti P, Consiglio G, Cui BK, Dai DQ, Dai YC, Daranagama DA, Das K, Dayaratne MC, Crop ED, De Oliveira RJV, De Souza CAF, De Souza JI, Dentinger BTM, Dissanayake AJ, Doilom M, Drechsler-Santos ER, Ghobad-Nejhad M, Gilmore SP, Góes-Neto A, Gorczak M, Haitjema GH, Hapuarachchi KK, Hashimoto A, He MQ, Henske JK, Hirayama K, Iribarren MJ, Jayasiri SC, Jayawardena RS, Jeon SJ, Jerônimo GH, Jesus AL, Jones EBG, Kang JC, Karunaratne SC, Kirk PM, Konta S, Kuhnert E, Langer E, Lee HS, Lee HB, Li WJ, Li XH, Liimatainen K, Lima DX, Lin CG, Liu JK, Liu XZ, Liu ZY, Luangsa-Ard JJ, Lücking R, Lumbsch HT, Lumyong S, Leaño EM, Marano AV, Matsumura M, McKenzie EHC, Mongkolsamrit S, Mortimer PE, Nguyen TTT, Niskanen T, Norphanphoun C, O'malley MA, Parnmen S, Pawłowska J, Perera RH, Phookamsak R, Phukhamsakda C, PiresZottarelli CLA, Raspé O, Reck MA, Rocha SCO, De Santiago ALCMA, Senanayake IC, Setti L, Shang QJ, Singh SK, Sir EB, Solomon KV, Song J, Srikitkulchai P, Stadler M, Suetrong S, Takahashi H, Takahashi T, Tanaka K, Tang LP, Thambugala KM, Thanakitpipattana D, Theodorou MK, Thongbai B, Thummarukcharoen T, Tian Q, Tibpromma S, Verbeken A, Vizini A, Vlasák J, Voigt K, Wanasinghe DN, Wang Y, Weerakoon G, Wen HA, Wen TC, Wijayawardene NN, Wongkanoun S, Wrzosek M, Xiao YP, Xu JC, Yan JY, Yang J, Yang SD, Hu Y, Zhang

- JF, Zhao J, Zhou LW, Peršoh D, Phillips AJL, Maharachchikumbura SSN (2016) Fungal Divers notes 253–366: taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 78: 1–237. <https://doi.org/10.1007/s13225-016-0366-9>
- Liu JK, Hyde KD, Gareth EBG, Ariyawansa HA, Bhat DJ, Boonmee S, Maharachchikumbura SS, Mckenzie EH, Phookamsak R, Phukhamsakda C, Shenoy BD, AbdelWahab MA, Buyck B, Chen J, Chethana KWT, Singtripop C, Dai DQ, Dai YC, Daranagama DA, Dis-sanayake AJ, Doilom M, D'souza MJ, Fan XL, Goonasekara ID, Hirayama K, Hongsanan S, Jayasiri SC, Jayawardena RS, Karunaratna SC, Li WJ, Mapook A, Norphanphoun C, Pang KL, Perera RH, Persoh D, Pinruan U, Senanayake IC, Somrithipol S, Suetrong S, Tanaka K, Thambugala KM, Tian Q, Tibpromma S, Udayanga D, Wijayawardene NN, Wanasinghe D, Wisitrassameewong K, Zeng XY, Abdel-Aziz FA, Adamcík S, Bahkali AH, Boonyuen N, Bulgakov T, Callac P, Chomnunti P, Greiner K, Hashimoto A, Hofstetter V, Kang JC, Xing DL, Li H, Liu XZ, Liu ZY, Matsumura M, Mortimer PE, Rambold G, Randrianjohany E, Sato G, Sri-Indrasutdh V, Tian CM, Verbeken A, Brackel W, Wang Y, Wen TC, Xu JC, Yan JY, Zhao RL, Camporesi, E (2015) Fungal diversity notes 1–110: taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72: 1–197. <https://doi.org/10.1007/s13225-015-0324-y>
- Luque J, Garcia-Figueroes F, Legorburu FJ, Muruamendiaraz A, Armengol J, Trouillas FP (2012) Species of Diatrypaceae associated with grapevine trunk diseases in Eastern Spain. *Phytopathologia Mediterranea* 51: 528–540. https://doi.org/10.14601/Phytopathol_Mediterr-9953
- Lynch SC, Eskalen A, Zambino PJ, Mayorquin JS, Wang DH (2013) Identification and pathogenicity of Botryosphaeriaceae species associated with coast live oak (*Quercus agrifolia*) decline in southern California. *Mycologia* 105: 125–140. <https://doi.org/10.3852/12-047>
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EHC, Bhat JD, Dayarathne MC, Huang SK, Norphanphoun C, Senanayake IC, Perera RH, Shang QJ, Xiao Y, D'souza MJ, Hongsanan S, Jayawardena RS, Daranagama DA, Konta S, Goonasekara ID, Zhuang WY, Jeewon R, Phillips AJL, Abdel-Wahab MA, Al-Sadi AM, Bahkali AH, Boonmee S, Boonyuen N, Cheewangkoon R, Dissanayake AJ, Kang J, Li QR, Liu JK, Liu XZ, Liu ZY, Luangsa-Ard JJ, Pang KL, Phookamsak R, Promputtha I, Suetrong S, Stadler M, Wen T, Wijayawardene NN (2016) Families of Sordariomycetes. *Fungal Diversity* 79: 1–317. <https://doi.org/10.1007/s13225-016-0369-6>
- Mehrabi M, Asgari B, Hemmati R (2019) Two new species of *Eutypella* and a new combination in the genus *Peroneutypa* (Diatrypaceae). *Mycological Progress* 18: 1057–1069. <https://doi.org/10.1007/s11557-019-01503-4>
- Mehrabi M, Hemmati R, Vasilyeva LN, Trouillas FP (2015) A new species and a new record of Diatrypaceae from Iran. *Mycosphere* 6: 60–68. <https://doi.org/10.5943/mycosphere/6/1/7>
- Mehrabi M, Hemmati R, Vasilyeva LN, Trouillas FP (2016) *Diatrypella macrospora* sp. nov. and new records of diatrypaceous fungi from Iran. *Phytotaxa* 252: 43–55. <https://doi.org/10.2307/3762061>
- Miao Q, Wang Y, Li J, Yuan S, Shi L, Gu X (2007) Study on the spring drought rule in the karst region of yunnan and guizhou plateau in china. *International Society for Optics and Photonics* 6790: 67903Z. <https://doi.org/10.1117/12.746860>

- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Gateway Computing Environments Workshop 2010 (GCE), New Orleans, Louisiana, November 2010: 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Moyo P, Damm U, Mostert L, Halleen F (2018a) *Eutypa*, *Eutypella*, and *Cryptovalsa* Species (Diatrypaceae) associated with *Prunus* species in South Africa. *Plant Disease* 102: 1402–1409. <https://doi.org/10.1094/PDIS-11-17-1696-RE>
- Moyo P, Mostert L, Spies CF, Damm U, Hallen F (2018b) Diversity of Diatrypaceae species associated with dieback of grapevines in South Africa, with the description of *Eutypa cremea* sp. nov. *Plant Disease* 102: 220–230. <https://doi.org/10.1094/PDIS-05-17-0738-RE>
- Nylander JAA (2004) MrModeltest v2.2. Program distributed by the author: 2. Evolutionary Biology Centre, Uppsala University 1–2.
- O'Donnell K, Cigelnik E (1997) Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* 7: 103–116. <https://doi.org/10.1006/mpev.1996.0376>
- Paolinelli-Alfonso M, Serrano-Gomez C, Hernandez-Martinez R (2015) Occurrence of *Eutypella microtheca* in grapevine cankers in Mexico. *Phytopathologia Mediterranea* 54: 86–93. https://doi.org/10.14601/Phytopathol_Mediterr-14998
- Perera RH, Hyde KD, Maharachchikumbura S, Jones EBG, McKenzie EHC, Stadler M, Lee HB, Samarakoon MC, Ekanayaka AH, Camporesi E, Liu JK, Liu ZY (2020) Fungi on wild seeds and fruits. *mycosphere* 11: 2108–2480. <https://doi.org/10.5943/mycosphere/11/1/14>
- Peršoh D, Melcher M, Graf K, Fournier J, Stadler M, Rmbold G (2009) Molecular and morphological evidence for the delimitation of *Xylaria hypoxylon*. *Mycologia* 101: 256–268. <https://doi.org/10.3852/08-108>
- Phookamsak R, Hyde KD, Jeewon R, Bhat DJ, Jones EBG, Maharachchikumbura SSN, Raspé O, Karunarathna SC, Wanasinghe DN, Hongsanan S, Doilom M, Tennakoon DS, Machado AR, Firmino AL, Ghosh A, Karunarathna A, Mešić A, Dutta AK, Thongbai B, Devadatha B, Norphanphoun C, Senwanna C, Wei DP, Pem D, Ackah FK, Wang GN, Jiang HB, Madrid H, Lee HB, Goonasekara ID, Manawasinghe IS, Kušan I, Cano J, Gené J, Li JF, Das K, Acharya K, Anil Raj KN, Deepna Latha KP, Thilini Chethana KW, He MQ, Dueñas M, Jadan M, Martín MP, Samarakoon MC, Dayarathne MC, Raza M, Park MS, Teresa Telleria M, Chaiwan N, Matočec N, de Silva NO, Pereira OL, Singh PN, Manimohan P, Uniyal P, Shang QJ, Bhatt RP, Perera RH, Alvarenga RLM, Nogal-Prata S, Singh SK, Vadhanarat S, Oh SY, Huang SK, Rana S, Konta S, Paloi S, Jayasiri SC, Jeon JS, Mehmood T, Gibertoni TB, Nguyen TT, Singh U, Thiagaraja V, Sarma VV, Dong W, Yu XD, Lu YZ, Lim YW, Chen Y, Tkalc̆ec Z, Zhang ZF, Luo ZL, Daranagama DA, Thambugala KM, Tibpromma S, Camporesi E, Bulgakov TS, Dissanayake AJ, Senanayake IC, Dai DQ, Tang LZ, Khan S, Zhang H, Promputtha I, Cai L, Chomnunti P, Zhao RL, Lumyong S, Boonmee S, Wen TC, Mortimer PE, Xu JC (2019) Fungal diversity notes 929–1036: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 95: 1–273. <https://doi.org/10.1007/s13225-019-00421-w>
- Phukhamsakda C, McKenzie EHC, Phillips AJL, Gareth Jones EB, Jayarama Bhat D, Stadler M, Bhunjun CS, Wanasinghe DN, Thongbai B, Camporesi E, Ertz D, Jayawardena RS,

- Perera RH, Ekanayake AH, Tibpromma S, Doilom M, Xu J, Hyde KD (2020) Microfungi associated with *Clematis* (Ranunculaceae) with an integrated approach to delimiting species boundaries. *Fungal Diversity* 102: 1–203. <https://doi.org/10.1007/s13225-020-00448-4>
- Rambaut A (2012) FigTree: Tree Figure Drawing Tool Version 1.4.0 2006–2012, Institute of Evolutionary Biology, University of Edinburgh. <http://tree.bio.ed.ac.uk/software/figtree/>
- Rappaz F (1987) Taxonomy and nomenclature of the octosporous Diatrypaceae. *Mycologia Helvetica* 2: 285–648.
- Rolshausen PE, Mahoney NE, Molyneux RJ, Gubler WD (2006) A reassessment of the species concept in *Eutypa lata*, the causal agent of *Eutypa* dieback of grapevine. *Phytopathology* 96: 369–377. <https://doi.org/10.1094/PHYTO-96-0369>
- Senanayake IC, Maharachchikumbura SN, Hyde KD, Bhat JD, Jones EG, Mckenzie EH, Dai DQ, Daranagama DA, Dayarathne MC, Goonasekara ID, Konta S, Li WJ, Shang QJ, Stadler M, Wijayawardene NN, Xiao YP, Norphanphoun C, Li Q, Liu XY, Bahkali AH, Kang JC, Wang Y, Wen TC, Wendt I, Xu JC, Camporesi E (2015) Towards unraveling relationships in Xylariomycetidae (Sordariomycetes). *Fungal Diversity* 73: 73–144. <https://doi.org/10.1007/s13225-015-0340-y>
- Senwanna C, Phookamsak R, Doilom M, Hyde KD, Cheewangkoon R (2017) Novel taxa of Diatrypaceae from Para rubber (*Hevea brasiliensis*) in northern Thailand; introducing a novel genus *Allocryptovalsa*. *Mycosphere* 8: 1835–1855. <https://doi.org/10.5943/mycosphere/8/10/9>
- Shang QJ, Hyde KD, Jeewon R, Khan S, Promputtha I, Phookamsak R (2018) Morpho-molecular characterization of *Peroneutypa* (Diatrypaceae, Xylariales) with two novel species from Thailand. *Phytotaxa* 356: 1–18. <https://doi.org/10.1080/15572536.2005.11832975>
- Shang QJ, Hyde KD, Phookamsak R, Doilom M, Bhat DJ, Maharachchikumbura SS, Promputtha I (2017) *Diatrypella tectonae* and *Peroneutypa mackenziei* spp. nov. (Diatrypaceae) from northern Thailand. *Mycological progress* 16: 463–476. <https://doi.org/10.1007/s11557-017-1294-0>
- Thiyagaraja V, Senanayake IC, Wanasinghe DN, Karunaratna SC, Worthy FR, To-Anun C (2019) Phylogenetic and morphological appraisal of *Diatrype lijiangensis* sp. nov. (Diatrypaceae, Xylariales) from China. *Asian Journal of Mycology* 2: 198–208. <https://doi.org/10.5943/ajom/2/1/10>
- Trouillas FP, Gubler WD (2004) Identification and characterization of *Eutypa leptoplaca*, a new pathogen of grapevine in Northern California. *Mycological Research* 108: 1195–1204. <https://doi.org/10.1017/S0953756204000863>
- Trouillas FP, Urbez-Torres JR, Gubler WD (2010) Diversity of diatrypaceous fungi associated with grapevine canker diseases in California. *Mycologia* 102: 319–336. <https://doi.org/10.3852/08-185>
- Trouillas FP, Hand FP, Inderbitzin P, Gubler WD (2015) The genus *Cryptosphaeria* in the western United States: taxonomy, multilocus phylogeny and a new species, *C. multicontinentalis*. *Mycologia* 107: 1304–1313. <https://doi.org/10.3852/15-115>
- Trouillas FP, Wayne MP, Sosnowski MR, Huang R, Peduto F, Loschiavo A, Savocchia S, Scott ES, Gubler WD (2011) Taxonomy and DNA phylogeny of Diatrypaceae associated with *Vitis vinifera* and other woody plants in Australia. *Fungal Diversity* 49: 203–223. <https://doi.org/10.1007/s13225-011-0094-0>

- Tulasne L-R, Tulasne C (1863) Selecta Fungorum carpologia, Paris, 2, 56.
- U'ren JM, Miadlikowska J, Zimmerman NB, Ltzoni F, Stajich JE, Arnold AE (2016) Contributions of North American endophytes to the phylogeny, ecology, and taxonomy of Xylariaceae (Sordariomycetes, Ascomycota). Molecular Phylogenetics and Evolution 98: 210–232. <https://doi.org/10.1016/j.ympev.2016.02.010>
- Úrbez-Torres JR, Adams P, Kamas J, Gubler WD (2009) Identification, incidence, and pathogenicity of fungal species associated with grapevine dieback in Texas. American Journal of Enology and Viticulture 60: 497–507.
- Úrbez-torres JR, Peduto F, Striegler RK, Urrearomero KE, Rupe JC, Cartwright RD, Gubler WD (2012) Characterization of fungal pathogens associated with grapevine trunk diseases in Arkansas and Missouri. Fungal Diversity 52: 169–189. <https://doi.org/10.1007/s13225-011-0110-4>
- Vasilyeva LN, Ma HX (2014) Diatrypaceous fungi in north-eastern China. 1. *Cryptosphaeria* and *Diatrype*. Phytotaxa 186(5): 261–270. <https://doi.org/10.11646/phytotaxa.186.5.3>
- Vasilyeva LN, Stephenson SL (2005) Pyrenomycetes of the Great Smoky Mountains National Park. II. *Cryptovalsa* Ces. et De Not. and *Diatrypella* (Ces. et De Not.) Nitschke (Diatrypaceae). Fungal Diversity 19: 189–200.
- Vieira MLA, Hughes AFS, Gil VB, Vaz AB, Alves TM, Zani CL, Rosa CA, Rosa LH (2011) Diversity and antimicrobial activities of the fungal endophyte community associated with the traditional Brazilian medicinal plant Solanum cernuum Vell. (Solanaceae). Canadian Journal of Microbiology 58: 54–56. <https://doi.org/10.1139/w11-105>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- White TJ, Bruns T, Lee SJWT, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, Lumbsch HT, Kukwa M, Issi IV, Madrid H, Phillips AJL, Selbmann L, Pfiegl WP, Horváth E, Bensch K, Kirk P, Kolaříková Z, Raja HA, Radek R, Papp V, Dima B, Ma J, Malosso E, Takamatsu S, Rambold G, Gannibal PB, Triebel D, Gautam AK, Avasthi S, Suetrong S, Timdal E, Fryar SC, Delgado G, Réblová M, Doilom M, Dolatabadi S, Pawłowska J, Humber RA, Kodsueb R, Sánchez-Castro I, Goto BT, Silva DKA, De Souza FA, Oehl F, Da Silva GA, Silva IR, Błaszkowski J, Jobim K, Maia LC, Barbosa FR, Fiua PO, Divakar PK, Shenoy BD, Castañeda-Ruiz RF, Somrithipol S, Karunaratna SC, Tibpromma S, Mortimer PE, Wanasinghe DN, Phookamsak R, Xu J, Wang Y, Fenghua T, Alvarado P, Li DW, Kušan I, Matočec N, Maharanachchikumbura SSN, Papizadeh M, Heredia G, Wartchow F, Bakhshi M, Boehm E, Youssef N, Hustad VP, Lawrey JD, Santiago ALCMA, Bezerra JDP, Souza-Motta CM, Firmino AL, Tian Q, Houbraken J, Hongsanan S, Tanaka K, Dissanayake AJ, Monteiro JS, Grossart HP, Suija A, Weerakoon G, Etayo J, Tsurykau A, Kuhnert E, Vázquez V, Mungai P, Damm U, Li QR, Zhang H, Boonmee S, Lu YZ, Becerra AG, Kendrick B, Brearley FQ, Motiejūnaitė J, Sharma B,

- Khare R, Gaikwad S, Wijesundara DSA, Tang LZ, He MQ, Flakus A, Rodriguez-Flakus P, Zhurbenko MP, McKenzie EHC, Stadler M, Bhat DJ, Liu JK, Raza M, Jeewon R, Nas-sanova ES, Prieto M, Jayalal RGU, Yurkov A, Schnittler M, Shchepin ON, Novozhilov YK, Liu P, Cavender JC, Kang Y, Mohammad S, Zhang LF, Xu RF, Li YM, Dayarathne MC, Ekanayaka AH, Wen TC, Deng CY, Lateef AA, Pereira OL, Navathe S, Hawksworth DL, Fan XL, Dissanayake LS, Erdođdu M (2020) Outline of Fungi and fungus-like taxa. *Mycosphere* 11: 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- Zhu H, Pan M, Wijayawardene NN, Jiang N, Ma R, Dai D, Tian C, Fan X (2021) The Hidden Diversity of Diatrypaceous Fungi in China. *frontiers in Microbiology* 12: 646262. <https://doi.org/10.3389/fmicb.2021.646262>