

Three new species of *Iodosphaeria* (Xylariomycetidae): *I. chiayiensis*, *I. jinghongensis* and *I. thailandica*

Lakmali S. Dissanayake¹, Diana S. Marasinghe^{2,6}, Milan C. Samarakoon²,
Sajeewa S.N. Maharachchikumbura³, Peter E. Mortimer⁴, Kevin D. Hyde^{2,5},
Chang-Hsin Kuo⁶, Ji-Chuan Kang¹

1 Engineering and Research Center for Southwest Bio-Pharmaceutical Resources of National Education Ministry of China, Guizhou University, Guiyang 550025, China **2** Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand **3** School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu 611731, China **4** Centre for Mountain Futures (CMF), Kunming Institute of Botany, Honghe County 654400, Yunnan, China **5** Innovative Institute for Plant Health, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, China **6** Department of Plant Medicine, National Chiayi University, 300 Syuefu Road, Chiayi City 60004, Taiwan

Corresponding author: Ji-Chuan Kang (jckang@gzu.edu.cn)

Academic editor: Andrew Miller | Received 27 September 2021 | Accepted 15 December 2021 | Published 7 January 2022

Citation: Dissanayake LS, Marasinghe DS, Samarakoon MC, Maharachchikumbura SSN, Mortimer PE, Hyde KD, Kuo C-H, Kang J-C (2022) Three new species of *Iodosphaeria* (Xylariomycetidae): *I. chiayiensis*, *I. jinghongensis* and *I. thailandica*. MycoKeys 86: 1–17. <https://doi.org/10.3897/mycokeys.86.75801>

Abstract

Three fungal specimens (two sexual and one asexual) were collected during fieldwork conducted in China, Taiwan and Thailand. Both sexual morphs share superficial, black ascocata surrounded by flexuous setae; 8-spored, unitunicate, cylindrical asci, with J+, apical ring, and ellipsoidal to allantoid, aseptate, guttulate ascospores. The asexual morph has ceratosporium-like conidia arising from aerial hyphae with a single arm and are usually attached or with 2–3 arms, brown, often with a subglobose to conical cell at the point of attachment. Morphological examinations and phylogenetic analyses of a combined LSU-ITS dataset *via* maximum likelihood and Bayesian analyses indicated that these three collections were new species. *Iodosphaeria chiayiensis* (sexual morph), *I. thailandica* (sexual morph) and *I. jinghongensis* (aseexual morph) are therefore introduced as new species in this study. *Iodosphaeria chiayiensis* has small, hyaline and ellipsoidal to allantoid ascospores, while *I. thailandica* has large ascocata, cylindrical to allantoid asci and hyaline to pale brown ascospores.

Keywords

Ceratosporium-like asexual morph, Sordariomycetes, taxonomy, three new taxa

Introduction

Iodosphaeria was introduced by Samuels et al. (1987) with its type *I. phyllophila* on a rachis of *Cyathea* sp., from Brazil. Only five of the nine *Iodosphaeria* species have been sequenced (Li et al. 2015; Marasinghe et al. 2019; Miller and Réblová 2021) and several species of them lack DNA-based sequence data. The sexual morph of *Iodosphaeria* is characterized by superficial, black, apapillate ascomata with unbranched, brown radial flexuous hairs, a two layered peridium composed of a pigmented outer layer and a hyaline inner layer; unitunicate, amyloid or non-amyloid, cylindrical to narrowly clavate, 8-spored ascii; and mostly allantoid to ellipsoidal, aseptate, hyaline ascospores with or without a gelatinous sheath (Miller and Réblová 2021). The asexual morphs of *Iodosphaeria* are considered selenosporella-like or ceratosporium-like (Samuels et al. 1987; Li et al. 2015; Miller and Réblová 2021). Members of *Iodosphaeria* are regarded as cosmopolitan species (Li et al. 2015). These species are usually saprobic on dead branches, twigs, stems, and petioles of economically important plants, such as *Alnus* sp., *Archontophoenix alexandrae*, *Arundinaria* sp., *Corylus* sp., *Cyathea dealbata*, *Podocarpus parlatorei*, *Polygonum chinense* and *Ripogonum scandens* (Samuels et al. 1987; Barr 1993; Hyde 1995; Candoussau et al. 1996; Hsieh et al. 1997; Taylor and Hyde 1999; Catania and Romero 2012; Li et al. 2015; Miller and Réblová 2021), but have never been reported as pathogens (Hyde et al. 2020a).

Samuels et al. (1987) accepted *Iodosphaeria* in Amphisphaeriaceae, and later, various authors placed it in Lasiosphaeriaceae and Trichosphaeriaceae (Barr 1990, 1994; Kang et al. 1998; Hilber and Hilber 2002; Jeewon et al. 2003). Again, Eriksson et al. (2001) placed *Iodosphaeria* in Amphisphaeriaceae. Later, Hilber and Hilber (2002) accommodated *Iodosphaeria* in the newly introduced family Iodosphaeriaceae. Maharachchikumbura et al. (2016) and Samarakoon et al. (2016) provided multigene phylogenies and accepted Iodosphaeriaceae in Xylariales. Hongsanan et al. (2017) treated it as Xylariomycetidae family *incertae sedis*, while Hyde et al. (2020a) and Wijayawardene et al. (2020) accepted Iodosphaeriaceae in Amphisphaeriales. In the most recent study of Miller and Réblová (2021), Iodosphaeriaceae is accounted as a family in Xylariales.

This study introduces three novel *Iodosphaeria* species from China, Taiwan, and Thailand. Detailed morphological descriptions, illustrations and a key are provided, and phylogenetic affinities of the new taxa are discussed.

Materials and methods

Morphological observations

Dead leaves were collected from Dahu Forest (Chiayi City, Taiwan) during autumn (September 2019), from dead twigs in Jinghong City (Yunnan Province, China) during winter (December 2019) and from dead leaves at MRC (Mushroom Research Centre, Chiang Mai, Thailand) during the rainy season (September 2020). Specimens

were treated following the methods outlined in Senanayake et al. (2020). A Motic SMZ 168 Series microscope was used to examine fruiting structures. Hand sections of the fruiting structures were mounted in water and 5% KOH for microscopic studies and microphotography. Indian ink was used to stain any gelatinous sheath around the ascospores and Melzer's reagent for ascal apical ring reaction. The micro-morphologies were examined using a Nikon ECLIPSE 80i compound microscope and photographed using a Canon 750D digital camera fitted to the microscope. Tarosoft (R) Image Frame Work program (IFW 0.97 version) and Adobe Photoshop CS6 software (Adobe Systems, USA) were used for image processing and measurements. The type specimens were deposited in the Mae Fah Luang University Herbarium (MFLU), Chiang Rai, Thailand and the Cryptogamic Herbarium, Kunming Institute of Botany Academia Sinica (HKAS), Chinese Academy of Sciences, Kunming, China. The new taxa were linked with Facesoffungi (Jayasiri et al. 2015) and Index Fungorum (<http://www.indexfungorum.org>).

DNA extraction, PCR amplification and sequencing

DNA extraction, PCR amplification and sequencing were carried out following the methods described in Dissanayake et al. (2020). Direct DNA extraction was done using a Biospin Fungus Genomic DNA Extraction Kit-BSC14S1 (BioFlux, P.R. China) with 15–20 fruiting bodies of the fungus as described in Wanasinghe et al. (2018). PCR amplification was done using LSU and ITS DNA regions with LR0R/LR5 (Vilgalys and Hester 1990) and ITS5/ITS4 (White et al. 1990) primer pairs, respectively. The thermal cycling program was followed by Wanasinghe et al. (2020). Purified PCR products were sent to a commercial sequencing provider, Beijing Biomed Gene Technology Co., Ltd., Shijingshan District, TsingKe Biological Technology Co., Beijing, China.

Phylogenetic analyses

Newly generated sequences were assembled and subjected to the standard BLAST search to identify the closest matches in GenBank. The accession numbers of taxa used in our analyses are shown in Table 1. Single datasets (LSU and ITS) were aligned using MAFFT v. 6.864b (<http://mafft.cbrc.jp/alignment/server/index.html>, Katoh and Standley 2013; Katoh et al. 2019), combined and manually improved using BioEdit v. 7.0.5.2 (Hall 1999). Maximum likelihood analysis and Bayesian inference (BI) were performed using RAxML-HPC2 on the XSEDE v. 8.2.10 tool and MrBayes 3.2.2 on the XSEDE tool in the CIPRES Science Gateway portal (Miller et al. 2012; Ronquist et al. 2012; Stamatakis 2014). The optimal ML tree was obtained with 1,000 separate runs under the GTR+GAMMA substitution model resulting from model tests using MrModeltest v. 2.3 (Nylander 2004) under the AIC (Akaike Information Criterion) implemented in PAUP v. 4.0b10. Maximum Likelihood bootstrap values (ML) equal or greater than 60% and Bayesian posterior probabilities (BYPP) equal or greater than 0.95 are presented above each node (Figure 1). All trees were visualized with FigTree v1.4.0 (Rambaut 2012), and

Table I. Taxa used in the phylogenetic analyses and corresponding GenBank accession numbers.

Taxon	Specimen/Strain	GenBank accession numbers ITS	GenBank accession numbers LSU	References
<i>Delonichicola siamense</i>	MFLUCC 15-0670 T	MF167586	MF158345	Perera et al. (2017)
<i>Furfurella luteostiolata</i>	CBS 143620 T	MK527842	MK527842	Voglmayr et al. (2019)
<i>Iodosphaeria chiayiensis</i>	MFLU 21-0042 T	MZ918994	MZ918992	This study
<i>I. foliicola</i>	NBM-F-07096 T	MZ509148	MZ509160	Miller and Réblová (2021)
<i>I. honghensis</i>	MFLU 19-0719 T	MK737501	MK722172	Marasinghe et al. (2019)
<i>I. jinghongensis</i>	HKAS 115761 T	MZ918989	MZ923776	This study
<i>I. phyllophilta</i>	PDD 56626	MZ509149	MZ509149	Miller and Réblová (2021)
<i>I. phyllophilta</i>	FC 5099-2d	MZ509150	N/A	Miller and Réblová (2021)
<i>I. phyllophilta</i>	ILL00121493 T	MZ509151	N/A	Miller and Réblová (2021)
<i>I. ripogoni</i>	PDD 103350	MZ509152	MZ509152	Miller and Réblová (2021)
<i>I. thailandica</i>	MFLU 21-0041 T	MZ923759	MZ923758	This study
<i>I. tongrenensis</i>	MFLU 15-0393 T	KR095282	KR095283	Li et al. (2015)
<i>Oxydothis metroxylonica</i>	MFLUCC 15-0281 T	KY206776	KY206765	Konta et al. (2016)
<i>O. palmicola</i>	MFLUCC 15-0806 T	KY206774	KY206763	Konta et al. (2016)
<i>O. phoenicis</i>	MFLUCC 18-0270 T	MK088066	MK088062	Hyde et al. (2020a)
<i>Pseudosporidesmium knawiae</i>	CBS 123529 T	FJ349609	FJ349610	Vu et al. (2019)
<i>P. lambertiae</i>	CBS 143169 T	MG386034	MG386087	Crous et al. (2017)
<i>Vialaea insculpta</i>	DAOM 240257	KC181926	KC181924	McTaggart et al. (2013)
<i>V. mangiferae</i>	MFLUCC 12-0808 T	KF724974	KF724975	Senanayake et al. (2014)
<i>V. minutella</i>	BRIP 56959	JX139726	JX139726	Shoemaker et al. (2013)

Types strains are indicated with (T). Newly generated sequences are indicated in bold. “N/A” sequences are unavailable.

Abbreviations: **BRIP**: Queensland Plant Pathology Herbarium, Australia; **CBS**: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; **DAOM**: Plant Research Institute, Department of Agriculture (Mycology), Ottawa, Canada; **HKAS**: Chinese Academy of Sciences, Kunming, China; **KUMCC**: Kunming Institute of Botany Culture Collection, Chinese Academy of Science, Kunming, China; **MFLUCC**: Mae Fah Luang University Culture Collection, Chiang Rai, Thailand; **MFLU**: Mae Fah Luang University Herbarium, Chiang Rai, Thailand; **Others**: information not available.

the final layout was done with Microsoft PowerPoint (2016). The finalized alignment and tree were registered in TreeBASE (submission ID TB2: S29095). Reviewer access URL: <http://purl.org/phylo/treebase/phylows/study/TB2:S29095?x-access-code=43fac9fe7622929c65c2bd4120a2c10a&format=html>

Results

Phylogenetic analyses

The combined LSU and ITS comprise 20 taxa including the outgroup taxa. The best scoring RAxML tree is shown in Figure with a final ML optimization likelihood value of -7278.703992. The matrix had 575 distinct alignment patterns, with 19.44% undetermined characters or gaps. Estimated base frequencies were: A = 0.245534,




Figure 1. RAxML tree based on a combined dataset of partial LSU and ITS sequence analyses. The tree is rooted to *Delonicicola siamense* (MFLUCC 15-0670) and *Furturella luteostiolata* (CBS 143620). Type strains are in bold, and the newly generated strains are in red.

$C = 0.244177$, $G = 0.286855$, $T = 0.223434$ substitution rates $AC = 1.190714$, $AG = 2.269637$, $AT = 1.889784$, $CG = 1.069908$, $CT = 5.997198$, $GT = 1.000000$; proportion of invariable sites $I = 0.39717$; gamma distribution shape parameters $\alpha = 0.578305$. Both trees (ML and BYPP) were similar in topology and did not differ in species relationships, which is in agreement with multi-gene phylogenies of previous studies (Marasinghe et al. 2019; Miller and Réblová 2021).

In the combined multi-gene phylogenetic analysis, Iodosphaeriaceae received 100% ML and 1.00 BYPP support values (Figure 1). Three strains of *Iodosphaeria phylophila* grouped as a monophyletic clade with 82% ML and 1.00 BYPP support. *Iodosphaeria honghensis* (MFLU 19-0719) nested as a sister clade to *I. phylophila* with 82% ML and 1.00 BYPP support. Within the *Iodosphaeria* clade, our new collections *viz.* HKAS 115761 (*I. jinghongensis*), MFLU 21-0042 (*I. chiayiensis*) and MFLU 21-0041 (*I. thailandica*) grouped as distinct lineages (Figure 1). *Iodosphaeria jinghongensis* was distinct from *I. ripogoni* by 100% ML and 1.00 BYPP support values. *Iodosphaeria chiayiensis* nested between *I. thailandica* and *I. jinghongensis*. However, this relationship is statistically not supported. *Iodosphaeria thailandica* received 100% ML and 1.00 BYPP support values. *Iodosphaeria foliicola* (NBM-F-07096) is grouped as the basal taxon in the Iodosphaeriaceae.

Taxonomy

Iodosphaeria chiayiensis Marasinghe, C.H. Kuo & K.D. Hyde, sp. nov.

IndexFungorum number: IF558412

Facesoffungi Number No: FoF09711

Figure 2

Etymology. The specific epithet *chiayiensis* refers to the city name where the fungus was collected.

Holotype. MFLU 21-0042.

Description. *Saprobic* on dead leaves of an unidentified host. **Sexual morph:** *Ascomata* 150–190 × 160–200 µm ($\bar{x} = 170 \times 180$ µm, n = 10), globose to subglobose, superficial, black, solitary to gregarious, consisting of numerous long, flexuous setae. *Setae* 3–5 µm wide, arising from cells at the peridium surface, brown, unbranched, septate, apex flattened. *Ostiole* periphysate, apapillate. *Peridium* 50–55 µm wide ($\bar{x} = 53.4$ µm, n = 10), comprises two layers of *textura angularis* cells, outer layer of dark brown to black thick-walled cells, and an inner layer of flattened, light brown. *Paraphyses* 2–4 µm wide, shorter than ascii, hyaline, embedded in a gelatinous matrix. *Asci* 60–90 × 8–10 µm ($\bar{x} = 72.9 \times 9.2$ µm, n = 30), 8-spored, unitunicate, cylindrical, shortly pedicellate, apex rounded, with a J+ apical ring. *Ascospores* 15–20 × 4–6 µm ($\bar{x} = 17.2 \times 5.2$ µm, n = 30), overlapping uni-seriate, ellipsoidal to allantoid, aseptate, hyaline, guttulate. **Asexual morph:** Undetermined

Material examined. Taiwan, Chiayi, Fanlu Township area, on dead leaves of an undetermined species, 10 September 2019, D.S Marasinghe, DTF018 (MFLU 21-0042, *holotype*).

Notes. *Iodosphaeria chiayiensis* resembles *I. polygoni* which has globose to sub globose, superficial, solitary to gregarious ascomata, cylindrical, short pedicellate ascii with J+, apical rings and ellipsoidal to allantoid, aseptate, guttulate ascospores. However, *I. chiayiensis* differs from *I. polygoni* in having smaller ascomata (150–190 × 160–200 µm vs. 270–475 × 250–500 µm) and shorter ascii (60–90 × 8–10 µm vs. 150–180 × 10–13 µm) (Hsieh et al. 1997). In the multi-gene phylogenetic analyses (Figure 1), our collection (*Iodosphaeria chiayiensis*, MFLU 21-0042) has close affinity to *I. thailandica*. However, it was not possible to compare *I. chiayiensis* and *I. jinghongensis* as they occur as different morphs.

Iodosphaeria jinghongensis L.S. Dissan., J.C. Kang & K.D. Hyde, sp. nov.

IndexFungorum number: IF558800

Facesoffungi Number No: FoF09712

Figure 3

Etymology. The specific epithet *jinghongensis* refers to the city name where the fungus was collected.

Holotype. HKAS 115761.




Figure 2. *Iodosphaeria chiayiensis* (MFLU 21-0042, holotype) **a** substrate **b** ascocarps on the host surface **c** section of ascoma **d** appearance of setae on peridium **e** peridium **f, g** asci **h** J+ apical ring (in Melzer's reagent) **i** paraphyses **j-m** ascospores (**m** stained in lactophenol cotton blue). Scale bars: 50 µm (**c, e**); 5 µm (**d**); 20 µm (**f-g**); 5 µm (**h, i**); 10 µm (**j-m**).

Description. Saprobic on dead twigs of an unidentified host. **Sexual morph:** Undetermined. **Asexual morph:** Colonies on natural substrate effuse, punctiform, scattered, blackish brown, mycelium mostly superficial, non-branched, hyaline,




Figure 3. *Iodosphaeria jinghongensis* (HKAS 115761, holotype) **a, b** colonies on the host surface **c–f** conidia, conidiogenous cells and conidiophores (black arrow shows hyphae, red arrow shows conidiophore). Scale bars: 20 μm (**c–f**).

smooth hyphae. *Conidiophores* micronematous, smooth, flexuous, pale brown. *Conidia* ceratosporium-like, arising from aerial hyphae, solitary, dry, composed of a central cell and 2–3 arms. *Arms* 70–93 \times 9–14 μm ($\bar{x} = 79.8 \times 12.1 \mu\text{m}$, $n = 20$), wide at the tip 5–8 μm ($\bar{x} = 6.9 \mu\text{m}$), radiating from the centrally located attachment




Figure 4. Asexual morph of *Iodosphaeria ripogoni* (ceratosporium-like conidia). Redrawn from: Samuels et al. (1987). Scale bar: 20 μm .

point, multi-septate (9–10), each septum with a central pore, brown, often with a sub-globose to conical cell at the point of attachment, dehiscence scar circular 3–4 μm diam. ($\bar{x} = 3.5 \mu\text{m}$).

Material examined. China, Yunnan Province, Xishuangbanna Dai Autonomous Prefecture, Jinghong City, Jinghaxiang ($21^{\circ}78'06.17''\text{N}$, $101^{\circ}05'61.22''\text{E}$), on a dead twig of undetermined species, 19 December 2019, D.N. Wanasinghe, DW060 (HKAS 115761, *holotype*).

Notes. *Iodosphaeria jinghongensis* is similar to *I. ripogoni* in having septate, brown, subglobose to conical conidia with 2–3 arms (Figure 4; Samuels et al. 1987). However, *I. jinghongensis* differs from *I. ripogoni* in having smaller arms ($70\text{--}93 \times 9\text{--}14 \mu\text{m}$ vs $95\text{--}120 \times 14\text{--}16 \mu\text{m}$). *Iodosphaeria ripogoni* was collected from the stem of *Ripogonum scandens* from New Zealand, and *I. jinghongensis* was collected from twigs of undetermined species from China.

***Iodosphaeria thailandica* L.S. Dissan., Marasinghe, & K.D. Hyde, sp. nov.**

IndexFungorum number: IF558411

Facesoffungi number No: FoF09710

Figure 5

Etymology. The specific epithet *thailandica* refers to the country where the fungus was collected.

Holotype. MFLU 21-0041

Description. *Saprobic* on dead leaves of unidentified host. **Sexual morph:** *Ascomata* 250–285 × 250–295 µm ($\bar{x} = 267.3 \times 272$ µm, n = 10), globose to subglobose, superficial, black, solitary to gregarious, consisting of numerous long, flexuous setae. *Setae* 4.5 µm wide, arising from cells at the peridium surface, dark brown to brown, unbranched, septate. *Ostiole* periphysate, apapillate. *Peridium* 40–50 µm wide ($\bar{x} = 44.6$ µm, n = 10), comprising two layers of cells of *textura angularis*, outer layer of dark brown to black thick-walled cells and an inner layer of flattened, hyaline cells. *Paraphyses* 5–8 µm wide, length as longer than ascii, septate, hyaline, branched, embedded in a gelatinous matrix. *Asci* 65–100 × 8–10 µm ($\bar{x} = 84.3 \times 8.9$ µm, n = 30), 8-spored, unitunicate, cylindrical, short pedicellate, apex rounded, with a J+ apical ring. *Ascospores* 20–35 × 2–4 µm ($\bar{x} = 29.1 \times 3.2$ µm, n = 30), overlapping uni-seriate, cylindrical to allantoid, aseptate, hyaline to pale brown, guttulate, slightly curved. **Asexual morph:** Undetermined.

Material examined. Thailand, Chiang Mai, Mushroom Research Centre, on dead leaves of an undetermined species, 11 September 2020, D.S Marasinghe, DMRC011 (MFLU 21-0041, *holotype*)

Notes. *Iodosphaeria thailandica* shares similar characteristics with *I. honghensis* in having globose to subglobose, superficial, solitary to gregarious ascomata, cylindrical, short pedicellate, J+, apical ring and cylindrical to allantoid ascospores with aseptate, guttulate ascospores (Marasinghe et al. 2019). However, *I. thailandica* differs from *I. honghensis* in having long, narrow (20–35 × 2–4 µm) and hyaline to pale brown ascospores versus short, broad (18.5–22.5 × 4.5–6.5 µm) and hyaline ascospores. In the phylogenetic analyses, *I. thailandica* is distinct from other species in the genus by 100 % ML and 1.00BYPP and sister to the *I. chiayiensis*. *Iodosphaeria thailandica* has larger ascomata (250–285 × 250–295 µm), cylindrical to allantoid ascospores, while the ascomata of *I. chiayiensis* are smaller (150–190 × 160–200 µm) and ascospores are hyaline and ellipsoidal to allantoid. *Iodosphaeria thailandica* is the first report of *Iodosphaeria* from Thailand.

Key to the accepted *Iodosphaeria* species based on known sexual morph

- | | | |
|---|---|------------------------|
| 1 | Asci with a distinct apical ring | 2 |
| - | Asci lacking a distinct apical ring | 10 |
| 2 | Apical ring not staining blue in Melzer's reagent | <i>I. arundinariae</i> |
| - | Apical ring staining blue in Melzer's reagent..... | 3 |

3	Ascomata immersed to erumpent.....	<i>I. aquatica</i>
—	Ascomata superficial	4
4	Ascospores guttulate	5
—	Ascospores eguttulate.....	8
5	Ascospores ellipsoidal.....	6
—	Ascospores cylindrical	7
6	Ascomata 270–475 × 250–500 µm.....	<i>I. polygoni</i>
—	Ascomata 150–190 × 160–200 µm.....	<i>I. chiayiensis</i>
7	Ascospores 18.5–22.5 × 4.5–6.5 µm, hyaline.....	<i>I. honghensis</i>
—	Ascospores 20–35 × 2–4 µm, hyaline to pale brown.....	<i>I. thailandica</i>
8	Asci shorter than 150 µm.....	9
—	Asci longer than 150 µm.....	<i>I. tongrenensis</i>
9	Ascospores allantoid.....	11
—	Ascospores ellipsoidal.....	<i>I. podocarpi</i>
10	Ascospores with a mucilaginous sheath	<i>I. ripogoni</i>
—	Ascospores without a mucilaginous sheath	<i>I. hongkongensis</i>
11	Paraphyses of similar length to asci	<i>I. foliicola</i>
—	Paraphyses longer than asci	<i>I. phyllophila</i>

Discussion

Iodosphaeria is seldom collected. In 15 years of studying fungi in Hong Kong, only a single collection was found despite intensive collection efforts (Taylor and Hyde 1999). *Iodosphaeria* is widely distributed in temperate and tropical regions, e.g., China (Guizhou, Yunnan), Europe (Belgium, Germany), Great Britain, Canada, Hong Kong, New Zealand, South America (Brazil, Argentina, French Guiana), Taiwan and USA (Louisiana) (Samuels et al. 1987; Barr 1993; Hyde 1995; Candoussau et al. 1996; Hsieh et al. 1997; Taylor and Hyde 1999; Catania and Romero 2012; Li et al. 2015; Marasinghe et al. 2019; Miller and Réblová 2021). This genus is saprobic on dead plant substrates in terrestrial grassland habitats (Barr 1993), on fern rachides (Samuels et al. 1987), on dead petioles of palms (Taylor and Hyde 1999), and on submerged wood in freshwater (Hyde 1995) but has never been reported as pathogenic on hosts. They are likely endophytes that become saprobes during leaf senescence (Hyde et al. 2020a). *Iodosphaeria* species may not be host-specific due to their wide distribution range (Miller and Réblová 2021). The genus may be much more diverse than presently known, as is true for many other microfungal genera (Hyde et al. 2020b).

The asexual morphs of this genus were recorded as selenosporella- or ceratosporium-like (Samuels et al. 1987; Li et al. 2015; Marasinghe et al. 2019). *Iodosphaeria phyllophila*, *I. polygoni* and *I. ripogoni* (Figure 4) were introduced with both sexual and asexual morphs (Hsieh et al. 1997; Samuels et al. 1987). *Iodosphaeria honghensis* and *I. tongrenensis* were observed to have ceratosporium-like conidia on their host surface




Figure 5. *Iodosphaeria thailandica* (MFLU 21-0041, holotype) **a** substrate **b, c** ascomata on the host surface **d** peridium **e** section of ascocarpi **f** appearance of setae (black arrow) on peridium **g** setae **h** paraphyses **i, j** ascii **k** J+ apical ring (in Melzer's reagent) **l–q** ascospores (**p, q**, stained in Lactophenol Cotton Blue). Scale bars: 10 µm (**d**); 100 µm (**e**); 5 µm (**f–h**); 20 µm (**i, j**); 10 µm (**k–q**).

(Li et al. 2015; Marasinghe et al. 2019). Samuels et al. (1987) observed another asexual morph of selenosporella-like conidia that was different from ceratosporium-like conidia. In present study, we establish ceratosporium-like conidia as an asexual morph of *Iodosphaeria*.

Acknowledgements

Ji-Chuan Kang thanks the National Natural Science Foundation of China (NSFC Grants Nos. 31670027 & 31460011) and the Open Fund Program of Engineering Research Center of Southwest Bio-Pharmaceutical Resources, Ministry of Education, Guizhou University (No. GZUKEY 20160705). Shaun Pennycook is thanked for the nomenclatural advice. Dhanushka Wanasinghe is thanked for the specimen collections and for providing sequence data. Peter Mortimer thanks the National Science Foundation of China (grant no. 41761144055) and High-End Foreign Experts” in the High-Level Talent Recruitment Plan of Yunnan Province (2021). Austin G. Smith at World Agroforestry (ICRAF), Kunming Institute of Botany, China, is thanked for English editing.

References

- Barr ME (1990) Prodromus to non-lichenized pyrenomycetous members of the class Hymenomycetes. *Mycotaxon* 39: 43–184.
- Barr ME (1993) Redisposition of some taxa described by J. B. Ellis. *Mycotaxon* 46: 45–76.
- Barr ME (1994) Notes on the Amphisphaeriaceae and related families. *Mycotaxon* 51: 191–224.
- Candoussau F, Magni JF, Petrini LE, Barr ME, Petrini O (1996) Bambusicolous fungi collected in Southwestern France: an annotated list. *Mycologia Helvetica* 8: 11–20.
- Catania MV, Romero AI (2012) *Crassoascus monocaudatus* and *Iodosphaeria podocarpi*, two new species on *Podocarpus parlatorei* from “Las Yungas”, Argentina. *Mycosphere* 3: 37–44. <https://doi.org/10.5943/mycosphere/3/1/4>
- Crous PW, Wingfield MJ, Burgess TI, Carnegie AJ, Hardy GEStJ, Smith D, Summerell BA, Cano-Lira JF, Guarro J, Houbraken J, Lombard L, Martín MP, Sandoval-Denis M, Alexandrova AV, Barnes CW, Baseia IG, Bezerra JDP, Guarnaccia V, May TW, Hernández-Restrepo M, Stchigel AM, Miller AN, Ordoñez ME, Abreu VP, Accioly T, Agnello C, Agustín Colmán A, Albuquerque CC, Alfredo DS, Alvarado P, Araújo-Magalhães GR, Arauzo S, Atkinson T, Barili A, Barreto RW, Bezerra JL, Cabral TS, Camello Rodríguez F, Cruz RHSF, Daniëls PP, da Silva BDB, de Almeida DAC, de Carvalho Júnior AA, Decock CA, Delgat L, Denman S, Dimitrov RA, Edwards J, Fedosova AG, Ferreira RJ, Firmino AL, Flores JA, García D, Gené J, Góis JS, Gomes AAM, Gonçalves CM, Gouliamova DE, Groenewald M, Guéorguiev BV, Guevara-Suarez M, Gusmão LFP, Hosaka K, Hubka V, Huhndorf SM, Jadan M, Jurjević Ž, Kraak B, Kučera V, Kumar TKA, Kušan I, Lacerda SR, Lamertthon S, Lisboa WS, Loizides M, Luangsa-ard JJ, Lysková P, Mac

- Cormack WP, Macedo DM, Machado AR, Malysheva EF, Marinho P, Matočec N, Meijer M, Mešić A, Mongkolsamrit S, Moreira KA, Morozova OV, Nair KU, Nakamura N, Noisripoom W, Olariaga I, Oliveira RJV, Paiva LM, Pawar P, Pereira OL, Peterson SW, Prieto M, Rodríguez-Andrade E, Rojo De Blas C, Roy M, Santos ES, Sharma R, Silva GA, Souza-Motta CM, Takeuchi-Kaneko Y, Tanaka C, Thakur A, Smith MT, Tkalčec Z, Valenzuela-Lopez N, Van der Kleij P, Verbeken A, Viana MG, Wang XW, Groenewald JZ (2017) Fungal Planet description sheets: 625–715. Persoonia 39: 270–467. <https://doi.org/10.3767/persoonia.2017.39.11>
- Dissanayake AJ, Bhunjun CS, Maharachchikumbura SSN, Liu JK (2020) Applied aspects of methods to infer phylogenetic relationships amongst fungi. Mycosphere 11: 2652–2676. <https://doi.org/10.5943/mycosphere/11/1/18>
- Eriksson OE, Baral HO, Currah RS, Hansen K, Kurtzman CP, Rambold G, Laessøe T (2001) Outline of Ascomycota-2001. Myconet 7: 1–88. <https://doi.org/10.1038/npg.els.0000346>
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium, Series 41: 95–98.
- Hilber R, Hilber O (2002) The genus *Lasiosphaeria* and allied taxa. Publ 9. Germany, 1–18.
- Hongsanan S, Maharachchikumbura SSN, Hyde KD, Samarakoon MC, Jeewon R, Zhao Q, Al-Sadi AM, Bahkali AH (2017) An updated phylogeny of Sordariomycetes based on phylogenetic and molecular clock evidence. Fungal Diversity 84: 25–41. <https://doi.org/10.1007/s13225-017-0384-2>
- Hsieh WH, Chen CY, Sivanesan A (1997) *Iodosphaeria polygoni* sp. nov. a new pyrenomycte from Taiwan. Mycological Research 101: 841–842. <https://doi.org/10.1017/S095375629600319X>
- Hyde KD (1995) Tropical Australian freshwater fungi. VII. New genera and species of ascomycetes. Nova Hedwigia 61: 119–140.
- Hyde KD, Norphanphoun C, Maharachchikumbura SSN, Bhat DJ, Jones EBG, Bundhun D, Chen YJ, Bao DF, Boonmee S, Calabon MS, Chaiwan N, Chethana KWT, Dai DQ, Da-yarathne MC, Devadatha B, Dissanayake AJ, Dissanayake LS, Doilom M, Dong W, Fan XL, Goonasekara ID, Hongsanan S, Huang SK, Jayawardena RS, Jeewon R, Karunaratna A, Konta S, Kumar V, Lin CG, Liu JK, Liu NG, Luangsa-ard J, Lumyong S, Luo ZL, Marasinghe DS, McKenzie EHC, Niego AGT, Niranjan M, Perera RH, Phukhamsakda C, Rathnayaka AR, Samarakoon MC, Samarakoon SMBC, Sarma VV, Senanayake IC, Shang QJ, Stadler M, Tibpromma S, Wanasinghe DN, Wei DP, Wijayawardene NN, Xiao YP, Yang J, Zeng XY, Zhang SN, Xiang MM (2020a) Refined families of Sordariomycetes. Mycosphere 11: 305–1059. <https://doi.org/10.5943/mycosphere/11/1/7>
- Hyde KD, Jeewon R, Chen YJ, Bhunjun CS, Calabon MS, Jiang HB, Lin CG, Norphanphoun C, Sysouphanthong P, Pem D, Tibpromma S, Zhang Q, Doilom M, Jayawardena RS, Liu JK, Maharachchikumbura SSN, Phukhamsakda C, Phookamsak R, Al-Sadi AM, Naritsada Thongklang N, Wang Y, Gafforov Y, Jones EBG, Lumyong S (2020b) The numbers of fungi: is the descriptive curve flattening?. Fungal Diversity 103: 219–271. <https://doi.org/10.1007/s13225-020-00458-2>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunaratna SC, Liu JK, Luangsa-ard

- JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang K, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, De Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74: 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jeewon R, Liew ECY, Hyde KD (2003) Molecular systematics of the Amphisphaeriaceae based on cladistic analyses of partial LSU rDNA gene sequences. *Mycological Research* 107: 1392–1402. <https://doi.org/10.1017/S095375620300875X>
- Kang JC, Kong RYC, Hyde KD (1998) Studies on the Amphisphaerales 1. Amphisphaeriaceae (*sensu stricto*) and its phylogenetic relationships inferred from 5.8S rDNA and ITS2 sequences. *Fungal Diversity* 1: 147–157.
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20: 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Konta S, Hongsanan S, Tibpromma S, Thongbai B, Maharachchikumbura SSN, Bahkali AH, Hyde KD, Boonmee S (2016) An advance in the endophyte story: Oxydothidaceae fam. nov. with six new species of *Oxydothis*. *Mycosphere* 9: 1425–1446. <https://doi.org/10.5943/mycosphere/7/9/15>
- Li Q, Kang JC, Hyde KD (2015) A multiple gene genealogy reveals the phylogenetic placement of *Iodosphaeria tongrenensis* sp. nov. in Iodosphaeriaceae (Xylariales). *Phytotaxa* 234: 121–132. <http://dx.doi.org/10.11646/phytotaxa.234.2.2>
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EHC, Bhat JD, Dayarathne MC, Huang SK, Norphanphoun C, Senanayake IC, Perera RH, Shang QJ, Xiao Y, D'souza MJ, Hongsanan S, Jayawardena RS, Daranagama DA, Konta S, Goonasekara ID, Zhuang WY, Jeewon R, Phillips AJL, Abdel-Wahab MA, Al-Sadi AM, Bahkali AH, Boonmee S, Boonyuen N, Cheewangkoon R, Dissanayake AJ, Kang J, Li QR, Liu JK, Liu XZ, Liu ZY, Luangsa-ard JJ, Pang KL, Phookamsak R, Promputtha I, Suetrong S, Stadler M, Wen T, Wijayawardene NN (2016) Families of Sordariomycetes. *Fungal Diversity* 79: 1–317. <https://doi.org/10.1007/s13225-016-0369-6>
- McTaggart AR, Grice KR, Shivas RG (2013) First report of *Vialaea minutella* in Australia, its association with mango branch dieback and systematic placement of *Vialaea* in the Xylariales. *Australasian Plant Disease Notes* 8: 63–66. <https://doi.org/10.1007/s13314-013-0096-8>
- Miller AN, Réblová M (2021) Phylogenetic placement of Iodosphaeriaceae (Xylariales, Ascomycota), designation of an epitype for the type species of *Iodosphaeria*, *I. phyllophila*, and description of *I. foliicola* sp. nov. *Fungal Systematics and Evolution* 8: 49–64. <https://doi.org/10.3114/fuse.2021.08.05>

- Miller MA, Pfeiffer W, Schwartz T (2012) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Gateway Computing Environments Workshop 2010 (GCE). Ieee, 1–8. <https://doi.org/10.1109/GCE.2010.5676129> [Accessed on 20 September 2021]
- Nylander JA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden, 1–3.
- Perera RH, Maharachchikumbura SSN, Jones EBG, Bahkali AH, Elgorban AM, Liu JK, Liu ZY, Hyde KD (2017) *Delonicicola siamense* gen. & sp. nov. (*Delonicolaceae* fam. nov., *Delonicolales* ord. nov.), a saprobic species from *Delonix regia* seed pods. Cryptogamie, Mycologie 38: 321–340. <https://doi.org/10.7872/crym/v38.iss3.2017.321>
- Rambaut A (2012) FigTree v. 1.4. Molecular evolution, phylogenetics and epidemiology. Edinburgh, UK: University of Edinburgh, Institute of Evolutionary Biology. <http://tree.bio.ed.ac.uk/software/figtree/> [Accessed on 20 September 2021]
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Samarakoon MC, Hyde KD, Promputtha I, Hongsanan S, Ariyawansa HA, Maharachchikumbura SSN, Daranagama DA, Stadler M, Mapook A (2016) Evolution of Xylariomycetidae (Ascomycota: Sordariomycetes). Mycosphere 7: 1746–1761. <https://doi.org/10.5943/mycosphere/7/11/9>
- Samuels GJ, Müller E, Petrini O (1987) Studies in the Amphisphaeriaceae (*sensu lato*) 3. New species of *Monographella* and *Pestalosphaeria* and two new genera. Mycotaxon 28: 473–499.
- Senanayake IC, Maharachchikumbura SSN, Mortimer PE, Xu J, Bhat JD, Hyde KD (2014) Phylogenetic studies on Vialaeaceae reveals a novel species, *Vialaea mangiferae*. Sydowia 66: 203–216. [https://doi.org/10.12905/0380.sydowia66\(2\)2014-0203](https://doi.org/10.12905/0380.sydowia66(2)2014-0203)
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Wanasinghe DN, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Chomnunti P, Boonmee S, Jayawardena RS, Wijayawardene NN, Doilom M, Jeewon R, Bhat JD, Zhang HX, Xie N (2020) Morphological approaches in studying fungi: collection, examination, isolation, sporulation and preservation. Mycosphere 11: 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Shoemaker RA, Hambleton S, Liu M (2013) *Vialaea insculpta* revisited. North American Fungi 8: 1–13. <https://doi.org/10.2509/naf2013.008.010>
- Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Taylor JE, Hyde KD (1999) Fungi from palms. XL. *Iodosphaeria*. Sydowia 51: 127–132.
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172: 4238–4246. <https://doi.org/10.1128/JB.172.8.4238-4246.1990>
- Vu D, Groenewald M, de Vries M, Gehrmann T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes

- boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92: 135–154. <https://doi.org/10.1016/j.simyco.2018.05.001>
- Wanasinghe DN, Phukhamsakda C, Hyde KD, Jeewon R, Lee HB, Jones EBG, Tibpromma S, Tennakoon DS, Dissanayake AJ, Jayasiri SC, Gafforov Y, Camporesi E, Bulgakov TS, Ekanayake AH, Perera RH, Samarakoon MC, Goonasekara ID, Mapook A, Li WJ, Senanayake IC, Li JF, Norphanphoun C, Doilom M, Bahkali AH, Xu JC, Mortimer PE, Tibell L, Savic ST, Karunarathna SC (2018) Fungal Diversity notes 709–839: taxonomic and phylogenetic contributions to fungal taxa with an emphasis on fungi on Rosaceae. *Fungal Diversity* 89: 1–236. <https://doi.org/10.1007/s13225-018-0395-7>
- Wanasinghe DN, Wijayawardene NN, Xu J, Cheewangkoon R, Mortimer PE (2020) Taxonomic novelties in *Magnolia*-associated pleosporalean fungi in the Kunming Botanical Gardens (Yunnan, China). *PLoS ONE* 15(7): e0235855. <https://doi.org/10.1371/journal.pone.0235855>
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, Lumbsch HT, Kukwa M, Issi IV, Madrid H, Phillips AJL, Selbmann L, Pfleigler WP, Horváth E, Bensch K, Kirk P, Kolaříková Z, Raja HA, Radek R, Papp V, Dima B, Ma J, Malosso E, Takamatsu S, Rambold G, Gannibal PB, Triebel D, Gautam AK, Avasthi S, Suetrong S, Timdal E, Fryar SC, Delgado G, Réblová M, Doilom M, Dolatabadi S, Pawłowska J, Humber RA, Kodsuē R, Sánchez-Castro I, Goto BT, Silva DKA, De Souza FA, Oehl F, Da Silva GA, Silva IR, Błaszkowski J, Jobim K, Maia LC, Barbosa FR, Fiúza PO, Divakar PK, Shenoy BD, Castañeda-Ruiz RF, Somrithipol S, Karunarathna SC, Tibpromma S, Mortimer PE, Wanasinghe DN, Phookamsak R, Xu J, Wang Y, Fenghua T, Alvarado P, Li DW, Kušan I, Matočec N, Maharachchikumbura SSN, Papizadeh M, Heredia G, Wartchow F, Bakhshi M, Boehm E, Youssef N, Hustad VP, Lawrey JD, Santiago ALCMA, Bezerra JDP, Souza-Motta CM, Firmino AL, Tian Q, Houbraken J, Hongsanan S, Tanaka K, Dissanayake AJ, Monteiro JS, Grossart HP, Suija A, Weerakoon G, Etayo J, Tsurykau A, Kuhnert E, Vázquez V, Mungai P, Damm U, Li QR, Zhang H, Boonmee S, Lu YZ, Becerra AG, Kendrick B, Brearley FQ, Motiejūnaitė J, Sharma B, Khare R, Gaikwad S, Wijesundara DSA, Tang LZ, He MQ, Flakus A, Rodriguez-Flakus P, Zhurbenko MP, McKenzie EHC, Stadler M, Bhat DJ, Liu JK, Raza M, Jeewon R, Nasanova ES, Prieto M, Jayalal RGU, Yurkov A, Schnittler M, Shchepin ON, Novozhilov YK, Liu P, Cavender JC, Kang Y, Mohammad S, Zhang LF, Xu RF, Li YM, Dayarathne MC, Ekanayaka AH, Wen TC, Deng CY, Lateef AA, Pereira OL, Navathe S, Hawksworth DL, Fan XL, Dissanayake LS, Erdöđu M (2020) Outline of Fungi and fungus-like taxa. *Mycosphere* 11: 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- White TJ, Bruns T, Lee S, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: a guide to methods and applications. Academic Press, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>