

# Two novel freshwater hyphomycetes, in *Acrogenospora* (Minutisphaerales, Dothideomycetes) and *Conioscypha* (Conioscyphales, Sordariomycetes) from Southwestern China

Lu Li<sup>1,2,3\*</sup>, Hong-Zhi Du<sup>1,2,4\*</sup>, Vinodhini Thiyagaraja<sup>3</sup>, Darbhe Jayarama Bhat<sup>5,6</sup>,  
Rungtiwa Phookamsak<sup>3,7</sup>, Ratchadawan Cheewangkoon<sup>1,2</sup>

1 Department of Entomology and Plant Pathology, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand

2 Innovative Agriculture Research Centre, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand

3 Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan Province, China

4 School of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang 550025, Guizhou Province, China

5 Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

6 Vishnugupta Vishwavidyapeetam, Ashoke, Gokarna 581326, India

7 Honghe Center for Mountain Futures, Kunming Institute of Botany, Chinese Academy of Sciences, Honghe County 654400, Yunnan, China

Corresponding author: Ratchadawan Cheewangkoon (ratchadawan.c@cmu.ac.th)



Academic editor: Ning Jiang

Received: 4 November 2023

Accepted: 21 December 2023

Published: 31 January 2024

Citation: Li L, Du H-Z, Thiyagaraja V, Bhat DJ, Phookamsak R, Cheewangkoon R (2024) Two novel freshwater hyphomycetes, in *Acrogenospora* (Minutisphaerales, Dothideomycetes) and *Conioscypha* (Conioscyphales, Sordariomycetes) from Southwestern China. MycoKeys 101: 249–273. <https://doi.org/10.3897/mycokeys.101.115209>

Copyright: © Lu Li et al.

This is an open access article distributed under the terms of the CC0 Public Domain Dedication.

## Abstract

Freshwater fungi are highly diverse in China and frequently reported from submerged wood, freshwater insects, herbaceous substrates, sediments, leaves, foams, and living plants. In this study, we investigated two freshwater species that were collected from Yunnan and Guizhou provinces in China. Detailed morphological analysis complemented by multi-gene phylogenetic analyses based on LSU, SSU, ITS, *RPB2* and *TEF1-a* sequences data revealed them to be two new saprobic species, namely *Acrogenospora alangii* **sp. nov.** and *Conioscypha yunnanensis* **sp. nov.** in their asexual morphs. Additionally, *Acrogenospora alangii* **sp. nov.** is reported for the first time as a freshwater ascomycete associated with the medicinal plant *Alangium chinense* (Alangiaceae). Detailed morphological descriptions, illustrations and updated phylogenetic relationships of the new taxa are provided herein.

**Key words:** Acrogenosporaceae, Conioscyphaceae, freshwater fungi, new taxa, taxonomy

## Introduction

The freshwater fungi in China are taxonomically highly diverse which include members of Dothideomycetes, Eurotiomycetes, Laboulbeniomycetes, Leotiomycetes, Orbiliomycete, Pezizomycetes and Sordariomycetes (Hu et al. 2013; Calabon et al. 2022). The freshwater fungi are ecologically diverse, occurring on various substrates, including submerged wood, freshwater foams, herbaceous substrates, insects, leaves, sediments and other organic matter, and living plants (Hu et al. 2013; Shen et al. 2022; Calabon et al. 2023). Most species are well-known as saprobes and they play an important role in ecological func-

\* These authors contributed equally to this work.

tioning as decomposers but also can be pathogens as well as symbionts on humans and plants (Su et al. 2015; Su et al. 2016; Dong et al. 2020).

The order Minutisphaerales (Dothideomycetes) is known as the order for freshwater fungi and comprises two families, viz. Acrogenosporaceae and Minutisphaeraceae (Wijayawardene et al. 2022). Acrogenosporaceae was introduced by Jayasiri et al. (2018) to accommodate *Acrogenospora* based on morpho-molecular evidence. The genus *Acrogenospora* was introduced by Ellis (1971) for two species namely *A. sphaerocephala* (the type species), and *A. carmichaeliana* (as *Farlowiella carmichaeliana*; asexual morph). A year later, Ellis (1972) added another new species, *A. setiformis*. While Goh et al. (1998) revised the genus and accepted eight species, Bao et al. (2020) re-investigated *Acrogenospora* and added seven new species that were reported from freshwater habitat. Subsequently, two new species *A. guizhouensis* and *A. stellata* were introduced in asexual and sexual states, respectively (Tan et al. 2022; Hyde et al. 2023). Presently, there are 23 epithets for *Acrogenospora* in Index Fungorum (<http://indexfungorum.org/Names/Names.asp>; accessed on 20 Nov. 2023).

*Acrogenospora* was considered as the asexual morph of *Farlowiella* which was further supported by the morpho-molecular analyses conducted by Jayasiri et al. (2018) and the pleomorphic status of these two genera was confirmed by Rossman et al. (2015) who recommended protecting the name *Acrogenospora* over *Farlowiella* based on the wider use and fewer name changes. The sexual morph of this genus is characterized by hysterothecial, thick-walled, apparently solitary to gregarious, but remaining erect and elevated and presenting an almost stipitate ascomata with a prominent sunken slit, 8-spored, cylindrical-clavate, short pedicellate asci and 1–2-celled, hyaline or moderately pigmented ascospores (Sivanesan 1984; Boehm et al. 2009). The asexual morph is characterized by macronematous, mononematous, simple, brown, sometimes percurrently proliferating conidiophores; monoblastic, terminal or intercalary conidiogenous cells with globose, ellipsoid or obovoid, olivaceous to dark brown conidia (Hughes et al. 1978; Goh et al. 1998). The members of *Acrogenospora* mostly show similar morphology, but mainly distinguished by degree of pigmentation of the conidiophores, and conidial shape, size, color, guttules and basal cells (Hughes et al. 1978; Bao et al. 2020).

Conioscyphales (Sordariomycetes), a largely freshwater order, was introduced by Réblová et al. (2016) to accommodate a single family Conioscyphaeaceae and a genus *Conioscypha*. The order was placed within Hypocreomycetidae (Réblová et al. 2016). However, Conioscyphales clustered within the newly introduced subclass Savoryellomycetidae in the phylogenetic analyses conducted by Hongsanan et al. (2017). Höhnelt (1904) had introduced *Conioscypha* with *C. lignicola* as the type species and the genus currently accommodates 18 species (Höhnelt 1904; Matsushima 1975, 1993, 1996; Shearer 1973; Shearer and Motta 1973; Udagawa and Toyazaki 1983; Kirk 1984; Chen and Tzean 2000; Réblová and Seifert 2004; Crous et al. 2014, 2018; Zelski et al. 2015; Chuaseeharonnachai et al. 2017; Hernández et al. 2017; Feng and Yang 2018; Turland et al. 2018; Liu et al. 2019; Luo et al. 2019; Hyde et al. 2020; Jiang et al. 2022). Réblová and Seifert (2004) established *Conioscyphascus* based on *C. varius* which is the sexual morph of *Conioscypha varia* and the sexual-asexual linkage was further confirmed by culture studies and molecular data (Réblová and

Seifert 2004; Zelski et al. 2015). According to the nomenclatural priority, *Conioscyphascus* is synonymized under *Conioscypha* (Turland et al. 2018).

Species of *Conioscypha* are mostly reported from freshwater and terrestrial habitats and primarily recorded in their asexual morph. Only few species are reported in sexual morph (Shearer 1973; Shearer and Motta 1973; Kirk 1984; Zelski et al. 2015). The asexual morph is characterized by the enteroblastic percurrent conidiogenesis in distinct conidiogenous cells that retain successive wall layers at the same level as multi-collaretted as each conidium ruptures through the apex with dematiaceous aseptate conidia of various shapes (Shearer 1973; Shearer and Motta 1973; Kirk 1984; Zelski et al. 2015). The sexual morph is characterized by perithecial ascomata that are immersed to superficial, globose to subglobose, cylindrical-clavate asci with a pronounced non-amyloid apical annulus, transversely multi-septate and hyaline ascospores (Luo et al. 2019).

Guizhou and Yunnan provinces are mostly referred as part of the Southwestern China (Feng and Yang 2018; Jiang et al. 2022). This region is a center of biodiversity for freshwater fungi (Shen et al. 2022). Many new freshwater fungi have been reported in Yunnan and Guizhou provinces in recent years (Su et al. 2016; Wang et al. 2016; Li et al. 2017, 2020; Luo et al. 2018a, b, 2019; Zhao et al. 2018; Dong et al. 2020; Wan et al. 2021; Shen et al. 2022). In particular, Yunnan province stands out as a hotspot for freshwater fungal research (Luo et al. 2019; Dong et al. 2020; Shen et al. 2022). The diversity of freshwater fungi in streams and rivers in northwestern Yunnan has been intensely studied, resulting in the discovery of a large number of new species and new records in some highly diverse genera e.g. *Acrogenospora*, *Dictyosporium*, *Distoseptispora*, *Pleurotheciella*, *Sporidesmium* and *Sporoschisma* (Su et al. 2016; Wang et al. 2016; Li et al. 2017, 2020; Luo et al. 2018a, b, 2019; Zhao et al. 2018; Bao et al. 2020; Wan et al. 2021; Shen et al. 2022).

In this study, two collections were obtained from decaying submerged wood and dead branches of *Alangium chinense* in freshwater habitat in Southwestern China. Multi-gene phylogenetic analyses based on Maximum likelihood (ML) and Bayesian analyses along with morphological characters support the establishment of the new species. We also provided a comparative synoptic table for *Conioscypha*. This study adds new data to our knowledge on fungal diversity of freshwater streams in Southwestern China.

## Materials and methods

### Sample collection, isolation and morphological studies

Submerged decaying wood and branches were collected from Guizhou and Yunnan provinces, China. Fresh specimens were studied following the methods described by Luo et al. (2018b). The samples were incubated in plastic boxes at room temperature for one week. Micromorphological characters were observed using a stereomicroscope (SteREO Discovery.V12, Carl Zeiss Microscopy GmBH, Germany) and photographed using a Nikon ECLIPSE 80i compound microscope fitted with a NikonDS-Ri2 digital camera. Microscopic structures were measured using Tarosoft (R) Image Frame Work program and the photo-

micrographs were processed using Adobe Photoshop CS6 version 10.0 software (Adobe Systems, USA).

Single spore isolation was performed following the method described by Luo et al. (2018b). The germinated conidia were transferred to fresh PDA plates and incubated at room temperature. The specimens were dried under natural light, wrapped in absorbent paper, and placed in a Ziplock bag with mothballs. Herbarium specimens were deposited in the Herbarium of Cryptogams, Kunming Institute of Botany Academia Sinica (**KUN-HKAS**), Kunming, China, and Herbarium, University of Electronic Science and Technology (**HUEST**), Chengdu, China. The cultures were deposited in Kunming Institute of Botany, Chinese Academy of Sciences (**KUNCC**), Kunming, Yunnan, China and the University of Electronic Science and Technology Culture Collection (**UESTCC**), Chengdu, China. The novel species were registered in Faceoffungi (Jayasiri et al. 2015) and MycoBank databases (<https://www.mycobank.org/mycobank-deposit>; accessed on 22 September 2023).

### **DNA extraction, PCR amplification and sequencing**

Fresh mycelia were scraped from colonies grown on potato dextrose agar (PDA) medium. DNA extraction was carried out using DNA extraction kit following the manufacturer's instructions (TOLOBIO Plant Genomic DNA Extraction Kit, Tsingke Company, Beijing, P.R. China). PCR amplification was performed using primers pairs LR0R/LR5 (Vilgalys and Hester 1990) for the nuclear ribosomal large subunit 28S rDNA gene (LSU); NS1/NS4 (White et al. 1990) for the nuclear ribosomal small subunit 18S rDNA gene (SSU); ITS5/ITS4 (White et al. 1990) for the internal transcribed spacer rDNA region (ITS); fRPB2-5F/fRPB2-7cR (Liu et al. 1999) for the RNA polymerase second largest subunit (*RPB2*); and EF1-983F/EF1-2218R (Rehner and Buckley 2005) for the translation elongation factor 1-alpha (*TEF1- $\alpha$* ). The PCR amplification was carried out in a 25  $\mu$ L reaction volume containing 12.5  $\mu$ L of 2 $\times$  Power Taq PCR Master Mix, 1  $\mu$ L of each forward and reward primer (10  $\mu$ M), 1  $\mu$ L of genomic DNA template (30–50 ng/ $\mu$ L) and 9.5  $\mu$ L of sterilized double-distilled water. Amplifications were carried out using the BioTeke GT9612 thermocycler (Tsingke Company, Beijing, P.R. China). The PCR amplification conditions for ITS, LSU, and SSU consisted of initial denaturation at 98 °C for 3 minutes, followed by 35 cycles of denaturation at 98 °C for 20 seconds, annealing at 53 °C for 10 seconds, an extension at 72 °C for 20 seconds, and a final extension at 72 °C for 5 minutes. The PCR amplification condition for *RPB2* consisted of initial denaturation at 95 °C for 5 minutes, followed by 40 cycles of denaturation at 95 °C for 1 minute, annealing at 52 °C for 2 minutes, an extension at 72 °C for 90 seconds, and a final extension at 72 °C for 10 minutes. The amplification condition for *TEF1- $\alpha$*  consisted of initial denaturation at 94 °C for 3 minutes, followed by 35 cycles of 45 seconds at 94 °C, 50 seconds at 55 °C and 1 minute at 72 °C, and a final extension period of 10 minutes at 72 °C. Quality of PCR products were checked using 1% agarose gel electrophoresis and distinct bands were visualized in gel documentation system (Compact Desktop UV Transilluminator analyzer GL-3120). The PCR products were purified and obtained Sanger sequences by Tsingke Company, Beijing, P.R. China.

## Sequence alignments and phylogenetic analyses

The newly generated sequences were subjected to the nucleotide BLAST search via NCBI (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>; accessed on 1 September 2023) for searching the closely related taxa and confirming the correctness of the sequences. The closely related taxa of the novel species were retrieved from GenBank based on nucleotide BLAST ([www.ncbi.nlm.nih.gov/blast/](http://www.ncbi.nlm.nih.gov/blast/)) searches and recent publications (Liu et al. 2019; Bao et al. 2020). Outgroups were selected based on recently published data (Liu et al. 2019; Bao et al. 2020) (Tables 1, 2). Multiple sequence alignments were aligned with MAFFT v.7 (<http://mafft.cbrc.jp/alignment/server/index.html>; accessed on 2 September 2023) and automatically trimmed using TrimAl (<http://phylemon.bioinfo.cipf.es/utilities.html>; accessed on 2 September 2023). A combined sequence dataset was performed with SquenceMatrix v.1.7.8 (Capella et al. 2009; Vaidya

**Table 1.** Taxon names, strain numbers and GenBank accession numbers of the ITS, LSU, SSU, *RPB2* and *TEF1- $\alpha$*  sequences used in the phylogenetic analyses. Newly generated sequences are highlighted in black bold font. The ex-type strains are indicated by superscript T. “–” stands for no sequence data in GenBank.

Taxon	Voucher/Culture	GenBank accession number				
		ITS	LSU	SSU	<i>RPB2</i>	<i>TEF1-<math>\alpha</math></i>
<i>Acrogenospora alangii</i>	KUNCC 23–14553 <sup>T</sup>	<b>OR557426</b>	<b>OR553807</b>	<b>OR553806</b>	<b>OR575924</b>	<b>OR575926</b>
	UESTCC 23.0140	<b>OR578817</b>	<b>OR574254</b>	<b>OR574239</b>	<b>OR575925</b>	<b>OR575927</b>
<i>Acrogenospora aquatic</i>	MFLUCC 16–0949	–	MT340732	–	MT367160	MT367152
	MFLUCC 20–0097 <sup>T</sup>	–	–	MT340743	MT367159	MT367151
<i>Acrogenospora basalicellularispora</i>	MFLUCC 16–0992 <sup>T</sup>	–	MT340729	–	–	–
<i>Acrogenospora carmichaeliana</i>	CBS 206.36	–	MH867287	AY541482	–	–
	CBS 179.73	–	–	GU296148	–	–
	CBS 164.76	–	GU301791	GU296129	GU371748	GU349059
	FMR11021	HF677172	HF677191	–	–	–
<i>Acrogenospora guttulatispora</i>	MFLUCC 17–1674 <sup>T</sup>	–	MT340730	–	MT367157	–
<i>Acrogenospora obovoidispora</i>	MFLUCC 18–1622 <sup>T</sup>	–	MT340736	MT340747	MT367163	MT367155
<i>Acrogenospora olivaceospora</i>	MFLUCC 20–0096 <sup>T</sup>	–	MT340731	MT340742	MT367158	MT367150
<i>Acrogenospora sphaerocephala</i>	MFLUCC 16–0179	MH606233	MH606222	–	MH626448	–
<i>Acrogenospora submerse</i>	MFLUCC 18–1324 <sup>T</sup>	–	MT340735	MT340746	MT367162	MT367154
<i>Acrogenospora subprolata</i>	MFLUCC 18–1314	–	MT340739	MT340750	–	–
<i>Acrogenospora stellate</i>	AMI-SPL 1243	OP439740	OP439739	–	–	–
<i>Acrogenospora terricola</i>	PS3565	ON176299	ON176305	ON176286	–	–
	PS3417	ON176288	–	–	–	–
	PS3610 <sup>T</sup>	ON176304	ON176306	ON176287	–	–
<i>Acrogenospora thailandica</i>	MFLUCC 17–2396 <sup>T</sup>	MH606234	MH606223	MH606221	MH626449	–
<i>Acrogenospora verrucispora</i>	MFLUCC 20–0098	–	MT340737	MT340748	–	–
	MFLUCC 18–1617	–	MT340738	MT340749	MT367164	MT367156
<i>Acrogenospora yunnanensis</i>	MFLUCC 20–0099	–	MT340734	MT340745	MT367161	MT367153
	MFLUCC 18–1611 <sup>T</sup>	–	MT340733	MT340744	–	–
<i>Minutisphaera aspera</i>	DSM 29478 <sup>T</sup>	NR_154621	NG_060319	NG_065059	–	–
<i>Minutisphaera japonica</i>	HHUF30098 <sup>T</sup>	NR_119419	NG_042338	NG_064840	–	–

**Table 2.** Taxon names, strain numbers and GenBank accession numbers of the LSU, ITS, SSU and *RPB2* sequences used in the phylogenetic analyses. The newly generated sequences are highlighted in black bold font. The ex-type strains are indicated by superscript T. “–” stands for no sequence data in GenBank.

Taxon	Voucher/Culture	Gene accession numbers			
		LSU	ITS	SSU	<i>RPB2</i>
<i>Conioscypha aquatic</i>	MFLUCC 18–1333 <sup>T</sup>	MK835857	MK878383	–	MN194030
<i>Conioscypha bambusicola</i>	JCM 7245 <sup>T</sup>	NG059037	NR154660	–	–
<i>Conioscypha boutwelliae</i>	CBS 144928 <sup>T</sup>	LR025183	LR025182	–	–
<i>Conioscypha hoehnelii</i>	FMR 11592	KY853497	KY853437	HF937348	–
<i>Conioscypha japonica</i>	CBS 387.84 <sup>T</sup>	AY484514	–	JQ437438	JQ429259
<i>Conioscypha lignicola</i>	CBS 335.93	AY484513	–	JQ437439	JQ429260
<i>Conioscypha minutispora</i>	FMR 11245 <sup>T</sup>	KF924559	NR137847	HF937347	–
<i>Conioscypha nakagirii</i>	BCC77658 <sup>T</sup>	KU509985	KY859266	KU509984	KU513952
	BCC77659	KU509987	KY859267	KU509986	KU513952
<i>Conioscypha peruviana</i>	CBS 137657 <sup>T</sup>	NG058867	–	–	–
<i>Conioscypha pleiomorpha</i>	FMR 13134 <sup>T</sup>	KY853498	KY853438	–	–
<i>Conioscypha submerse</i>	MFLU 18–1639 <sup>T</sup>	MK835856	MK878382	–	–
<i>Conioscypha tenebrosa</i>	MFLU 19–0688 <sup>T</sup>	MK804508	MK804506	MK804510	MK828514
	MFLU 19–0687	MK804509	MK804507	MK804511	MK828515
<i>Conioscypha varia</i>	CBS 602.70	MH871654	MH859868	–	–
	CBS 436.70	MH871548	MH859785	–	–
	CBS 604.70	MH871656	MH859869	–	–
	CBS 603.70	MH871655	–	–	–
<i>Conioscypha verrucosa</i>	MFLUCC 18-0419 <sup>T</sup>	MN061364	MN061350	MN061352	MN061668
<b><i>Conioscypha yunnanensis</i></b>	<b>KUNCC23–13319<sup>T</sup></b>	<b>OR478379</b>	<b>OR234669</b>	<b>OR478381</b>	<b>OR487158</b>
	<b>KUNCC23–13172</b>	<b>OR478380</b>	<b>OR478183</b>	<b>OR478382</b>	<b>OR487157</b>
<i>Parafuscosporella Garethii</i>	BCC79986 <sup>T</sup>	KX958430	OK135602	KX958428	–
<i>Parafuscosporella moniliformis</i>	MFLUCC 15–0626 <sup>T</sup>	KX550895	NR152557	NG063614	–

**Abbreviation:** AMI-SPL: Collection of A. Mateos & S. De la Peña, Azores, Terceira, Portugal; **BCC:** BIOTEC Culture Collection, Thailand; **CBS:** CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands; **DSM:** Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Science Campus Braunschweig-Süd, Germany; **FMR:** Facultat de Medicina i Ciències de la Salut, Reus, Spain; **JCM:** Japan Collection of Microorganism, RIKEN BioResource Center, Japan; **HHUF:** Herbarium of Hiroshima University, Japan; **KUNCC:** Kunming Institute of Botany, Chinese Academy of Sciences Culture Collection, Kunming, Yunnan, China; **MFLU:** the herbarium of Mae Fah Luang University, Chiang Rai, Thailand; **MFLUCC:** Mae Fah Luang University Culture Collection, Chiang Rai, Thailand; **PS:** the R. L. Gilbertson Mycological Herbarium at the University of Arizona (MYCO-ARIZ); **UESTCC:** University of Electronic Science and Technology Culture Collection, Chengdu, China.

et al. 2011; Katoh and Standley 2013). Phylogenetic relationships of the new species were performed based on Maximum likelihood (ML) and Bayesian inference (BI) analyses.

Maximum likelihood (ML) was performed by RAxML-HPC2 v.8.2.12 on the XSEDE (8.2.12) tool via the CIPRES Science Gateway (<http://www.phylo.org/portal2>; accessed on 4 September 2023) (Stamatakis 2006; Miller et al. 2015) following the default setting but adjusted by setting 1,000 bootstrap replications and GTRGAMMA model of nucleotide substitution.

The evolution model for the Bayesian inference (BI) analyses was performed using MrModeltest v2.3 (Ronquist et al. 2012). GTR+I+G was selected as the best-fit model for LSU, SSU, ITS, *RPB2* and *TEF1-α* dataset. Markov Chain Monte Carlo sampling (MCMC) was computed to estimate Bayesian posterior probabilities (BPP) using MrBayes v.3.2.7 (Ronquist et al. 2012). Six simultaneous Markov chains were run for random trees for 1,000,000 generations and trees

were sampled every 200<sup>th</sup> generation. The first 10% of the total trees were set as burn-in and were discarded. The remaining trees were used to calculate Bayesian posterior probabilities (BPP) in the majority rule consensus tree (when the final average standard deviation of split frequencies reached below 0.01). Phylograms were visualized using FigTree v1.4.0 (Rambaut 2006) and rearranged in Adobe Photoshop CS6 software (Adobe Systems, USA).

The newly generated sequences were deposited in GenBank (Tables 1, 2). The final alignment and phylogenetic tree was registered in TreeBASE (<http://www.treebase.org/>) under the submission ID: 30847 (*Acrogenospora*) and ID:30689 (*Conioscypha*).

## Results


### Phylogenetic analyses

Two phylogenetic analyses were conducted to resolve the phylogenetic affinities of the two new freshwater species, one each, within the genera *Acrogenospora* (Acrogenosporaceae/ Minutisphaerales/ Dothideomycetes; Analysis 1), and the other within *Conioscypha* (Conioscyphaceae/ Conioscyphales/ Sordariomycetes; Analysis 2), as follows:

**Analysis 1:** The phylogram generated from ML analysis based on combined LSU, SSU, ITS, *RPB2* and *TEF1- $\alpha$*  sequences data was selected to represent the relationship between the new species and other known species in *Acrogenospora*. Twenty-six strains were included in the combined dataset which comprised 4,527 characters (LSU: 987 bp, SSU: 1007 bp, ITS: 535 bp, *RPB2*: 1044 bp, *TEF1- $\alpha$* : 954 bp) after alignment (including gaps). *Minutisphaera aspera* (DSM29478) and *M. japonica* (HHUF30098) were selected as the outgroup taxa. The best RAxML tree with a final likelihood value of -15211.062629 is presented in Fig. 1. RAxML analysis yielded 1,028 distinct alignment patterns and 43.09% of undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.260065, C = 0.232516, G = 0.268900, T = 0.238519, with substitution rates AC = 1.050467, AG = 3.191516, AT = 1.485302, CG = 1.086194, CT = 7.658416, GT = 1.000000; gamma distribution shape parameter alpha = 0.180026. The final average standard deviation of split frequencies at the end of total MCMC generations for BI analysis was 0.009674 (the critical value for the topological convergence diagnostic is below 0.01).

Phylogenetic analyses retrieved from ML and BI analyses were not significantly different and showed similar topologies. Phylogenetic analyses showed that our new collection (KUNCC23–14553 and UESTCC 23.0140) formed an independent subclade with strong statistical support (100% MLBS/ 1.00 BPP) and shared the same clade with *Acrogenospora terricola* and *A. thailandica* with moderate statistical support (71% MLBS/ 0.95 BPP; Fig. 1).


**Analysis 2:** The phylogram generated from ML analysis based on combined LSU, ITS, SSU and *RPB2* sequences data was selected to represent the relationship between the new species and other known species in *Conioscypha*. Twenty-three strains were included in the combined dataset which comprised 3,679 characters (LSU: 904 bp, ITS: 696 bp, SSU: 1026 bp, *RPB2*: 1053 bp) after alignment (including gaps). *Parafuscosporella garthii* (BCC79986) and *P. moniliformis* (MFLUCC 15–0626) were selected as the outgroup taxa. The



**Figure 1.** Phylogenetic tree constructed from RAxML analysis of LSU, SSU, ITS, *RPB2* and *TEF1- $\alpha$*  sequences data. Bootstrap support values for ML equal or greater than 50% and Bayesian posterior probabilities greater than 0.95 BPP are indicated at the nodes. The tree is rooted to *Minutisphaera aspera* (DSM29478) and *Minutisphaera japonica* (HHUF30098). The new isolates are in red bold.

best RAxML tree with a final likelihood value of -14285.072957 is presented in Fig. 2. RAxML analysis yielded 1,112 distinct alignment patterns and 40.07% of undetermined characters or gaps. Estimated base frequencies were as follows: A = 0.236438, C = 0.267389, G = 0.295788, T = 0.200385, with substitution rates AC = 1.738303, AG = 2.933990, AT = 1.389088, CG = 1.593182, CT = 7.181256, GT = 1.000000; gamma distribution shape parameter alpha = 0.453781. The final average standard deviation of split frequencies at the end of total MCMC generations for BI analysis was 0.003901 (the critical value for the topological convergence diagnostic is below 0.01).

Phylogenetic analyses retrieved from ML and BI analyses were not significantly different and showed similar topologies. Phylogenetic analyses showed that our new collections (KUNCC 23-13319 and KUNCC 23-13172) formed an



**Figure 2.** Phylogenetic tree constructed from RAxML analysis of LSU, ITS, SSU and *RPB2* sequences data. Bootstrap support values for ML equal or greater than 50% and Bayesian posterior probabilities greater than 0.95 BPP are indicated at the nodes. The tree is rooted to *Parafuscosporella moniliformis* (MFLUCC 15-0626) and *P. Garethii* (BCC79986). The new isolates are in red bold.

independent subclade with strong statistical support (100% MLBS/ 0.99 BPP) and clustered with *Conioscypha peruviana* and *C. minutispora*. In this study, *C. aquatica* (MFLUCC 18-1333) shared the same branch length with *C. submersa* (MFLU 18-1636) with high statistic support (99% MLBS/ 0.99 BPP). Simultaneously, *C. pleiomorpha* (FMR 13134) shares the same branch length with *C. verrucosa* (MFLUCC 18-0419) with high support (100% MLBS/ 1.00 BPP). While *C. boutwelliae* (CBS 144928) also shares the same branch length with *C. japonica* (CBS 387.84), it exhibits low statistical support in both ML and BI analyses. Therefore, the conspecific status of these species is questionable.

## Taxonomy

### ***Acrogenospora alangii* H.Z. Du & Cheewangkoon, sp. nov.**

MycoBank No: 850015

Facesoffungi Number: FoF15041

Fig. 3

**Etymology.** The epithet '*alangii*' refers to the host genus *Alangium* on which the holotype was collected.

**Holotype.** KUN-HKAS 130312.

**Description.** *Saprobic* on submerged decaying branches of *Alangium chinense* (Alangiaceae). **Asexual morph:** Hyphomycetous. **Colonies** on natural substrate, effuse, hairy, black, glistening. **Mycelium** partly semi-immersed, composed of septate, brown to dark brown, branched, smooth hyphae. **Conidiophores** 179–687 × 2.7–5.5 μm ( $\bar{x}$  = 485 × 4.2 μm, n = 20), mononematous, macronematous, solitary, erect, straight or slightly flexuous, cylindrical, unbranched, brown to dark brown, paler toward apex, septate, proliferating percurrently, smooth. **Conidiogenous cells** monoblastic, integrated, initially terminal, later becoming intercalary, cylindrical, smooth, pale brown. **Conidia** 15–22 × 15–23 μm ( $\bar{x}$  = 19.5 × 19 μm, n = 30) acrogenous, solitary, spherical or subspherical, truncate at base, aseptate, with apical appendages, hyaline and pale gray when young, pale to dark brown when mature, smooth. **Sexual morph:** Undetermined.

**Culture characteristics.** Conidia germinating on PDA within 24 h and germ tubes produced from the conidial base. Colonies reaching 16 mm diam at the room temperature in natural light for one month. Colonies on PDA medium dense, irregular in shape, slightly raised to umbonate or convex, surface rough, radially striated with lobate edge, fairly fluffy to floccose, white at the center, white-gray to gray sparse towards the margin; in reverse, white to white-gray at the center, with dark gray to brown-gray in the middle, white to pale yellowish at the edge, radiating outwards with irregular ring; no pigmentation on PDA.

**Material examined.** CHINA, Guizhou Province, Guiyang City, Wudang District, Xiangzhigou scenic spot, (26°46'7"N, 106°54'55"E), on dead branches of medicinal plant *Alangium chinense* (Alangiaceae) from freshwater stream, 25 February 2022, H.Z. Du, S136 (KUN-HKAS 130312, **holotype**), ex-holotype living culture = KUNCC 23–14553; *ibid.*, S136A (HUEST 23.0140, isotype), ex-isotype living culture = UESTCC 23.0140.

**Notes.** In the combined multi-locus phylogenetic analyses, *Acrogenospora alangii* formed a distinct clade with *A. terricola* and *A. thailandica* with significant support (71% MLBS/ 0.95 BPP; Fig. 1). The nucleotide base pair comparison between *A. alangii* (KUNCC 23–14553) and *A. terricola* (PS 3610) revealed the differences as 25/829 bp (3.0%) of LSU and 5/1006 bp (0.50%) of SSU. While the differences between *A. alangii* (KUNCC 23–14553) and *A. thailandica* (MFLUCC 17–2396) showed 30/834 bp (3.6%) of LSU and 2/1029 bp (0.2%) of SSU and 131/1045 bp (12.5%) of RPB2. *Acrogenospora alangii* can be distinguished from *A. terricola* in having conidia that are hyaline to pale gray when young, becoming pale brown to dark brown when mature, while *A. terricola* has olive green to dark brown conidia. Additionally, *A. thailandica* differs from *A. alangii* in having deep brown to black conidia (Hyde et al. 2019; Harrington et al. 2022). Furthermore, *A. alangii* also differs from the type species *A. sphaerocephala* in conidial color which is dark reddish brown, or pale to mid brown in *A. sphaerocephala* (Hughes 1978). Both *A. alangii* and *A. guizhouensis* were collected from Guizhou Province. However, morphological comparison of *A. alangii* with *A. guizhouensis* shows their differences in conidial color (hyaline, to pale gray, becoming pale brown to dark brown vs. brown) and position of conidial development (acropleurogenous vs. acrogenous) (Hyde et al. 2023).



**Figure 3.** *Acrogenospora alangii* (KUN-HKAS 130312, holotype) **a** hostplant growing near water body **b, c** colonies on host substrate **d–h** conidiophores, conidiogenous cells and conidia **i** germinating conidium **j, k** colony on PDA (up-front, down-reverse) **l, n** conidia with apical appendages **l–p** conidia. Scale bars: 100  $\mu\text{m}$  (**d, e**), 40  $\mu\text{m}$  (**f–i**), 20  $\mu\text{m}$  (**l–p**).

***Conioscypha yunnanensis* L. Li, Bhat & Phookamsak, sp. nov.**

MycoBank No: 849830

Facesoffungi Number: FoF14746

Fig. 4

**Etymology.** The specific epithet “*yunnanensis*” refers to the name of the region, Yunnan Province (China), from where the holotype was collected.


**Holotype.** KUN-HKAS 129616.

**Description.** **Saprobic** on submerged wood and unidentified twigs from freshwater habitat. **Asexual morph:** Hyphomycetous. **Colonies** on natural substrates effuse, black, glistening. **Conidiophores** reduced to conidiogenous cells. **Conidiogenous cells** phialidic, integrated, terminal, globose to subglobose, cup-shaped, percurrently proliferating in the same level, becoming multi-layered, multi-collaretted with outwardly curved edge, hyaline, smooth-walled. **Conidia** 18–26 × 17–22 µm ( $\bar{x}$  = 22 × 20 µm, n = 20), acrogenous, brown to dark brown, globose to subglobose, smooth-walled, aseptate, rounded at apex, subtruncate at base. **Sexual morph:** Undetermined.

**Culture characteristics.** Conidia germinating on PDA within 48 h and germ tubes produced from the conidial base. Colonies reaching 4.3 mm diam at room temperature in natural light for three months. Colonies on PDA medium dense to dense, circular, white and gray in the center, with packed mycelium, becoming black mycelial patch in the middle, white to cream at the margin, slightly radiating with irregular edge, radially furrowed aspect; in reverse, dark brown to black at the center, radiated with pale yellowish and dark greenish furrowed ring, white to cream at the margin with furrows aspects; no pigmentation on PDA.

**Material examined.** CHINA, Yunnan Province, Xishuangbanna (21°10'–22°40'N, 99°55'–101°50'E), on decaying submerged wood in a freshwater stream, 9 September 2022, L. Li, LILU-117-1 (KUN-HKAS 129616, **holotype**), ex-type living culture = KUNCC 23–13319; Djuanh Lake (22°29'–25°30'N, 100°16'–103°16'E), on unidentified twigs, 26 August 2022, LILU-109-1 (KUN-HKAS 129617, paratype), living culture KUNCC = 23–13172.

**Notes.** *Conioscypha yunnanensis* has close phylogenetic relationships with *C. peruviana* and *C. minutispora*. The nucleotide base pair comparison between *C. yunnanensis* (KUNCC 23–13319) and *C. peruviana* (CBS 137657) revealed 95/828 bp (11.2%) of LSU differences. The nucleotide base pair comparison between *C. yunnanensis* (KUNCC 23–13319) and *C. minutispora* (FMR 11245) revealed 57/623 bp (9.2%) of LSU, 118/554 bp (22%) of ITS and 10/937 (1.1%) of SSU differences. The new taxon shares similar morphology with *C. peruviana* in having cup-like phialidic conidiogenous cells, and brown conidia but differing by varied shapes (globose to subglobose vs. ellipsoidal to allantoid or fabiform), the size (18–26 × 17–22 µm vs. 13.5–18 × 5–8.5 µm) and absence of lipid droplets (Zelski et al. 2015). *Conioscypha yunnanensis* also resembles *C. minutispora* in having subglobose conidia but differs in the size measurement (18–26 × 17–22 µm vs. 6–9 × 5–6 µm) (Crous et al. 2014). Furthermore, *C. yunnanensis* shares similar morphology to the type species *C. lignicola* in having micronematous conidiophores and globose to subglobose conidia that are brown. However, *C. yunnanensis* differs by the absence of dark brown ring



**Figure 4.** *Conioscypha yunnanensis* (KUN-HKAS 129616, holotype) **a** host specimen **b** colonies on submerged wood **c** conidiogenous cells bearing conidia (note: arrow points = cupulate conidiogenous cells) **d, e** conidiogenous cell attached with conidia **f–m** conidia **n** germinated conidium **o, p** colony on PDA (**o** = up-front, **p** = down-reverse). Scale bars: 20  $\mu$ m (**c–n**).

surrounded in the conidia and presence of guttules periphery of conidia (Shearer 1973). The morphological comparison with other *Conioscypha* species is also provided in Table 3.

**Table 3.** Synopsis and distribution of *Conioscypha* species. The new species is indicated by black bold.

Species	Conidiophores	Conidiogenous cells	Conidia	Hosts	References
				Habitats	
				Distribution	
<i>Conioscypha aquatica</i>	–	–	Globose to subglobose, dark brown to black, 19–23 × 17–21 µm	Submerged wood Freshwater China	Luo et al. 2019
<i>C. bambusicola</i>	Semi-macronematous to micronematous, mononematous	Percurrent, cuneiform, 1.6–8.0 × 2.3–4.8 µm	Ovoid or broadly obclavate, base truncate, apex apiculate, dark brown, 11–16 × 6–10 µm	Bamboo Terrestrial Japan	Matsushima 1975
<i>C. boutwelliae</i>	Reduced to conidiogenous cells	Monoblastic, endogenous, 11.5–20.5 × 8–15 µm	Ellipsoidal, obovoid or subglobose, base truncate with a central pore of 1–1.5 µm diam, brown, pitted-wall, 10.5–21 × 8–13.5 µm	Soil Terrestrial Netherlands	Crous et al. 2018
<i>C. dimorpha</i>	–	–	Macroconidia: oblong to cylindrical, apex round, base truncate, olivaceous to brown, 8–18 × 4–6.5 µm Microconidia: subglobose to oblong, apex round, base truncate, pale brown, 2.0–3.0 × 2.0–2.5 µm	Decayed leaves Terrestrial South Africa	Matsushima 1996
<i>C. fabiformis</i>	–	–	Oblong or round, slightly curved, olivaceous, black in mass, 10–16 × 4.5–6.6 µm	Decayed leaves Terrestrial Peru	Matsushima 1993
<i>C. gracilis</i>	–	–	Ellipsoidal to flammiform, base truncate, slightly tapering towards apex, reddish brown, 8.5–9.5 × 5.5–7 µm, L/W 1.6:1	Decayed wood Terrestrial and Freshwater Denmark and Japan	Zelski et al. 2015
<i>C. hoehnelii</i>	Semi-macronematous to micronematous, mononematous	Cuneiform, cylindrical, often with a conspicuous cup-shaped, multi-collarete at the apex	Globose to subglobose or sometimes irregular, with a central pore in the inconspicuous scar at the base, brown to dark brown, 12–17 × 11–15 µm	Bark of <i>Eucalyptus</i> sp., leaf of <i>Phormium tenax</i> and unidentified wood Terrestrial UK and China	Kirk 1984; Chen and Tzean 2000
<i>C. japonica</i>	Micronematous to semi-macronematous, mononematous	Percurrent, with a multi-layered, cup-like, collarete at the apex, 4.0–17.6 × 3.2–3.8 µm	Obpyriform or subglobose, sometimes elongate, base truncate, broadly rounded at apex, smooth but with irregular pigments deposited at the periphery of the wall to give the appearance of roughness, with a pore at the point of attachment to the conidiogenous cell, entirely covered by a thin gelatinous sheath, dark brown, 9–14 × 4.5–10 µm	Scraping and hair of male dog and rotten herbaceous stem Terrestrial Japan and China	Udagawa and Toyazaki 1983; Chen and Tzean 2000
<i>C. lignicola</i>	Micronematous to semi-macronematous, mononematous	Mostly cuneiform, doliiform, percurrent, often with a cup-shaped multi-collarete, up to 16.0 µm wide at the apex, 1.6–4.8 × 2.8–6.8 µm	Obovate or sometimes subglobose, truncate at the base, with reduced lumina, smooth but dark dots deposited at the periphery, at the base with a central pore, surrounded by a dark brown ring, 11–21.6 × 10.6–16.8 µm	Balsa wood and rotten leaf of <i>Phyllostachys pubescens</i> Freshwater and terrestrial USA and China	Shearer and Motta 1973; Shearer 1973; Chen and Tzean 2000
<i>C. minutispora</i>	Reduced to conidiogenous cells	Cuneiform, percurrent, with a cup-like collarete, up to 4.0 µm wide at the apex, 7–10 × 4–5 µm	Ellipsoidal, obovoid or subglobose, apex rounded, base truncate with a central pore, dark brown, 6–9 × 5–6 µm	Submerged wood Freshwater Spain	Crous et al. 2014
<i>C. nakagirii</i>	Micronematous to semi-macronematous, mononematous	Cuneiform, cylindrical, percurrent, with a cup-shaped multi-collarete, up to 50 µm wide at the apex, 7.5–15 × 5–7.5 µm	Turbinate to pyriform, rounded at apex, truncate with a basal pore	Submerged wood Freshwater Thailand	Chuaseeharonnachai et al. 2017
<i>C. peruviana</i>	–	–	Ellipsoidal to allantoid or fabiform, containing lipid droplets, brown, 13.5–18 × 5–8.5 µm	Submerged wood Freshwater Peru	Zelski et al. 2015
<i>C. pleiomorpha</i>	Micronematous, reduced to conidiogenous cells	Monoblastic, cupulate, endogenous, multilayer-cupulate collarete after several percurrent, enteroblastic, tiny elongations, 9–12 × 13–16 µm, up to 14.0 µm	Ellipsoidal, obovoid or subglobose, base truncate with a central pore, brown, 13–18 × 12–14 µm	Dead wood Unknown habitat Spain	Hernández-Restrepo et al. 2017

Species	Conidiophores	Conidiogenous cells	Conidia	Hosts	References
				Habitats	
				Distribution	
<i>C. submersa</i>	Reduced to conidiogenous cells	–	Globose to subglobose or ovoid, pale brown, guttulate, when young, dark brown to black when mature, 17–19 × 15–17 µm	Submerged wood Freshwater China	Luo et al. 2019
<i>C. tenebrosa</i>	Micronematous, mononematous, often reduced to conidiogenous cells	Phialidic, integrated, sessile or on short conidiophores, subcylindrical, percurrently proliferating, with cup-shaped multi-collarette	Globose to subglobose, obovoid, olivaceous, aseptate, broadly rounded at apex, base subtruncate, dark brown to black when mature, 18–25 × 14–20 µm	Submerged wood Freshwater China	Liu et al. 2019
<i>C. taiwaniana</i>	Micronematous to semi-macronematous, mononematous	Cuneiform, percurrent, smooth, hyaline, with a multilayered cup-like collarette, up to 25.0 µm wide at the apex, 2.8–6.4 × 4.0–7.2 µm	Ovoid or broadly obclavate, truncate at the base, often tapering towards a point at the apex, olive brown to yellowish brown or dark brown, 14.1–20.0 × 6.4–8.0 µm	Decaying stem Terrestrial China	Hyde et al. 2020
<i>C. varia</i>	–	–	Ovoid, flammiform, naviculiform, or subellipsoid, dark brown, 8.4–15 × 5.6–8.5 µm	Balsa wood Freshwater USA	Shearer and Motta 1973; Shearer 1973
<i>C. verrucosa</i>	Macronematous, mononematous, sometimes reduced to conidiogenous cells	Monoblastic, integrated, terminal, globose to ellipsoidal, 5.5–13 × 5–11.5 µm	Globose, subglobose, ellipsoidal or obovoid, aseptate, verrucose, guttulate, dark olivaceous to with a central basal pore, dark olivaceous to dark brown, 12.5–23 × 10.5–20 µm	Submerged wood Freshwater China	Hyde et al. 2020
<i>C. yunnanensis</i>	Reduced to conidiogenous cells	Monoblastic, phialidic, integrated, terminal, globose to subglobose, cup-shaped, percurrently proliferating to the same level, multi-collarette with outwardly curved edge, hyaline, smooth-walled	Globose, subglobose, smooth-walled, aseptate, rounded at apex, subtruncate at base, brown to dark brown, 18–26 × 17–22 µm	Submerged wood Freshwater China	This study

## Discussion

Dothideomycetes and Sordariomycetes are the two largest classes of lignicolous freshwater fungi (Luo et al. 2019; Calabon et al. 2022; Shen et al. 2022). In this study, two new freshwater species belonging to Dothideomycetes and Sordariomycetes were introduced which add to the fundamental knowledge on the diversity of freshwater fungi in Southwestern China. Furthermore, an updated phylogenetic information was provided and thus we attempted to resolve the taxonomic ambiguities of the genus *Acrogenospora* (Acrogenosporaceae, Minutisphaerales, Dothideomycetes) and *Conioscypha* (Conioscyphaceae, Conioscyphales, Sordariomycetes). The study will provide a better understanding of the taxonomic boundaries of these two genera with the illustration of two new species.

*Acrogenospora* species are mostly reported from freshwater habitats (Bao et al. 2020; Hyde et al. 2023). Recent studies have revealed that more than half of the new and interesting *Acrogenospora* species were observed from freshwater habitats in China, including *A. alangii* (in this study), *A. aquatica*, *A. basalicellularispora*, *A. ellipsoidea*, *A. guizhouensis*, *A. guttulatispora*, *A. hainanensis*, *A. obovoidspora*, *A. olivaceospora*, *A. ovalia*, *A. sphaerocephala*, *A. submersa*, *A. subprolata*, *A. verrucispora* and *A. yunnanensis* (Goh et al. 1998; Ho et al. 2001; Zhu et al. 2005; Hu et al. 2010; Bao et al. 2020; Hyde et al. 2023). Previous studies revealed that the highest number of *Acrogenospora* species were reported from Yunnan Province whereas a few species have been reported from Guizhou, Hainan, Hongkong, and Xizang. This suggests a high diversity of freshwater fungi in Yunnan Province, especially of the genus *Acrogenospora*. Simultaneously, Guizhou Province is located in southwestern China that shares

similar biogeographical environments with Yunnan Province and therefore, the province may also offer a potential diversity of *Acrogenospora*.

Morphologically, species of *Acrogenospora* are distinguished from each other with difficulty and previous studies made efforts to segregate them based on shape, size, and color of the conidia and the degree of pigmentation of the conidiophores (Hughes 1978; Bao et al. 2020). A comprehensive study of *Acrogenospora* was carried out by Bao et al. (2020) who provided an updated taxonomic treatment of *Acrogenospora* and introduced seven new *Acrogenospora* species from Yunnan, China. Bao et al. (2020) and Hughes (1978) also provided a synoptic table of morphological comparison for all known *Acrogenospora* species. No significant morphological differences have been observed among known *Acrogenospora* species according to the species delineation provided by Bao et al. (2020). However, phylogenetic evidence and nucleotide pairwise comparison provide adequate justification for our species novelty following the recommendation of Jeewon and Hyde (2016).

The host association of freshwater fungi is difficult to identify. Besides, *Acrogenospora* species were mostly reported on submerged wood. Interestingly, the host associations of some *Acrogenospora* species (e.g., *A. altissima*, *A. gigantospora*, *A. sphaerocephala*, and *A. verrucispora*) have been identified. In this study, we reported *A. alangii* from freshwater habitat and associated with the medicinal plant *Alangium chinense* for the first time.

Preliminary phylogenetic analyses of a combined LSU, SSU, ITS, *RPB2* and *TEF1- $\alpha$*  sequence dataset based on Maximum likelihood (ML) (Suppl. material 1: fig. S1) showed that *Acrogenospora* sp. (JX 43) [as *Farlowiella carmichaeliana*] is sister to *A. submersa* (MFLUCC 18–1324) with low support in this study. Hyde et al. (2019) identified *Acrogenospora* sp. (JX 43) as *A. thailandica* based on phylogenetic evidence of a combined LSU, SSU and ITS sequence dataset. However, we have rechecked the sequences of *Acrogenospora* sp. (JX 43) via NCBI nucleotide BLAST search. The nucleotide BLAST search of LSU (KF836062) showed the similarity of this strain to *Chaetomium globosum* (CBS 828.73) with 100% similarity (Identities: 894/894 bp with no gap), of SSU (KF836061) showed 100% similarity to *A. thailandica* (MFLUCC 17–2396) (Identities: 1025/1025 bp with no gap), and of ITS (KF836060) showed 96.31% similarity to *Camposporium cambrense* (CBS 132486) (Identities: 496/515 bp with 5 gaps). As the nucleotide BLAST results showed three different genes of *Acrogenospora* sp. (JX 43) aligning in different genera, we excluded this strain from our analysis to avoid misidentification.

Present phylogenetic analyses also showed that *Acrogenospora carmichaeliana* (CBS 206.36) formed a separated clade with other strains of *A. carmichaeliana* (CBS 164.76, CBS 179.73, FMR 11021). *Acrogenospora carmichaeliana* (CBS 206.36) was identified as *Farlowiella carmichaeliana* (sexual morph) by E.W. Mason ([https://wi.knaw.nl/fungal\\_table](https://wi.knaw.nl/fungal_table); accessed on 17 October 2023). While strain CBS 164.76 was priorly identified as *Acrogenospora sphaerocephala* (on decaying wood in Belgium), strain CBS 179.73 was identified as *Farlowiella carmichaeliana* (on decaying wood in Germany) and FMR 11021 was identified as *Farlowiella carmichaeliana* (unknown source). Hyde et al. (2019) introduced a new species *A. thailandica* and designated the reference specimen for the type species of *Acrogenospora*, *A. sphaerocephala*. Based on their phylogenetic analyses, these four unpublished strains were identified as *A. carmichaeliana*. However, the molecular data from the ex-type strain of *A. carmi-*

*chaeliana* is unavailable. Therefore, the phylogenetic affinity of *A. carmichaeliana* remains uncertain, pending further study.

According to current reports, the species of *Conioscypha* are distributed worldwide, including Africa (*C. dimorpha*) (Matsushima 1996), United States of America (*C. lignicola*, *C. fabiformis*, *C. peruviana* and *C. varia*) (Matsushima 1993; Shearer 1973; Shearer and Motta 1973; Chen and Tzean 2000; Zelski et al. 2015), Asia (*C. aquatica*, *C. bambusicola*, *C. gracilis*, *C. hoehnelii*, *C. japonica*, *C. lignicola*, *C. nakagirii*, *C. submersa*, *C. tenebrosa*, *C. taiwaniana*, *C. verrucosa* and *C. yunnanensis*) (Shearer 1973; Shearer and Motta 1973; Matsushima 1975; Udagawa and Toyazaki 1983; Kirk 1984; Chen and Tzean 2000; Zelski et al. 2015; Chuaseeharonnachai et al. 2017; Liu et al. 2019; Luo et al. 2019; Hyde et al. 2020) and Europe (*C. gracilis*, *C. boutwelliae*, *C. hoehnelii*, *C. minutispora* and *C. pleiomorpha*) (Kirk 1984; Chen and Tzean 2000; Crous et al. 2014, 2018; Zelski et al. 2015; Hernández et al. 2017). In China, so far nine species have been reported including *C. aquatica*, *C. hoehnelii*, *C. japonica*, *C. lignicola*, *C. submersa*, *C. taiwaniana*, *C. tenebrosa* and *C. verrucosa* (Shearer 1973; Shearer and Motta 1973; Udagawa and Toyazaki 1983; Kirk 1984; Chen and Tzean 2000; Liu et al. 2019; Luo et al. 2019; Hyde et al. 2020).

Through our research on *Conioscypha yunnanensis*, it has been observed that the species of *Conioscypha* are largely indistinguishable in morphology. Hence, it has become necessary to use the potential of phylogenetic markers for clarifying their phylogenetic relationships. In this study, the single gene trees of *Conioscypha* (ITS, LSU, SSU, *RPB2*) and combined sequence datasets (LSU-ITS, LSU-ITS-SSU, and LSU-ITS-*RPB2*) were priorly conducted for comparing the reliable phylogenetic markers (Suppl. material 1: figs S2–S8). The results of these prior analyses demonstrated that the addition of *RPB2* gene could provide a better phylogenetic resolution of *Conioscypha*. Therefore, *RPB2* gene is recommended as a genetic marker for resolving phylogenetic relationships among species in *Conioscypha*.

Present phylogenetic analyses indicated that our new species formed a stable subclade independently and clustered with *Conioscypha peruviana* (CBS 137657, ex-type strain) and *C. minutispora* (FMR 11245, ex-type strain). However, the phylogenetic relationships of these three species are not well-resolved in the present study. This may be due to the available sequence data wherein only LSU gene is available for *C. peruviana* while ITS, LSU, and SSU sequences are available for *C. minutispora*. Due to the recommendation of using *RPB2* gene for delineating species of *Conioscypha*, more sequence data of *C. peruviana* and *C. minutispora* are required for providing a better phylogenetic resolution on *Conioscypha*.

Meanwhile, *Conioscypha aquatica* and *C. submersa*, introduced by Luo et al. (2019), were shown to be conspecific in the present phylogenetic analyses. Comparison of nucleotide pairwise of ITS and *TEF1- $\alpha$*  also demonstrated that these two species were not significantly different (7/530 bp (1.32%) of ITS and 10/801 bp (1.24%) of *TEF1- $\alpha$* ). However, *C. submersa* lacks *RPB2* gene that could separate these two species. Therefore, we tentatively instate these two species as a distinct species until the reliable gene (*RPB2*) is analyzed for resolving their conspecific status. Simultaneously, *C. pleiomorpha* and *C. verrucosa* have also been shown to be conspecific in the present phylogenetic analyses. However, a comparison of nucleotide pairwise of ITS and LSU demonstrated that these two species are different in 24/515 bp (4.66%) of ITS and 7/852 bp (0.82%) of LSU which is ade-

quate to justify the species' novelty. Similarly, there are only ITS and LSU sequence data of *C. pleiomorpha* currently available. These two genes may be inadequate to resolve the phylogenetic relationship of *C. pleiomorpha* and *C. verrucosa*.

The phylogenetic relationship of *Conioscypha boutwelliae* and *C. japonica* is not well-resolved in the present study. This also may be affected by the available genes used in the analyses. There are only ITS and LSU sequences available for *C. boutwelliae* whereas LSU, SSU, and *RPB2* are available for *C. japonica*. Unfortunately, the nucleotide pairwise comparison between *C. boutwelliae* and *C. japonica* could not be done due to the LSU sequence of *C. japonica* is too short (531 bp) and lacking needful genetic information compared with *C. boutwelliae* (1,053 bp). Notably, *C. boutwelliae* was introduced by Crous et al. (2018). The species was isolated from soil in the Netherlands (holotype CBS H-23743, cultures ex-type CBS 144928 = JW203008, GenBank no. LR025182 (ITS) and LR025183 (LSU), MycoBank: MB828023). The search results of LR025182 (ITS) and LR025183 (LSU) via NCBI nucleotide search brought us to the species *C. pleiomorpha*. We have rechecked the detailed source information of LR025182 (ITS) and LR025183 (LSU) and resolved that the source information belongs to *C. boutwelliae*, resulting that the sequence name of *C. boutwelliae* is incorrect in NCBI database and the name "*C. boutwelliae*" should instead be referred to as "*C. pleiomorpha*" for the GenBank no. LR025182 (ITS) and LR025183 (LSU).

## Acknowledgments

The authors would like to thank Professor Qi Zhao for his generosity in providing the experimental platform and all the cost of the experiment. Rungtiwa Phookamsak thanks the Yunnan Revitalization Talent Support Program "Young Talent" Project (grant no. YNWR-QNBJ-2020-120) the Project on Key Technology for Ecological Restoration and Green Development in Tropical Dry-Hot Valley, under the Yunnan Department of Sciences and Technology of China (grant no: 202302AE090023) for financial research support. D. Jayarama Bhat gratefully acknowledges the financial support provided under the Distinguished Scientist Fellowship Programme (DSFP), at King Saud University, Riyadh, Saudi Arabia.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This study is supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) Program (Grant No. 2019QZKK0503).

### Author contributions

Conceptualization: LL, HZD. Data curation: LL, HZD, RP. Formal analysis: LL, HZD, RP, VT. Funding acquisition: LL, RC. Investigation: LL, HZD. Methodology: LL, HZD, DJB, VT. Project administration: LL, RC. Supervision: RP, RC. Writing – original draft: LL

## Author ORCIDs

Lu Li  <https://orcid.org/0000-0003-4695-2528>

Hong-Zhi Du  <https://orcid.org/0000-0003-0350-4530>

Vinodhini Thiyagaraja  <https://orcid.org/0000-0002-8091-4579>

Darbhe Jayarama Bhat  <https://orcid.org/0000-0002-3800-5910>

Rungtiwa Phookamsak  <https://orcid.org/0000-0002-6321-8416>

Ratchadawan Cheewangkoon  <https://orcid.org/0000-0001-8576-3696>

## Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

- Bao DF, McKenzie EHC, Bhat DJ, Hyde KD, Luo ZL, Shen H-W, Su H-Y (2020) *Acrogenospora* (Acrogenosporaceae, Minutisphaerales) appears to be a very diverse genus. *Frontiers in Microbiology* 11: e1606. <https://doi.org/10.3389/fmicb.2020.01606>
- Boehm EW, Mugambi GK, Miller AN, Huhndorf SM, Marinowitz S, Spatafora JW, Schoch CL (2009) A molecular phylogenetic reappraisal of the Hysteriaceae, Mytiliniaceae and Gloniaceae (Pleosporomycetidae, Dothideomycetes) with keys to world species. *Studies in Mycology* 64: 49–83. <https://doi.org/10.3114/sim.2009.64.03>
- Calabon MS, Hyde KD, Jones EBG, Luo ZL, Dong W, Hurdeal V, Gentekaki E, Rossi W, Leonardi M, Thiyagaraja V, Lestari A, Shen HW, Bao DF, Boonyuen N, Zeng M (2022) Freshwater fungal numbers. *Fungal Diversity* 114(1): 3–235. <https://doi.org/10.1007/s13225-022-00503-2>
- Calabon MS, Hyde KD, Jones EBG, Bao DF, Bhunjun CS, Phukhamsakda C, Shen HW, Gentekaki E, Al Sharie AH, Barros J, Chandrasiri KSU, Hu DM, Hurdeal VG, Rossi W, Valle LG, Zhang H, Figueroa M, Raja HA, Seena S, Song HY, Dong W, El-Elimat T, Leonardi M, Li Y, Li YJ, Luo ZL, Ritter CD, Strongman DB, Wei Mj, Balasuriya A (2023) Freshwater fungal biology. *Mycosphere* 14(1): 195–413. <https://doi.org/10.5943/mycosphere/14/1/4>
- Capella GS, Silla MJM, Gabaldón T (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Chen JL, Tzean SS (2000) *Conioscypha taiwaniana* sp. nov. and several new records of the genus from Taiwan. *Botanical Bulletin of Academia Sinica* 41: 315–322.
- Chuaseeharonnachai C, Somrithipol S, Suetrong S, Klaysuban A, Pornputtpong N, Jones EBG, Boonyuen N (2017) *Conioscypha nakagirii*, a new species from naturally submerged wood in Thailand based on morphological and molecular data. *Mycoscience* 58(6): 424–431. <https://doi.org/10.1016/j.myc.2017.06.003>
- Crous PW, Shivas RG, Quaedvlieg W, Bank M, Zhang Y, Summerell BA, Guarro J, Wingfield MJ, Wood AR, Alfenas AC, Braun U, Cano-Lira JF, García D, Marin-Felix Y, Alvarado P, Andrade JP, Armengol J, Assefa A, Breeÿen A, Camele I, Cheewangkoon R, De Souza JT, Duong TA, Esteve-Raventós F, Fournier J, Frisullo S, García-Jiménez J, Gardienet A, Gené J, Hernández-Restrepo M, Hirooka Y, Hospenthal DR, King A, Lechat C, Lombard L, Mang SM, Marbach PAS, Marinowitz S, Marin-Felix Y, Montañó-Mata NJ, Moreno G, Perez CA, Pérez Sierra AM, Robertson JL, Roux J, Rubio E, Schumacher RK, Stchigel AM, Sutton DA, Tan YP, Thompson EH, Linde E, Walker AK, Walker DM, Wickes BL, Wong PTW, Groenewald JZ (2014) Fungal Planet description sheets: 2142–2180. *Persoonia* 32(1): 184–306. <https://doi.org/10.3767/003158514X682395>

- Crous PW, Luangsa-Ard JJ, Wingfield MJ, Carnegie AJ, Hernández-Restrepo M, Lombard L, Roux J, Barreto RW, Baseia IG, Cano-Lira JF, Martín MP, Morozova OV, Stchigel AM, Summerell BA, Brandrud TE, Dima B, García D, Giraldo A, Guarro J, Gusmão LFP, Kham-suntorn P, Noordeloos ME, Nuankaew S, Pinruan U, Rodríguez-Andrade E, Souza-Motta CM, Thangavel R, van Iperen AL, Abreu VP, Accioly T, Alves JL, Andrade JP, Bahram M, Baral HO, Barbier E, Barnes CW, Bendiksen E, Bernard E, Bezerra JDP, Bezerra JL, Bizio E, Blair JE, Bulyonkova TM, Cabral TS, Caiafa MV, Cantillo T, Colmán AA, Conceição LB, Cruz S, Cunha AOB, Darveaux BA, da Silva AL, da Silva GA, da Silva GM, da Silva RMF, de Oliveira RJV, Oliveira RL, De Souza JT, Dueñas M, Evans HC, Epifani F, Felipe MTC, Fernández-López J, Ferreira BW, Figueiredo CN, Filippova NV, Flores JA, Gené J, Ghorbani G, Gibertoni TB, Glushakova AM, Healy R, Huhndorf SM, Iturrieta-González I, Javan-Nikkhah M, Juciano RF, Jurjević Ž, Kachalkin AV, Keochanpheng K, Krisai-Greilhuber I, Li YC, Lima AA, Machado AR, Madrid H, Magalhães OMC, Marbach PAS, Melanda GCS, Miller AN, Mongkolsamrit S, Nascimento RP, Oliveira TGL, Ordoñez ME, Orzes R, Palma MA, Pearce CJ, Pereira OL, Perrone G, Peterson SW, Pham THG, Piontelli E, Pordel A, Quijada L, Raja HA, Rosas de Paz E, Ryvarden L, Saitta A, Salcedo SS, Sandoval-Denis M, Santos TAB, Seifert KA, Silva BDB, Smith ME, Soares AM, Sommai S, Sousa JO, Suetrong S, Susca A, Tedersoo L, Telleria MT, Thanakitpipattana D, Valenzuela-Lopez N, Visagie CM, Zapata M, Groenewald JZ (2018) Fungal Planet description sheets: 785–867. *Persoonia* 41(1): 238–417. <https://doi.org/10.3767/persoonia.2018.41.12>
- Dong W, Wang B, Hyde KD, McKenzie EHC, Raja HA, Tanaka K, Abdel-Wahab MA, Abdel-Aziz FA, Doilom M, Phookamsak R, Hongsanan S, Wanasinghe DN, Yu XD, Wang GN, Yang H, Yang J, Thambugala KM, Tian Q, Luo ZL, Yang JB, Miller AN, Fournier J, Boonmee S, Hu DM, Nalumpang S, Zhang H (2020) Freshwater Dothideomycetes. *Fungal Diversity* 105(1): 319–575. <https://doi.org/10.1007/s13225-020-00463-5>
- Ellis MB (1971) *Dematiaceous Hyphomycetes*. Commonwealth Mycological Institute, Kew, 114–115. <https://doi.org/10.1079/9780851986180.0000>
- Ellis MB (1972) *Dematiaceous hyphomycetes*. XI. *Mycological Papers* 131: 1–25. <https://doi.org/10.1079/9780851986180.0000>
- Feng B, Yang ZL (2018) Studies on diversity of higher fungi in Yunnan, southwestern China: A review. *Plant Diversity* 40(4): 165–171. <https://doi.org/10.1016/j.pld.2018.07.001>
- Goh TK, Hyde KD, Tsui KM (1998) The hyphomycete genus *Acrogenospora*, with two new species and two new combinations. *Mycological Research* 102(11): 1309–1315. <https://doi.org/10.1017/S0953756298006790>
- Harrington AH, Sarmiento C, Zalamea PC, Dalling JW, Davis AS, Arnold AE (2022) *Acrogenospora terricola* sp. nov., a fungal species associated with seeds of pioneer trees in the soil seed bank of a lowland forest in Panama. *International Journal of Systematic and Evolutionary Microbiology* 72(10): e005558. <https://doi.org/10.1099/ijsem.0.005558>
- Hernández RM, Gené J, Castañeda RRF, Mena PJ, Crous PW, Guarro J (2017) Phylogeny of saprobic microfungi from Southern Europe. *Studies in Mycology* 86: 53–97. <https://doi.org/10.1016/j.simyco.2017.05.002>
- Ho WH, Hyde KD, Hodgkiss IJ (2001) Fungal communities on submerged wood from streams in Brunei, Hong Kong, and Malaysia. *Mycological Research* 105(12): 1492–1501. <https://doi.org/10.1017/S095375620100507X>
- Höhnelt FXR von (1904) *Mycologische Fragmente*. (Fortsetzung) [XLII–LXIX]. *Annales mycologici* 2(1): 38–60.
- Hongsanan S, Maharachchikumbura SSN, Hyde KD, Samarakoon MC, Jeewon R, Zhao Q, Al-Sadi AM, Bahkali AH (2017) An updated phylogeny of Sordariomycetes based on

- phylogenetic and molecular clock evidence. *Fungal Diversity* 84(1): 25–41. <https://doi.org/10.1007/s13225-017-0384-2>
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD (2010) Four new freshwater fungi associated with submerged wood from Southwest Asia. *Sydowia* 62: 191–203.
- Hu DM, Liu F, Cai L (2013) Biodiversity of aquatic fungi in China. *Mycology* 4(3): 125–168. <https://doi.org/10.1080/21501203.2013.835752>
- Hughes SJ (1978) New Zealand Fungi 25. Miscellaneous species. *New Zealand Journal of Botany* 16(3): 311–370. <https://doi.org/10.1080/0028825X.1978.10425143>
- Hyde KD, Tennakoon DS, Jeewon R, Bhat DJ, Maharachchikumbura SSN, Rossi W, Leonardi M, Lee HB, Mun HY, Houbraken J, Nguyen TTT, Jeon SJ, Frisvad JC, Wanasinghe DN, Lücking R, Aptroot A, Cáceres MES, Karunarathna SC, Hongsanan S, Phookamsak R, Silva NI, Thambugala KM, Jayawardena RS, Senanayake IC, Boonmee S, Chen J, Luo ZL, Phukhamsakda C, Pereira OL, Abreu VP, Rosado AWC, Bart B, Randrianjohany E, Hofstetter V, Gibertoni TB, Soares AMS, Plautz Jr HL, Sotão HMP, Xavier WKS, Bezerra JDP, Oliveira TGL, Souza-Motta CM, Magalhães OMC, Bundhun D, Harishchandra D, Manawasinghe IS, Dong W, Zhang SN, Bao DF, Samarakoon MC, Pem D, Karunarathna A, Lin CG, Yang J, Perera RH, Kumar V, Huang SK, Dayarathne MC, Ekanayaka AH, Jayasiri SC, Xiao Y, Konta S, Niskanen T, Liimatainen K, Dai YC, Ji XH, Tian XM, Mešić A, Singh SK, Phuthacharoen K, Cai L, Sorvongxay T, Thiyagaraja V, Norphanphoun C, Chaiwan N, Lu YZ, Jiang HB, Zhang JF, Abeywickrama PD, Aluthmuhandiram JVS, Brahmanage RS, Zeng M, Chethana T, Wei D, Réblová M, Fournier J, Nekvindová J, Barbosa RN, Santos JEF, Oliveira NT, Li GJ, Ertz D, Shang QJ, Phillips AJL, Kuo CH, Camporesi E, Bulgakov TS, Lumyong S, Jones EBG, Chomnunti P, Gentekaki E, Bungartz F, Zeng XY, Fryar S, Tkalčec Z, Liang J, Li G, Wen TC, Singh PN, Gafforov Y, Promputtha I, Yasanthika E, Goonasekara ID, Zhao RL, Zhao Q, Kirk PM, Liu JK, Yan JY, Mortimer PE, Xu J, Doilom M (2019) Fungal diversity notes 1036–1150: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 96(1): 1–242. <https://doi.org/10.1007/s13225-019-00429-2>
- Hyde KD, Dong Y, Phookamsak R, Jeewon R, Bhat DJ, Jones EBG, Liu NG, Abeywickrama PD, Mapook A, Wei D, Perera RH, Manawasinghe IS, Pem D, Bundhun D, Karunarathna A, Ekanayaka AH, Bao DF, Li JF, Samarakoon MC, Chaiwan N, Lin CG, Phuthacharoen K, Zhang SN, Senanayake IC, Goonasekara ID, Thambugala KM, Phukhamsakda C, Tennakoon DS, Jiang HB, Yang J, Zeng M, Huanraluek N, Liu JK, Wijesinghe SN, Tian Q, Tibpromma S, Brahmanage RS, Boonmee S, Huang SK, Thiyagaraja V, Lu YZ, Jayawardena RS, Dong W, Yang EF, Singh SK, Singh SM, Rana S, Lad SS, Anand G, Devadatha B, Niranjana M, Sarma VV, Liimatainen K, Aguirre-Hudson B, Niskanen T, Overall A, Alvarenga RLM, Gibertoni TB, Pfigler WP, Horváth E, Imre A, Alves AL, Santos ACS, Tiago PV, Bulgakov TS, Wanasinghe DN, Bahkali AH, Doilom M, Elgorban AM, Maharachchikumbura SSN, Rajeshkumar KC, Haelewaters D, Mortimer PE, Zhao Q, Lumyong S, Xu J, Sheng J (2020) Fungal diversity notes 1151–1276: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 100(1): 5–277. <https://doi.org/10.1007/s13225-020-00439-5>
- Hyde KD, Norphanphoun C, Ma J, Yang HD, Zhang JY, Du TY, Gao Y, Gomes de Farias AR, Gui H, He SC, He YK, Li CJY, Liu XF, Lu L, Su HL, Tang X, Tian XG, Wang SY, Wei DP, Xu RF, Xu RJ, Yang Q, Yang YY, Zhang F, Zhang Q, Bahkali AH, Boonmee S, Chethana KWT, Jayawardena RS, Lu YZ, Karunarathna SC, Tibpromma S, Wang Y, Zhao Q (2023) *Mycosphere* notes 387–412 – novel species of fungal taxa from around the world. *Mycosphere* 14(1): 663–744. <https://doi.org/10.5943/mycosphere/14/1/8>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat DJ, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC (2015) The faces

- of fungi database: Fungal names linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jayasiri SC, Hyde KD, Jones EBG, Persoh D (2018) Taxonomic novelties of hysteriform Dothideomycetes. *Mycosphere* 9(4): 803–837. <https://doi.org/10.5943/mycosphere/9/4/8>
- Jeewon R, Hyde KD (2016) Establishing species boundaries and new taxa among fungi: Recommendations to resolve taxonomic ambiguities. *Mycosphere* 7(11): 1669–1677. <https://doi.org/10.5943/mycosphere/7/11/4>
- Jiang HB, Phookamsak R, Hongsanan S, Bhat DJ, Mortimer PE, Suwannarach N, Karkumyan P, Xu JC (2022) A review of bambusicolous Ascomycota in China with an emphasis on species richness in southwest China. *Studies in Fungi* 7(1): 1–20. <https://doi.org/10.48130/SIF-2022-0020>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kirk PM (1984) New or interesting microfungi XII. A new species of *Conioscypha* (Hyphomycetes). *Transactions of the British Mycological Society* 82(1): 177–178. [https://doi.org/10.1016/S0007-1536\(84\)80230-2](https://doi.org/10.1016/S0007-1536(84)80230-2)
- Li WL, Luo ZL, Liu JK, Bhat D, Bao DF, Su HY, Hyde KD (2017) Lignicolous freshwater fungi from China I: *Aquadictyospora lignicola* gen. et sp. nov. and new record of *Pseudodictyosporium wauense* from northwestern Yunnan Province. *Mycosphere* 8(10): 1587–1597. <https://doi.org/10.5943/mycosphere/8/10/1>
- Li WL, Bao DF, Bhat DJ, Su HY (2020) *Tetraploa aquatica* (Tetraplosporaaceae), a new freshwater fungal species from Yunnan Province, China. *Phytotaxa* 459(2): 181–189. <https://doi.org/10.11646/phytotaxa.459.2.8>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liu NG, Bhat DJ, Hyde KD, Liu JK (2019) *Conioscypha tenebrosa* sp. nov. (Conioscyphaaceae) from China and notes on *Conioscypha* species. *Phytotaxa* 413(2): 159–171. <https://doi.org/10.11646/phytotaxa.413.2.5>
- Luo ZL, Hyde KD, Bhat DJ, Jeewon R, Maharachchikumbura SSN, Bao DF, Li WL, Su XJ, Yang XY, Su HY (2018a) Morphological and molecular taxonomy of novel species Pleurotheciaceae from freshwater habitats in Yunnan, China. *Mycological Progress* 17(5): 511–530. <https://doi.org/10.1007/s11557-018-1377-6>
- Luo ZL, Hyde KD, Liu JK, Bhat DJ, Bao DF, Li WL, Su HY (2018b) Lignicolous freshwater fungi from China II: Novel *Distoseptispora* (Distoseptisporaceae) species from northwestern Yunnan Province and a suggested unified method for studying lignicolous freshwater fungi. *Mycosphere* 9(3): 444–461. <https://doi.org/10.5943/mycosphere/9/3/2>
- Luo ZL, Hyde KD, Liu JK, Maharachchikumbura SSN, Jeewon R, Bao DF, Bhat DJ, Lin CG, Li WL, Yang J, Liu NG, Lu YZ, Jayawardena RS, Li JF, Su HY (2019) Freshwater Sordariomycetes. *Fungal Diversity* 99(1): 451–660. <https://doi.org/10.1007/s13225-019-00438-1>
- Matsushima T (1975) *Icones microfungorum a Matsushima lectorum*. Matsushima book, 209 pp.
- Matsushima T (1993) *Matsushima Mycological Memoirs*. No. 7. Published by the author, Kobe, Japan, 131 pp.
- Matsushima T (1996) *Matsushima Mycological Memoirs*. No. 9. Matsushima Fungus collection, Kobe, Japan, 120 pp.

- Miller MA, Schwartz T, Pickett BE, He S, Klem EB, Scheuermann RH, Passarotti M, Kaufman S, O'Leary MAA (2015) Restful API for access to phylogenetic tools via the CIPRES science gateway. *Evolutionary Bioinformatics Online* 11: 43–48. <https://doi.org/10.4137/EBO.S21501>
- Rambaut A (2006) FigTree. Tree figure drawing tool version 1.3.1. Institute of Evolutionary Biology, University of Edinburgh, UK.
- Réblová M, Seifert KA (2004) *Conioscyphascus*, a new ascomycetous genus for holomorphs with *Conioscypha* anamorphs. *Studies in Mycology* 50: 95–108.
- Réblová M, Seifert KA, Fournier J, Štěpánek V (2016) Newly recognised lineages of perithecial ascomycetes: The new orders Conioscyphales and Pleurotheciales. *Persoonia* 37(1): 57–81. <https://doi.org/10.3767/003158516X689819>
- Rehner SA, Buckley EA (2005) Phylogeny inferred from nuclear ITS and *EF1-a* sequences evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* 97(1): 84–98. <https://doi.org/10.3852/mycologia.97.1.84>
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Rossman AY, Crous PW, Hyde KD, Hawksworth DL, Aptroot A, Bezerra JL, Bhat JD, Boehm E, Braun U, Boonmee S, Camporesi E, Chomnunti P, Dai DQ, D'souza MJ, Dissanayake A, Gareth Jones EB, Groenewald JZ, Hernández-Restrepo M, Hongsanan S, Jaklitsch WM, Jayawardena R, Jing LW, Kirk PM, Lawrey JD, Mapook A, McKenzie EH, Monkai J, Phillips AJ, Phookamsak R, Raja HA, Seifert KA, Senanayake I, Slippers B, Suetrong S, Taylor JE, Thambugala KM, Tian Q, Tibpromma S, Wanasinghe DN, Wijayawardene NN, Wikee S, Woudenberg JH, Wu HX, Yan J, Yang T, Zhang Y (2015) Recommended names for pleomorphic genera in Dothideomycetes. *IMA Fungus* 6(2): 507–523. <https://doi.org/10.5598/imafungus.2015.06.02.14>
- Shearer CA (1973) Fungi of the Chesapeake Bay and its tributaries II. The genus *Conioscypha*. *Mycologia* 65(1): 128–136. <https://doi.org/10.1080/00275514.1973.12019411>
- Shearer CA, Motta JJ (1973) Ultrastructure and conidiogenesis in *Conioscypha* (Hyphomycetes). *Canadian Journal of Botany* 51(10): 1747–1751. <https://doi.org/10.1139/b73-226>
- Shen HW, Bao DF, Bhat DJ, Su HY, Luo ZL (2022) Lignicolous freshwater fungi in Yunnan Province, China: An overview. *Mycology* 13(2): 119–132. <https://doi.org/10.1080/21501203.2022.2058638>
- Sivanesan A (1984) The Bitunicate Ascomycetes and their anamorphs. *Strauss and Cramer GmbH. Hirschberg*, 280–282.
- Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22(21): 2688–2690. <https://doi.org/10.1093/bioinformatics/btl446>
- Su HY, Udayanga D, Luo ZL, Manamgoda DS, Zhao YC, Yang J, Liu XY, McKenzie EHC, Zhou DQ, Hyde KD (2015) Hyphomycetes from aquatic habitats in Southern China: species of *Curvularia* (Pleosporaceae) and *Phragmocephala* (Melanommataceae). *Phytotaxa* 226(3): 201–216. <https://doi.org/10.11646/phytotaxa.226.3.1>
- Su HY, Hyde KD, Maharachchikumbura SSN, Ariyawansa HA, Luo ZL, Promputtha I, Tian Q, Lin CG, Shang QJ, Zhao YC, Chai HM, Liu XY, Bahkali A, Bhat JD, McKenzie E, Zhou DQ (2016) The families Distoseptisporaceae fam. nov., Kirschsteinioteliaceae, Sporormiaceae and Torulaceae, with new species from freshwater in Yunnan Province, China. *Fungal Diversity* 80(1): 375–409. <https://doi.org/10.1007/s13225-016-0362-0>

- Tan YP, Bishop HSL, Shivas RG, Cowan DA (2022) Fungal Planet description sheets: 1436–1477. *Persoonia* 49: 261–350. <https://doi.org/10.3767/persoonia.2022.49.08>
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (2018) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- Udagawa S, Toyazaki N (1983) A new species of *Conioscypha*. *Mycotaxon* 18: 131–137.
- Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27(2): 171–180. <https://doi.org/10.1111/j.1096-0031.2010.00329.x>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wan YL, Bao DF, Luo ZL, Bhat DJ, Xu YX, Su HY, Hao YE (2021) Two new species of *Minimelanolocus* (Herpotrichiellaceae, Chaetothyriales) from submerged wood in Yunnan, China. *Phytotaxa* 480(1): 45–56. <https://doi.org/10.11646/phytotaxa.480.1.4>
- Wang RX, Luo ZL, Hyde KD, Bhat DJ, Su XJ, Su HY (2016) New species and records of *Dictyocheirospora* from submerged wood in north-western Yunnan, China. *Mycosphere: Journal of Fungal Biology* 7(9): 1357–1367. <https://doi.org/10.5943/mycosphere/7/9/9>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis M, Gelfand D, Shinsky J, White T (Eds) *PCR Protocols: A Guide to Methods and Applications*. Academic Press, New York, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunaratna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa-2021. *Mycosphere* 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Zelski SE, Raja HA, Miller AN, Shearer CA (2015) *Conioscypha peruviana* sp. nov., its phylogenetic placement based on 28S rRNA gene, and a report of *Conioscypha gracilis* comb. nov. from Peru. *Mycoscience* 56(3): 319–325. <https://doi.org/10.1016/j.myc.2014.09.002>
- Zhao N, Luo ZL, Hyde KD, Su HY, Bhat DJ, Liu JK, Bao DF, Hao YE (2018) *Helminthosporium submersum* sp. nov. (Massarinaceae) from submerged wood in north-western Yunnan Province, China. *Phytotaxa* 348(4): 269–278. <https://doi.org/10.11646/phytotaxa.348.4.3>
- Zhu H, Cai L, Hyde KD, Zhang KQ (2005) A new species of *Acrogenospora* from submerged Bamboo in Yunnan, China. *Mycotaxon* 92: 383–386.

## Supplementary material 1

### Supplementary document

Authors: Lu Li, Hong-Zhi Du, Vinodhini Thiyagaraja, Darbhe Jayarama Bhat, Rungtiwa Phookamsak, Ratchadawan Cheewangkoon

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.101.115209.suppl1>