

Revision of *Immersaria* and a new lecanorine genus in Lecideaceae (lichenised Ascomycota, Lecanoromycetes)

Cong-Miao Xie¹, Li-Song Wang², Zun-Tian Zhao¹,
Yan-Yun Zhang², Xin-Yu Wang², Lu-Lu Zhang³

1 Key Laboratory of Plant Stress Research, College of Life Sciences, Shandong Normal University, Jinan, Shandong, 250014, China **2** CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Heilongtan, Kunming, Yunnan, 650204, China **3** Institute of Environment and Ecology, Shandong Normal University, Jinan, Shandong, 250014, China

Corresponding authors: Xin-Yu Wang (wangxinyu@mail.kib.ac.cn), Lu-Lu Zhang (612038@sdnu.edu.cn)

Academic editor: Gerhard Rambold | Received 5 August 2021 | Accepted 17 January 2022 | Published 15 February 2022

Citation: Xie C-M, Wang L-S, Zhao Z-T, Zhang Y-Y, Wang X-Y, Zhang L-L (2022) Revision of *Immersaria* and a new lecanorine genus in Lecideaceae (lichenised Ascomycota, Lecanoromycetes). MycoKeys 87: 99–132. <https://doi.org/10.3897/mycokeys.87.72614>

Abstract

The species *Immersaria cupreoatra* has been included in *Bellemerea*. This caused us to reconsider the relationships between *Bellemerea* and the lecanorine species of *Immersaria* and to question the monophyly of *Immersaria*. Amongst 25 genera of the family Lecideaceae, most have lecideine apothecia, the exceptions being *Bellemerea* and *Koerberiella*, which have lecanorine apothecia. According to previous classifications, *Immersaria* included species with both lecanorine and lecideine apothecia. A five-loci phylogenetic tree (nrITS, nrLSU, RPB1, RPB2, and mtSSU) for Lecideaceae showed that *Immersaria* was split into two clades: firstly, all the lecideine apotheciate species and secondly, all the lecanorine apotheciate species. The latter clade was closely related to the remaining lecanorine apotheciate genera: *Bellemerea* and *Koerberiella*. Therefore, the genus concept of *Immersaria* is revised accordingly and a new lecanorine genus *Lecaimmeria* is proposed. Furthermore, four new species for *Immersaria* and seven new species and three new combinations for the new genus *Lecaimmeria* are proposed. Keys to *Immersaria* and the new genus *Lecaimmeria* are provided.

Keywords

China, generic classification, lecanorine apothecia, lichen, phylogeny, taxonomy

Introduction

The lichen genus *Immersaria* Rambold & Pietschm. (Rambold 1989) was originally split from the genus *Lecidea* Ach. in order to accommodate the species *Immersaria athroocarpa* (Ach.) Rambold & Pietschm. The genus *Immersaria* was characterised by its brown thallus with an epinecral layer, a pruinose margin and an amyloid medulla, immersed apothecia with a somewhat reduced proper margin and *Porpidia*-type asci with eight, simple, halonate ascospores (Rambold 1989). Subsequently, Calatayud and Rambold (1998) enlarged the scope of the genus by including the lecanorine species, *Immersaria mehadiana* Calat. & Rambold and *I. cupreoatra* (Nyl.) Calat. & Rambold, based on morphological characters only. Currently, eight species of *Immersaria* are known worldwide (Lücking et al. 2017), three of which have lecanorine apothecia. Four of these species were previously reported from China (Hertel 1977; Zhang et al. 2015).

The species *Immersaria cupreoatra* (Nyl.) Calat. & Rambold (= *Lecanora cupreoatra* Nyl.) was previously included in *Bellemerea* Hafellner & Cl. Roux (Clauzade and Roux 1984), then into *Immersaria* by Calatayud and Rambold (1998). This caused us to reconsider the relationships between *Bellemerea* and the lecanorine species of *Immersaria* and to question the monophyly of *Immersaria*. The family Lecideaceae Chevall originally included all the crustose lecideoid genera, but now only 25 genera have been retained. Most of these are monospecific genera or small genera with under five species (Fryday and Hertel 2014; McCune et al. 2017). Most genera in Lecideaceae have lecideine apothecia. Three exceptions are *Bellemerea*, *Immersaria* and *Koerberiella* Stein, which have lecanorine apothecia. Only *Immersaria* has both lecanorine and lecideine apothecia, according to the previous circumscription (Calatayud and Rambold 1998; Valadbeigi et al. 2011). Calatayud and Rambold (1998) indicated that the presence of “two types of ascomata” represent different stages of ontogeny. However, there was no molecular evidence that could clarify the species-level phylogenetic relationships within *Immersaria*. In the two-loci phylogenetic tree of Buschbom and Mueller (2004), the lecideine species *Immersaria usbekica* (Hertel) M. Barbero, Nav.-Ros. & Cl. Roux was related to *Lecidea tessellata* Flörke. However, because only two loci of one lecideine species were included, this tree was insufficient to clarify the relationship of the lecanorine apotheciate species in *Immersaria*.

In this study, a phylogenetic tree of Lecideaceae, based on five loci, is established in order to verify the monophyly of *Immersaria*. The results show that *Immersaria* is split into two clades. One clade includes all the lecideine apotheciate species, which is sister to *Lecidea tessellata*, *L. auriculata* Th. Fr., *Cyclohymania epilithica* McCune & M.J. Curtis and the *Porpidia albocaerulescens* group and the *Porpidia speirea* group. The second clade contains all the lecanorine apotheciate species and is closely related to the rest of the lecanorine apotheciate genera within this family: *Bellemerea* and *Koerberiella*. Therefore, the genus concept of *Immersaria* is revised, retaining only the species with lecideine apothecia. The lecanorine species of *Immersaria* are excluded and proposed as a new genus, *Lecaimmeria* C.M. Xie, Lu L. Zhang & Li S. Wang. Furthermore, four

new species for *Immersaria* and seven new species and three new combinations for the new genus *Lecaimmeria* are proposed, based on the four-loci phylogenetic trees. Keys to *Immersaria* and the new genus are provided below.

Methods

Morphological analysis

All the materials for this study were collected in mainland China, mostly from the Qinghai-Tibetan Plateau, during the authors' participation in The Second Tibetan Plateau Scientific Expedition and Research Program. These specimens were stored in the Herbarium of the Kunming Institute of Botany, Chinese Academy of Sciences (KUN) and the Lichen Section of the Botanical Herbarium, Shandong Normal University (SDNU). Type specimens were loaned from the University of Helsinki (H) and Universität Wien (WU). High-resolution photographs of type specimens were provided by the curators of H or obtained from the website Global Plants (<https://plants.jstor.org/>). Morphological descriptions were made from under a dissecting microscope COIX. Anatomical descriptions were based on observations made from hand-cut sections, mounted in water and using a NIKON microscope. Usually, twenty ascospores were measured and the values of measurement means smallest measured-largest measured, with outlying values in brackets. Photographs were captured with a NIKON Eclipse 50i microscope, equipped with a NIKON digital camera (DSFi2 high-definition colour camera head, NIKON, Japan). The specimens were tested with a 10% aqueous solution of potassium hydroxide (K), a solution of aqueous sodium hypochlorite (C) and 3% Lugol's iodine (I) in the medulla and the surface of the thallus. Secondary metabolites of all the specimens were examined by thin-layer chromatography (TLC) methods, using Solvents A, B and C, as described by Orange et al. (2001).

Phylogenetic analysis

Molecular analysis was carried out on the selected specimens. Genomic DNA was extracted from dry or fresh specimens using a DNAsecure Plant Kit (Tiangen), following the manufacturer's instructions. Five gene loci were amplified by using the following primers: ITS1F (Larena et al. 1999), ITS4 (White et al. 1990), LR0R (Rehner and Samuels 1994), LR5 (Vilgalys and Hester 1990), gRPB1a (Stiller and Hall 1997), rRPB1c (Matheny et al. 2002), RPB2-6f, RPB2-7cr (Liu et al. 1999), mrSSU1 and mrSSU3R (Zoller et al. 1999). The 25 µl PCR mixture consisted of 2 µl DNA, 1 µl of each primer, 12.5 µl 2 × Taq PCR MasterMix (Aidlab) (Taq DNA Polymerase [0.1 unit/ml]; 4 mM MgCl₂; and 0.4 mM dNTPs) and 8.5 µl ddH₂O. Conditions for PCR of nrITS, nrLSU and mtSSU were set for an initial denaturation at 94 °C for 10 min, followed by 34 cycles of denaturation at 95 °C for 45 s, annealing at 50 °C for 45 s, extension at 72 °C for 90 s and a final extension at 72 °C for 10 min. For RPB1 and

RPB2, the parameters were set to an initial denaturation at 94 °C for 10 min, followed by 34 cycles of denaturation at 95 °C for 45 s, annealing at 52 °C for 50 s, extension at 72 °C for 60 s and a final extension at 72 °C for 5 min. The PCR products were sequenced using Sanger technology by the company of Tsingke Biological Technology (Beijing).

The raw sequences were assembled and edited using SeqMan v.7.0 (DNASTar packages). Sequences extracted from new materials with each gene locus were aligned with additional sequences that were available from GenBank (Suppl. material 1: Table S1), by using MEGA v.10.0 and an online version of MAFFT v.7.0 to generate nrITS-nrLSU-RPB1-RPB2-mtSSU or nrITS-nrLSU-RPB1-RPB2 matrices. The five or four gene matrices were combined by SequenceMatrix v.1.7.8. and the concatenated alignments were estimated by PartitionFinder 2 (Lanfear et al. 2017), based on the Bayesian Information Criterion (BIC), to find the most appropriate nucleotide substitution model for each of the five loci.

Phylogenetic relationships were inferred using Bayesian Inference (BI) and Maximum Likelihood (ML). ML analyses were performed with RAxMLHPC using the general time reversible model of nucleotide substitution with the gamma model of rate heterogeneity (GTRGAMMA or GTRCAT). The analyses were run with a rapid bootstrap analysis using 1000 replicates with data partitioned. The Bayesian method was performed with MrBayes v.3.1.2 (Huelsenbeck and Ronquist 2001). Four Markov chains were run with 2 million generations for each dataset and trees were sampled every 100 generations. It was ensured that the average standard deviation of split frequencies was lower than 0.01. Posterior probabilities above 0.9 and bootstrap support above 70% were considered significant supporting values. All the trees were visualised with FigTree v. 1.4.0 (Rambaut 2012).

Results

A total of 172 sequences of the nrITS, nrLSU, RPB1, RPB2, and mtSSU were generated from 61 specimens representing 57 species. Although the five-loci tree only poorly resolved the hierarchy of genera within the family Lecideaceae and the split between the lecanorine and lecideine genera of Lecideaceae was without robust support, nonetheless the results revealed that the genus *Immersaria* was not a monophyletic lineage. Rather, it was divided into two distant and well-supported lineages: clade 1 which contained the lecideine apotheciate species and clade 2 which contained the lecanorine apotheciate species (Fig. 1).

Clade 1, together with *Amygdalaria* Norman, *Cyclohymentia* McCune & M.J. Curtis, *Lecidea* s. str. and *Porpidia* Körb. (Fig. 1) formed a well-supported clade (93%

MLBS and 0.99 PP), all of which have lecideine apothecia. *Amygdalaria*, *Porpidia* and *Lecidea* s. str. were nested together, which was consistent with the results of previous research (Buschbom and Mueller 2004; Fryday et al. 2014). However, the relationships between these genera still need further research. There was a high level of support for a monophyletic lineage of lecideine apotheciate species of *Immersaria*, with these being




Figure 1. Phylogenetic tree constructed from Maximum Likelihood analyses in Lecideaceae, based on the concatenated nrITS-nrLSU-RPB1-RPB2-mtSSU dataset. Maximum Likelihood bootstrap probabilities above 70% (left) and Bayesian Inference posterior probabilities above 0.9 (right) are given at the nodes.

sister to the *Lecidea tessellata*, *L. auriculata*, *Cyclohymania*, the *Porpidia albocaerulescens* group and the *Porpidia speirea* group. The type species of *Immersaria* (*I. athrocarpa*) was included in the lineage. Thus, only those *Immersaria* species with lecideine apothecia belong to *Immersaria* s. str. This revised concept of the genus *Immersaria* is as follows: a glossy surface of thallus with an epinecral layer, immersed lecideine apothecia with a reduced margin and *Porpidia*-type asci with halonate ascospores.

There was also a high level of support for clade 2 as a monophyletic lineage (100% MLBS and 1.00 PP), which was clustered with other genera of *Lecideales* with lecanorine apothecia: *Bellemerea* and *Koerberiella* (Fig. 1). *Bellemerea* could be distinguished from Clade 2 by its amyloid ascospores and *Koerberiella* by its adnate apothecia. Although the topology of clade 2 for *Bellemerea* and *Koerberiella* is not robust, there are conspicuous differences in their morphology and significant differences between the bases in their nucleotide sequences. Since clade 2 is monophyletic with strong support, a new genus, *Lecaimmeria*, is proposed to accommodate clade 2. The new genus has immersed lecanorine apothecia with a white margin and a distinct plectenchyma developed on top of the orange epihymenium.

Two additional phylogenetic trees were constructed, based on four loci (nrITS, nrLSU, RPB1, and RPB2), in order to assess the phylogenetic position of species within *Immersaria* and *Lecaimmeria*, respectively. The phylogenetic tree of *Immersaria* was comprised of one highly supported clade with five separate lineages, based on 105 sequences from 37 specimens (Fig. 2). All the species with brown, orange, irregular or aggregate thalli formed respective monophyletic lineages. *Immersaria shangrilaensis* C.M. Xie & Lu L. Zhang formed a well-supported clade and the aggregate areolae clearly distinguished *I. shangrilaensis* from other species. *Immersaria ferruginea* C.M. Xie & Li S. Wang also formed a well-supported clade and differed from other species by its greyish-brown thallus. It seems that *Immersaria shangrilaensis* is sister to *I. ferruginea*, but the nodes were without support. In addition, the morphology is distinct between *Immersaria shangrilaensis* and *I. ferruginea*. The robust lineage *Immersaria aurantia* C.M. Xie & Li S. Wang was distinguished by its irregular, conspicuously orange thallus and green epihymenium. *Immersaria athroocarpa* was sister to *I. venusta* C.M. Xie & Xin Y. Wang, but differed in its convex, polygon areolae and densely crowded apothecia.

The phylogenetic tree of *Lecaimmeria* was comprised of one well-supported clade with nine separate lineages, based on 140 sequences from 61 specimens (Fig. 3). “*Immersaria*” sp. and *Lecaimmeria iranica* (Valadb., Sipman & Rambold) C.M. Xie comprised the basal group. “*Immersaria*” sp. has only been recorded from Macedonia and *Lecaimmeria iranica* has been recorded from Inner Mongolia in China and from Iran. *Lecaimmeria tuberculosa* C.M. Xie & Xin Y. Wang was sister to *L. qinghaiensis* C.M. Xie & Li S. Wang, but conspicuously differed in its tuberculiform conidiomata. *Lecaimmeria mongolica* C.M. Xie & Lu L. Zhang formed a well-supported monophyletic lineage and its population was mainly recognised by its orange, irregular areolae and gyrophoric acid content. *Lecaimmeria botryoides* C.M. Xie & Li S. Wang formed a highly supported sister group to *L. orbicularis* C.M. Xie & Lu L. Zhang, *L. lygaea* C.M. Xie & Lu L. Zhang and *L. tibetica* C.M. Xie & Xin Y. Wang, but differed in its crowded apothecia. *Lecaimmeria orbicularis* formed a highly supported sister group to *L. lygaea* and *L. tibetica*, but differed in its round apothecia and the white margin of the apothecia. *Lecaimmeria lygaea* was seemingly sister to *L. tibetica* and differed in its areolae having a black margin and with a well-developed prothallus between areolae.




Figure 2. Phylogenetic tree constructed from Maximum Likelihood analyses in *Immersaria*, based on the concatenated nrITS-nrLSU-RPB1-RPB2 dataset. Maximum Likelihood bootstrap probabilities above 70% (left) and Bayesian Inference posterior probabilities above 0.9 (right) are given at the nodes. Solid brown rectangle: thallus brown; solid orange rectangle: thallus yellow brown to orange brown; solid red rectangle: thallus rusty; hollow brown rectangle: thallus pale yellow brown. Solid green circle: green epihymenium; solid grey circle: without apothecia; hollow green circular: brown epihymenium.

Discussion

Revised boundaries of *Immersaria*

Formerly, the boundaries for *Immersaria* species were: lecanorine or lecideine type of immersed apothecia, production of confluent acid and gyrophoric acid and ostiole or stellate shapes of conidiomata. However, these characters were not good characters by which to distinguish this genus. The lecanorine species *Immersaria cupreoatra* was previously included in *Bellemerea*. Based on many specimens from China, it was also discovered that the ostiole or stellate shapes of conidiomata appeared in different stages of ontogeny. The main substances produced in the genus are confluent acid and gyrophoric acid; confluent acid only occurs in lecideine species, whereas gyrophoric acid appears in lecanorine species, with the exception of one lecideine species *Immersaria usbekica*. Furthermore, these characters, the types of apothecia and the shapes of conidiomata could not be applied as proper delimitations to classify species within *Immersaria*, neither were they supported by the phylogeny.




Figure 3. Phylogenetic tree constructed from Maximum Likelihood analyses in *Lecaimmeria*, based on the concatenated nrITS-nrLSU-RPB1-RPB2 dataset. Maximum Likelihood bootstrap probabilities above 70% (left) and Bayesian Inference posterior probabilities above 0.9 (right) are given at the nodes. Solid purple rectangle: areolae margin white; solid grey rectangle: areolae margin black; hollow purple rectangle: areolae margin jade green pigmented. Solid blue circle: gyrophoric acid; hollow blue circle: none.

The five-loci based analysis (Fig. 1) was incompatible with previous circumscriptions of the genus *Immersaria*, the members of which in this study are defined by their lecideine immersed apothecia, brown areolae with an epinecral layer and brown/green epihymenium without a plectenchyma. Consequently, a new genus, *Lecaimmeria*,

is established to accommodate the excluded lecanorine species. The new taxonomic system, proposed here, revised the classification boundaries between *Immersaria* and *Lecaimmeria*, but it may still be difficult to distinguish between them in cases when apothecia are absent. In this case, they could be distinguished by the substances produced or by molecular methods.

Diagnostic traits within species of *Immersaria* and *Lecaimmeria*

Species of *Immersaria* could be identified by their different thallus colours (indicated in Fig. 2). *Immersaria ferruginea* has a conspicuously greyish-brown thallus, whereas *I. athroocarpa*, *I. aurantia*, *I. shangrilaensis* and *I. venusta* have a reddish-brown thallus. *Immersaria athroocarpa* (indicated in Fig. 2) is the species that mostly has a green epihymenium, whereas the other species mostly have a brown epihymenium. Almost all these species contain confluent acid, which is often accompanied by 2'-O-methylmicrophyllinic acid. Planaic acid, which is newly reported from this genus, is only presented in specimens of *Immersaria aurantia*, *I. shangrilaensis* and *I. venusta*. All the characters discussed above were supported by the phylogeny, thus could be used as key characters to differentiate species in *Immersaria*.

Species of the new genus *Lecaimmeria* could be delimited by the colours of their areolae and margins, the existence of an apothecial margin and usually by the lack of substances. The margin of areolae (indicated in Fig. 3) was usually white, but rarely black or jade green. *Lecaimmeria lygaea* could be easily distinguished by the black margin of the areolae. The jade green margin occurs in *Lecaimmeria tuberculosa*, which grows on Qilian jade. The areolae margin of *Lecaimmeria qinghaiensis* is white, but is occasionally pigmented with very slightly green colour. The margin of the apothecia is absent in *Lecaimmeria tuberculosa* and *L. iranica*, whereas the apothecia of the other species have white margins. Most species of *Lecaimmeria* lack secondary metabolites, while gyrophoric acid was detected only in *Lecaimmeria botryoides*, *L. iranica* and *L. mongolica* (indicated in Fig. 3). In addition, an orange thallus appeared only in *Lecaimmeria mongolica* and *L. tibetica*, whereas the remaining species were brownish.

Taxonomy

***Immersaria* Rambold & Pietschm., Bibliotheca Lichenologica 34: 239 (1989).**

Type species. *Immersaria athroocarpa* (Ach.) Rambold & Pietschm., in Rambold, Biblioth. Lichenol. 34: 240 (1989).

Description. Thallus crustose, yellow-brown, red-brown, orange-brown or brown, sometimes rust coloured, continuous; areolae irregular or tending to rectangular, with a glossy surface (*atrobrunnea*-type) caused by a layer of dead, colourless cells above the upper cortex, areolae sometimes aggregate with black

prothallus and forming larger areolae; margin pruinose; prothallus distinct at the margin of thallus or absent. Upper cortex orange pigmented; epinecral layer colourless; algal layer continuous; medulla filled with grey granules. Apothecia lecideine, immersed, sometimes aggregate, round or irregular; disc black, flat, less concave, sometimes slightly raised, often poorly developed in section, pruinose or not; margin reduced. Exciple almost absent, sometimes developed, brown. Hymenium colourless; paraphyses simple, rarely branched, anastomosing or not; epihymenium brown, green or brown green, without plectenchyma; subhymenium colourless, sometimes pale brown; hypothecium pale brown to brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores ellipsoid, halonate, non-amyloid. Conidiomata present or not, immersed, linear or stellate, black, margin pruinose; conidia bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. Confluent acid, often accompanied with 2'-O-methylmicophyllinic acid, planaic acid or no substances detected by TLC. The compound planaic acid is newly reported in this genus.

Ecology and distribution. In China, growing on bare rock, sandstone or granite, from elevations of 3800 to 4500 m in the alpine zone of west China and elevations of 1200 to 1900 m in the steppe of north China. Worldwide distribution.

Notes. Species with lecanorine apothecia were previously included in *Immersaria* (Calatayud and Rambold 1998; Valadbeigi et al. 2011), but the five-loci phylogenetic analysis excluded these species from *Immersaria*. This exclusion entails a restricted concept of the genus. *Immersaria* is now defined by its orange-brown, yellow-brown, sometimes rusty coloured thallus, the amyloid medulla, the glossy surface of areolae with a pruinose margin, the black immersed lecideine apothecia with a reduced proper margin, the brown epihymenium and the *Porpidia*-type asci with eight halonate and non-amyloid ascospores. The members of this genus occur in alpine habitats.

Species of *Sporastatia* A. Massal. might be misidentified as members of *Immersaria* because of field observations of their glossy areolae and the immersed lecideine apothecia. However, they are characterised by multi-spored asci and their yellow-brown thallus. Additionally, *Miriquidica* Hertel & Rambold resembles *Immersaria* by its glossy areolae and the lecideine apothecia, but differs in its black brown thallus, its *Lecanora*-type asci with non-halonate ascospores and often containing miriquidic acid. The immersed apothecia of *Immersaria* may resemble *Aspicilia* A. Massal. and *Acarospora* A. Massal., but *Aspicilia* has a white or grey thallus, the *Aspicilia*-type asci with non-halonate ascospores; *Acarospora* has multi-spored asci.

Although four known species, *Immersaria carbonoides* (J.W. Thomson) Esnault & Cl. Roux, *I. fuliginosa* Fryday, *I. olivacea* Calat. & Rambold and *I. usbekica*, currently lack molecular data, they are temporarily left in *Immersaria* due to their morphology which corresponds to that of *Immersaria*. Our morphological comparisons were based on high-resolution photographs of type materials and the original descriptions.

***Immersaria athroocarpa* (Ach.) Rambold & Pietschm., in Rambold, Biblioth. Lichenol. 34: 240 (1989).**

Figure 4a–e

Type. SWEDEN [no locality, no date, no collector], H9508237 (H-Ach-lectotype!–designated in Hertel 1977). High-resolution photographs seen.

Description. Thallus areolate, yellow-brown, orange-brown, continuous; areolae 0.2–1.0 mm across, often convex, regular polygons, tends to be squamulose at the margin, epruinose; margin pruinose; prothallus black, not distinct, sometimes absent. Upper cortex ca. 32.0 μm thick, yellow-brown; epinecral layer ca. 7.0 μm thick; algal layer ca. 82.0 μm thick, cells 8.0–10.0 \times 7.5 μm in diam., ellipsoid. Apothecia frequent, densely crowded, immersed, 0.3–1.3 mm in diam.; disc black, rare pruinose, flat, epruinose; margin reduced. Exciple sometimes developed, 25.0–30.0 μm wide, brown. Hymenium 100–115 μm thick, colourless; paraphyses 1.0–2.0 μm wide, branched, not anastomosing; epihymenium 20.0–25.0 μm thick, brown, rarely green; subhymenium ca. 90.0 μm thick, colourless; hypothecium pale brown to brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 17.5–20.0 \times 10.0 μm , ellipsoid, halonate. Conidiomata immersed, stellate, black, margin pruinose; conidia 7.5–10.0 \times 1.0 μm in diam., bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. Chemotype I: Confluent acid. Chemotype II: Unknown substance.

Ecology and distribution. In China, growing on granite in arid and semi-arid steppe habitats at elevations of 1200–1950 m. Worldwide distribution. This species is known from Inner Mongolia and Mt. Changbai (Hertel and Zhao 1982) in China.

Notes. The lectotype grows on siliceous rock and contains several intact apothecia. The materials from Inner Mongolia are identical with the lectotype, based on comparisons with high-resolution photographs and the description given by Hertel (1977). It is, therefore, treated as *Immersaria athroocarpa* at present. Some Inner Mongolian materials contain an unknown substance, but form a well-supported clade with other materials. *Immersaria athroocarpa* is characterised by the convex, yellow-brown areolae and the large sizes of ascospores. In this genus, only this species has ascospores up to 20.0 μm long.

Immersaria usbekica is similar to *I. athroocarpa* in its brown thallus and dense apothecia, but differs in its flat areolae, the brown epihymenium and the presence of confluent acid and gyrophoric acid. By comparison with high-resolution photographs and the original descriptions (Hertel 1977) of *Immersaria usbekica*, we discovered that previous reports of this species from China (Zhang et al. 2015) were due to misidentification. It is known from Algeria, Iran, Spain, and the USSR (Barbero et al. 1990).

Specimens examined (SDNU). CHINA. Inner Mongolia: Chifeng City, Balin Youqi, Hongshilazi, 1403.2 m elev., 44°13'N, 118°44'E, on rock, 2019, Ling Hu et al. SDNU20190035; Rongshen, Wangfengou, 1217.4 m elev., 44°16'N, 118°22'E, Ling Hu et al. SDNU20190140, SDNU20190143; Erlinba, 1915.2 m elev., 44°26'N, 118°41'E, Ling Hu et al. SDNU20190227.




Figure 4. *Immersaria athrocarpa* (a–e SDNU20190227): a–b thallus c apothecial anatomy d ascus e ascospores. *I. aurantia* (f–j KUN XY19–1290): f–g thallus h apothecial anatomy i ascus j ascospores. Scale bars: 1 mm (a–b, f–g); 20 µm (c, h); 10 µm (d–e, i–j).

***Immersaria aurantia* C.M. Xie & Li S. Wang, sp. nov.**

Mycobank No: 839738

Figure 4f–j

Etymology. The name “aurantia” refers to the orange thallus.

Type. CHINA. Tibet: Sajia Co., Mula Village, 4752 m elev., 28°40'N, 88°45'E, on rock, 28 Jun 2019, Xin-Yu Wang et al. XY19–1814 (KUN-holotype).

Description. Thallus areolate, orange, dark orange, pale orange to pale red-brown, continuous; areolae 0.7–1.3 mm across, flat, epruinose, irregular; margin thin pruinose; prothallus not seen. Upper cortex 25.0–45.0 μm thick, orange; epinecral layer (12.0–) 37.0–63.0 μm thick, uneven; algal layer 50.0–93.0 μm thick, cells 5.0–15.0 \times 5.0–10.0 μm in diam., round to ellipsoid. Apothecia frequent, scattered, immersed or isolated from areolae, 0.3–1.3 mm in diam.; disc black, flat or concave, sometimes pruinose; margin reduced. Exciple sometimes developed, ca. 30.0 μm wide, brown. Hymenium 55.0–83.0 μm thick, colourless; paraphyses 2.0–3.0 μm wide, only branched and anastomosing at apex; epihymenium ca. 20.0 μm thick, green or green-brown; subhymenium colourless, not distinct or absent; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 8.0–15.0 \times 5.0–7.5 μm in diam., ellipsoid, halonate. Conidiomata rare, immersed, oblate, black, margin white; conidia 7.5 \times 1.0 μm , bacilliform.

Chemistry. Thallus K–, C–. Medulla I + violet. Chemotype I: Confluent acid, often accompanied with 2'-O-methylmicrophyllinic acid. Chemotype II: Planaic acid. Chemotype III: none (rare).

Ecology and distribution. In China, growing on rock at elevations of 3900–4300 m in the alpine zone. This species is known from Qinghai, Sichuan Province and Tibet of China.

Notes. *Immersaria aurantia* is characterised by its distinct orange, irregular areolae and the mostly green epihymenium. *Immersaria athroocarpa* and *I. venusta* are similar to *I. aurantia*, but *I. athroocarpa* differs in the convex, regularly polygonal areolae and the more crowded apothecia; *I. venusta* differs in having yellow-brown, often rusty, cracked areolae and flat apothecia. Additionally, confluent acid and planaic acid do not appear simultaneously in *Immersaria aurantia*, whereas *I. venusta* always contains both compounds.

Specimens examined (KUN). CHINA. Qinghai Province: Banma Co., 3933 m elev., 32°40'N, 100°48'E, on rock, 2020, Li-Song Wang et al. 20–66886, 3932 m elev., Li-Song Wang et al. 20–66897; Jiuzhi Co., Baiyu Village, 4285 m elev., 33°14'N, 100°58'E, Li-Song Wang et al. 20–67809. Sichuan Province: Rantang Co., Mt. Haizi, 4223 m elev., 32°20'N, 101°25'E, on rock, 2020, Li-Song Wang et al. 20–66701, 4229 m elev., Li-Song Wang et al. 20–66693, 4217 m elev., Li-Song Wang et al. 20–66680, 4221 m elev., Li-Song Wang et al. 20–66692. Tibet: Changdu City, Mangkang Co., Luoni Village, 4145 m elev., 29°56'N, 98°33'E, on rock, 2020, Li-Song Wang et al. 20–69091, 4138 m elev., Li-Song Wang et al. 20–69091, 20–69094; Gatuo Town, 29°39'N, 98°35'E, 3831 m elev., Li-Song Wang et al. 20–69114, 3850 m elev.,

Li-Song Wang et al. 20–69122; Gongga Co., Jiangtang Town, 29°12'N, 90°38'E, 2019.7.23, 4560 m elev., Xin-Yu Wang et al. XY19–1287, 4556 m elev., XY19–1290; Sajia Co., Mula Village, 28°40'N, 88°45'E, 2019.7.28, 4752 m elev., Xin-Yu Wang et al. XY19–1814; Angren Co., Kerangla, 29°19'N, 87°01'E, 4530 m elev., Li-Song Wang et al. 19–63635.

***Immersaria ferruginea* C.M. Xie & Li S. Wang, sp. nov.**

MycoBank No: 839739

Figure 5a–c

Etymology. The name “ferruginea” refers to the rusty brown colour of the thallus.

Type. CHINA. Tibet: Changdu City, Mangkang Co., Quzika Village, 4093 m elev., 29°15'N, 98°40'E, on rock, 25 Sept 2020, Li-Song Wang et al. 20–69144 (KUN-holotype).

Description. Thallus areolate, greyish-brown, continuous; areolae 0.5–1.3 mm across, flat, less often convex, rectangular to polygonal, epruinose; margin pruinose; prothallus black, not distinct. Upper cortex 50.0–68.0 µm thick, brown; epinecral layer 17.0–40.0 µm thick; algal layer 75.0–78.0 µm thick, cells (4.0–) 7.0–13.0 µm diam., round. Apothecia frequent, densely crowded, immersed, 0.7–1.3 mm in diam.; disc black, flat, pruinose; margin pruinose, slightly raised. Exciple sometimes developed, 25.0–28.0 µm wide, brown. Hymenium 57.0–100.0 µm thick, colourless; paraphyses 1.0–3.0 µm wide, not branched, anastomosing; epihymenium 15.0–33.0 µm thick, brown; subhymenium 25.0–63.0 µm thick, colourless to pale brown, rusty or dark pink; hypothecium pale brown. Asci *Porpidia*-type, cylindrical; ascospores rare, 7.5–10.0 × 5.0 µm in diam., ellipsoid, halonate. Conidiomata not seen.

Chemistry. Thallus K–, C–. Medulla I+ violet. Confluent acid, often accompanied with 2'-O-methylmicrophyllinic acid.

Ecology and distribution. In China, growing on quartz sandstone or granite at elevations of 3800–4300 m in the alpine zone. This species is known from Sichuan Province and Tibet of China.

Notes. *Immersaria ferruginea* is characterised by its brown, rusty thallus, its densely crowded apothecia and its brown epihymenium. The morphology of *Immersaria ferruginea* resembles *I. carbonioidea*, but the latter differs in its dark black-brown thallus containing norstictic acid and black-brown hypothecium.

Specimens examined (KUN). CHINA. Sichuan Province: Rangtang Co., Mt. Haizi, 4227 m elev., 32°20'N, 101°25'E, on rock, 2020, Li-Song Wang et al. 20–66697, 4221 m elev., Li-Song Wang et al. 20–67670. Tibet: Changdu City, Mangkang Co., Quzika Village, 4093 m elev., 29°15'N, 98°40'E, Li-Song Wang et al. 20–69144, 4101, Li-Song Wang et al. 20–69146, 4122 m elev., Li-Song Wang et al. 20–69148; Gatuo Town, 3848 m elev., 29°39'N, 98°35'E, Li-Song Wang et al. 20–69105.

***Immersaria shangrilaensis* C.M. Xie & Lu L. Zhang, sp. nov.**

MycoBank No: 839741

Figure 5d–f

Etymology. The name “shangrilaensis” refers to the location at which the holotype was collected: “Shangri-La”, a county of Yunnan Province in China.

Type. CHINA. Yunnan Province: Shangri-La County, Mt. Hong Shan, 4363 m elev., 28°7'N, 99°54'E, on rock, 18 Aug 2018, Li-Song Wang et al. 18–60447 (KUN-holotype).

Description. Thallus areolate, yellow-brown, orange-brown, often appears as greyish-brown, generally heavily pruinose, continuous, 5.7–10.0 cm across; areolae aggregated by 4–14 small areolae (often surrounded by black prothallus), small areolae up to 0.1 mm across, concave or flat, irregular, pruinose; margin pruinose; prothallus black, distinct. Upper cortex 32.0–50.0 µm thick, yellow-brown granules pigmented; epinecral layer 15.0–20.0 µm thick; algal layer 47.5–65.0 µm thick, cells 7.5–8.0 × 5.0 µm in diam., ellipsoid. Apothecia frequent, crowded, immersed or isolated from areolae, 0.3–0.8 mm in diam.; disc black, concave to flat, aggregated, cracked once mature, thin pruinose; margin reduced, slightly raised. Exciple almost absent. Hymenium 100.0–138.0 µm thick, colourless; paraphyses ca. 2.5 µm wide, branched, anastomosing or not; epihymenium ca. 15.0 µm thick, brown; subhymenium ca. 55.0 µm thick, colourless; hypothecium pale brown to brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 7.0–9.0 × 3.0–4.0 µm, ellipsoid, halonate (sometimes not distinct). Conidiomata immersed, oblate, black, margin heavily pruinose; conidia 7.5 × 1.0 µm, bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. Confluent acid, planaic acid and/or 2'-O-methylmicophyllinic acid.

Ecology and distribution. In China, growing on granite at elevations of 4300–4500 m in the alpine zone. This species is known from Yunnan Province of China.

Notes. The materials of *Immersaria athroocarpa* from the Shangri-La County of Yunnan Province are morphologically identical with the specimen Hertel (1977) reported from the same locality, but differ from the lectotype in its aggregate areolae, the aggregate apothecia and the smaller size of ascospores (7.0–9.0 × 3.0–4.0 µm). Based on the phenotypic and phylogenetic results, the material from Shangri-La is treated as a new species, *Immersaria shangrilaensis*. It is characterised by its large thallus, up to 10.0 cm in diam., the aggregate areolae and apothecia and the small size of ascospores.

Specimens examined. CHINA. Yunnan Province: Shangri-La County, 4350–4500 m elev., on rock, 1915, Handel-Mazzetti no. 6945 = WU-Lichenes0037752 (WU); Mt. Hong Shan, 4363 m elev., 28°7'N, 99°54'E, on rock, 2018, Li-Song Wang et al. 18–60430 (KUN), Li-Song Wang et al. 18–60447 (KUN) 4503.1 m elev., Chun-Xiao Wang et al. SDNU20181696 (SDNU), 4361.9 m elev., Chun-Xiao Wang et al. SDNU20181675 (SDNU); Luquan Co., Mt. Jiaozixueshan, 3800 m elev., 2008, Hai-Ying Wang SDNU20082253 (SDNU); Lijiang City, Mt. Laojun, 3981 m elev., 26°37'N, 99°43'E, 2018, Li-Song Wang et al. 18–60555, 18–60602 (KUN).




Figure 5. *Immersaria ferruginea* (a–c KUN 20–69144): a–b thallus c apothecial anatomy. *I. shangrilaensis* (d–f KUN 18–60430): d–e thallus f apothecial anatomy. Scale bars: 1 mm (a–b, d–e); 20 μm (c, f).

***Immersaria venusta* C.M. Xie & Xin Y. Wang, sp. nov.**

Mycobank No: 839742

Figure 6a–d

Etymology. The name “venusta” refers to the beautiful appearance of the thallus.

Type. CHINA. Qinghai Province: Maqing Co., Xueshan Village, 4187 m elev., 34°37'N, 99°42'E, on rock, 11 Sept 2020, Li-Song Wang et al. 20–67969 (KUN-holotype).

Description. Thallus areolate, brown, orange-brown, more or less rusty, continuous; areolae 0.5–1.3 mm across, flat or slightly convex, irregular, tending to rectangular, often cracked, sometimes pruinose; margin pruinose; prothallus not seen. Upper cortex ca. 38.0 μm thick, yellow brown granules pigmented; epinecral layer ca. 12.0 μm thick; algal layer ca. 128.0 μm thick, cells 5.0–10.0 \times 5.0–7.5 μm in diam., round to ellipsoid. Apothecia frequent, crowded, immersed or isolated from areolae, 0.6–1.0 mm in diam.; disc black, flat, epruinose; margin reduced, sometimes developed. Exciple sometimes developed, ca. 35.0 μm wide, brown. Hymenium 92.0–113.0 μm thick, colourless; paraphyses ca. 2.0 μm wide, anastomosing; epihymenium 27.5–30.0 μm thick, brown; subhymenium ca. 62.0 μm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 10.0–12.5 \times 5.0–7.5 μm , ellipsoid, halonate. Conidiomata immersed, linear, black, margin pruinose; conidia not seen.

Chemistry. Thallus K–, C–. Medulla I+ violet. Chemotype I: Confluent acid, often accompanied with 2'-O-methylmicrophyllinic acid. Chemotype II: Planaic acid. Chemotype III: none (rare).

Ecology and distribution. In China, growing on quartz sandstone or granite at elevations of 3900–4300 m in the alpine zone. This species is known from Qinghai Province of China.

Notes. *Immersaria venusta* is characterised by its yellow-brown, cracked areolae, its flat apothecia and brown epihymenium. It resembles *Immersaria shangrilaensis* by its cracked areolae, but its areolae have the tendency to split into several patches, but not aggregate like those of *I. shangrilaensis*. *Immersaria athroocarpa* is similar to *I. venusta* in the brown appearance of its thallus and in forming a sister group in the phylogenetic tree, but it differs in its yellow brown thallus, convex areolae, densely crowded apothecia and larger ascospores (17.5–20.0 \times 10.0 μm). *Immersaria venusta* is also similar to *I. aurantia* (see notes for *I. aurantia*). The brown thallus of *Immersaria venusta* possibly resembles that of *I. olivacea*, but the latter differs in its simple or one-septate ascospores, pyriform conidia and dark bluish-green epihymenium.

Specimens examined (KUN). CHINA. Qinghai Province: Maqing Co., Xueshan Village, 4187 m elev., 34°37'N, 99°42'E, on rock, 2020, Li-Song Wang et al. 20–67969, 20–67965; Banma Co., Yaertang Village, 3930 m elev., 32°42'N, 100°42'E, Li-Song Wang et al. 20–66940. Sichuan Province: Shiqu Co., Xinrong Village, 4043 m elev., 32°59'N, 98°19'E, on rock, 2020, Li-Song Wang et al. 20–68802; Rangtang Co., Mt. Haizi, 4246 m elev., 32°21'N, 101°24'E, Li-Song Wang et al. 20–66721, 20–66725.

Selected additional comparative material was examined.

Bellemerea alpina (Sommerf.) Clauzade & Cl. Roux RUSSIA, Lps. Petsamo, Pummangin vuonon N-puoli, 1938, Räsänen, V., H9503269 (H); Lps. Petsamo, inter Vaitolahti et Kervanto, 1938, Räsänen, V., H9503270 (H).

Bellemerea cinereorufescens (Ach.) Clauzade & Cl. Roux FINLAND, Ob. Simo. Anteroinen. Rantakivellä, 1920, Räsänen, V., H9503267 (H); Le. Enontekiö, Kirkonkylä, 1925, Kari, L.E., H9503268 (H).

High-resolution photographs seen.

Immersaria carbonoides (J.W. Thomson) Esnault & Cl. Roux USA, Alaska, along the Pitmegea River, 15 miles upstream from Cape Sabine, 1958, Thomson, J.W., M0082171 (M-isotype!), G00126754 (G-isotype!).

Immersaria olivacea Calat. & Rambold SPAIN, Espana, Castelló: Benicàssim, Parreta Alta, 390 m elev., 1993, Calatayud, V., M0101779 (M-isotype!).

Immersaria usbekica (Hertel) M. Barbero, Nav.-Ros. & Cl. Roux ALGERIA Algerie-Atlas Tellieu, Larba, Piste de Bougara á Tablat au S-E de l'arboretum de Meindja, 1985, Esnault, J., M0101787 (M-paratype!).

***Lecaimmeria* C.M. Xie, Lu L. Zhang & Li S. Wang, gen. nov.**

MycoBank No: 839743

Etymology. The name “*Lecaimmeria*” refers to the immersed lecanorine apothecia of all known species.

Type species. *Lecaimmeria orbicularis* C.M. Xie & Lu L. Zhang, sp. nov.

Description. Thallus crustose, red-brown, orange-brown or dark brown, continuous or not; areolae irregular or tending to rectangular, with a glossy surface (*atrobrunnea*-type) caused by a layer of dead, colourless cells above the upper cortex; margin white or black; prothallus distinct at the margin of thallus or absent, sometimes developed between areolae. Upper cortex orange; epinecral layer colourless; algal layer continuous; medulla filled with grey granules. Apothecia lecanorine, immersed, round or irregular; disc red-brown, dark red-brown or dark orange-brown, flat or concave; margin present or absent, black or white, rarely green, pruinose or not. Exciple reduced, tissue at the lateral sides of the hymenium corresponding to the upper cortex and the algal layer of the vegetative areolae and to hypothecial hyphal cells when apothecia reach the margin of the areole (indicated in Figs. 7c, g, 8c, 9c, h, 10c, h). Hymenium colourless; paraphyses simple, rarely branched, anastomosing or not; epihymenium orange, orange-brown, rarely brown, with a plectenchyma. Asci *Porpidia*-type (indicated in Fig 9d), cylindrical, eight-spored; ascospores ellipsoid, halonate, non-amyloid. Conidiomata present or absent, immersed, rarely convex, linear or stellate, rarely tuberculiform; conidia bacilliform.

Chemistry. Thallus K–, C+/- . Medulla I+ violet. Gyrophoric acid, 4-O-demethylplanaiic acid or no substances detected by TLC.

Ecology and distribution. In China, growing on rock, sandstone, granite or Qilian jade (rare), from elevations of 3100 to 4800 m in the alpine zone of west China and from 1200 to 1900 m in the steppe of north China. This genus is known from China, Europe, Iran, Mongolia, Romania, Russia, and USA.

Notes. The five-loci phylogenetic analysis showed that the species with lecanorine apothecia formed a novel lineage and should be excluded from *Immersaria*; thus, they are here treated as a new genus *Lecaimmeria*. *Lecaimmeria* is distinguished from related genera by its glossy surface, orange or red-brown areolae with margins, the amyloid




Figure 6. *Immersaria venusta* (a–d KUN 20–66725): a–b thallus c apothecial anatomy d ascus. Scale bars: 1 mm (a–b); 20 μ m (c); 10 μ m (d).

medulla, the red-brown immersed lecanorine apothecia, the orange epihymenium with a plectenchyma and the *Porpidia*-type asci with eight halonate and non-amyloid ascospores. In China, the genus is distributed in alpine areas, high altitude desert-steppe areas or high latitude steppe. Almost all the species of *Lecaimmeria* grow on granite or sandstone, with the exception of one species, *L. tuberculosa*, which grows on jade. Interestingly, the margin of conidiomata and areolae of *Lecaimmeria tuberculosa* appear with heavily jade-green pruinose.

The immersed apothecia and brown thallus of *Lecaimmeria* often resemble those of *Immersaria*, but *Lecaimmeria* differs in its red-brown lecanorine apothecia, often with a white margin, their orange epihymenium with a plectenchyma and the thallus containing gyrophoric acid. This genus might be confused with *Bellemerea* by its lecanorine apothecia and the *Porpidia*-type asci with halonate ascospores, but the latter genus differs in its amyloid ascospores.

Three species, previously included in *Immersaria*, *I. cupreoatra*, *I. iranica* and *I. mehadiana*, have lecanorine apothecia, but two of these, *I. cupreoatra* and *I. mehadiana*, currently lack molecular sequences. We suggest that these three species should be transferred to *Lecaimmeria*, based on the following factors. Their morphology is consistent with *Lecaimmeria* according to molecular results and comparisons with type specimens, high-resolution photographs of the type materials and the original descriptions. One unknown “*Immersaria*” species from Macedonia is sister to *Lecaimmeria iranica* in

the phylogenetic tree (Fig. 3), but comparison with high-resolution photograph and previous records (Malíček and Mayrhofer 2017) show that it differs in its black margin of areolae and absence of gyrophoric acid. This unknown species with lecanorine apothecia is possibly a member of *Lecaimmeria*, but descriptions are lacking and the specimens were not seen. Thus, this species is temporarily retained in *Immersaria*.

***Lecaimmeria botryoides* C.M. Xie & Li S. Wang, sp. nov.**

MycoBank No: 839744

Figure 7a–d

Etymology. The name “botryoides” refers to the crowded apothecia while immature.

Type. CHINA. Sichuan Province: Aba City, Rangtang County, Haizi Mt., 4225 m elev., 32°21'N, 101°24'E, on rock, 6 Sept 2020, Li-Song Wang et al. 20–66730 (KUN-holotype).

Description. Thallus areolate, red-brown, discontinuous; areolae 0.2–1.0 mm across, flat, slightly concave or convex, pruinose, polygonal, tending to be rectangular, margins heavily pruinose. Prothallus black, distinct in the margin of thallus. Upper cortex 20.0–25.0 µm thick, brown; epinecral layer 22.0–48.0 µm thick; algal layer ca. 37.0 µm thick, cells 7.5–10.0 µm diam., round. Apothecia frequent, irregular, densely crowded while immature (3–6/areolae), aggregate once mature, immersed, 0.2–1.3 mm in diam.; disc red-brown, flat, or concave, epruinose; margin pruinose, slightly raised. Hymenium 67.0–100.0 (–155.0) µm thick, colourless; paraphyses ca. 2.0 µm wide, simple, only branched at the top, not anastomosing; epihymenium 25.0–30.0 µm thick, orange; plectenchyma 2.0–8.0 µm thick; subhymenium 17.0–38.0 µm thick, colourless; hypothecium pale brown to brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 7.5–8.0 × 4.0–6.0 µm in diam., ellipsoid, halonate. Conidiomata not seen; conidia not seen.

Chemistry. Thallus K–, C+/- . Medulla I+ violet. Chemotype I: Gyrophoric acid. Chemotype II: none.

Ecology and distribution. In China, growing on rock at elevations of 3100–4300 m in the alpine zone. This species is known from Qinghai and Sichuan Provinces of China.

Notes. *Lecaimmeria botryoides* is characterised by its discontinuous thallus, densely crowded apothecia while immature and the orange epihymenium. *Lecaimmeria orbicularis* is similar to *L. botryoides* in its red-brown thallus, but differs in its round, flat apothecia and continuous thallus. The red-brown thallus of *Lecaimmeria botryoides* resembles *L. cupreoatra*, but the latter differs in the black margin of its apothecia and its distinct black prothallus between areolae.

Specimens examined (KUN). CHINA. Qinghai Province: Banma Co., 3958 m elev., 32°40'N, 100°48'E, on rock, 2020, Li-Song Wang et al. 20–66900, 3932 m elev., Li-Song Wang et al. 20–66898, 3935 m elev., Li-Song Wang et al. 20–66891, 3178 m elev., Li-Song Wang et al. 20–66765. Sichuan Province: Rangtang Co., Mt. Haizi,

4256 m elev., 32°21'N, 101°24'E, on rock, 2020, Li-Song Wang et al. 20–66721, 4300 m elev., Li-Song Wang et al. 20–67706, 4276 m elev., Li-Song Wang et al. 20–66706, 4255 m elev., Li-Song Wang et al. 20–66707, 4274 m elev., Li-Song Wang et al. 20–66713, 4274 m elev., Li-Song Wang et al. 20–66711, 20–66705, 4225 m elev., Li-Song Wang et al. 20–66730, 4220 m elev., 32°20'N, 101°25'E, Li-Song Wang et al. 20–66683.

***Lecaimmeria cupreoatra* (Nyl.) C.M. Xie, comb. nov.**

MycoBank No: 839745

Basionyms. *Lecanora cupreoatra* Nyl., Lichens Lapponiae orientalis: 181 (1866).

Type. RUSSIA. “Medvæschiihora, ad Onegam”, 13 June 1863, Th. Simming, H9508237 (H-lectotype!).

Description. Nylander (1866) and Clauzade and Roux (1985).

Notes. The lectotype grows on siliceous rock and contains several intact apothecia. As “*Immersaria*” *cupreoatra* has lecanorine apothecia and is related to *I. lygeae* in our phylogeny, it is, therefore, transferred to *Lecaimmeria*. This species has not been correctly recorded in China (see notes for *Lecaimmeria mongolica*). The species is known from Europe, Mongolia, Russia, and USA (Calatayud and Rambold 1998).

Specimens examined (H). RUSSIA. Kl. Kurkijoki, Kuuppala, Himohirsi, 12 May 1934, Räsänen, V., H9503417, H9510194.

***Lecaimmeria iranica* (Valadb., Sipman & Rambold) C.M. Xie, comb. nov.**

MycoBank No: 839746

Basionyms. *Immersaria iranica* Valadb., Sipman & Rambold, Lichenologist 43(3): 204 (2011).

Type. IRAN. Mazandaran, Haraz Road, 20 km to Aamol, 36°17'N, 52°21'E, on calcareous rock, 1475 m, 7 Apr 2006, T. Valadbeigi 9008 (TARI-holotype; B, hb. Valadbeigi-isotype). Not seen.

Description. Valadbeigi et al. (2011).

Notes. “*Immersaria*” *iranica* has lecanorine apothecia, a distinct epinecral layer and halonate ascospores (Valadbeigi et al. 2011). The materials from China are in accordance with the materials of Iran, based on comparisons with the original descriptions and the photographs given by Valadbeigi et al. (2011). The characters of this species are consistent with the new genus and the phylogenetic results showed that it was clustered with species of *Lecaimmeria*. Therefore, it was transferred to *Lecaimmeria*. This species is currently known from Iran and China.

Specimens examined (SDNU). CHINA. Xinjiang: Urumqi, Mt. Tianshan-glacier No.1, alt. 3800 m, on rock, 2011, Z.L. Huang SDNU20126106, SDNU20129049.




Figure 7. *Lecaimmeria botryoides* (a–d KUN 20–66730): a–b thallus c apothecial anatomy. *L. lyagea* (e–h KUN 20–69070): e–f thallus g apothecial anatomy h scospores. Scale bars: 1 mm (a–b, e–f); 20 µm (c, g); 10 µm (d, h).

***Lecaimmeria lygaea* C.M. Xie & Lu L. Zhang, sp. nov.**

Mycobank No: 839747

Figure 7e–h

Etymology. The name “lygaea” refers to the dark appearance of the thallus.

Type. CHINA. Tibet: Changdu City, Mangkang County, Luoni Village, 4127 m elev., 29°56'N, 98°33'E, on rock, 24 Sept 2020, Li-Song Wang et al. 20–69072 (KUN-holotype).

Description. Thallus areolate, dark red-brown, dark brown, continuous; areolae 0.5–1.0 mm across, flat, epruinose, irregular pentagonal, sometimes rectangular, fissures between areolae often filled with black prothallus; margin black, thinly pruinose; prothallus black, developed between areolae, also distinct in the margin. Upper cortex ca. 20.0 µm thick, orange-brown; epinecral layer ca. 15.0 µm thick; algal layer ca. 50.0 µm thick, cells 7.0–13.0 µm in diam., round. Apothecia frequent, round, crowded, immersed, 0.2–0.8 mm in diam.; disc red-brown, flat, or concave, epruinose; margin black, moderately thick, pruinose, raised. Hymenium 75.0–93.0 µm thick, colourless; paraphyses ca. 2.0 µm wide, simple, unbranched, not anastomosing; epihymenium 25.0–38.0 µm thick, orange brown; plectenchyma ca. 7.0 µm thick; subhymenium 20.0–25.0 µm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 12.5–20.0 × 5.0–7.5 µm in diam., ellipsoid, halonate. Conidiomata immersed, stellate, black, margin pruinose; conidia 5.0 × 1.0 µm, bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. Unknown fatty acid by TLC.

Ecology and distribution. In China, growing on sandstone at elevations of 4000–4200 m in the alpine zone. This species is known from the Tibet Region of China.

Notes. *Lecaimmeria lygaea* is characterised by its dark brown thallus, black margin of its areolae, black prothallus which fills the fissures between areolae, dark orange apothecia and its orange brown epihymenium. *Lecaimmeria cupreoatra* and *L. mehadiana* are similar to *L. lygaea*, but *L. cupreoatra* has a discontinuous thallus, with each areola surrounded by black prothallus, dark red-brown to black-brown apothecia without a margin. *Lecaimmeria mehadiana* has areolae with a white margin, black-brown apothecia, brown epihymenium and contains 4-O-demethylplanaic acid. The phylogenetic results show that *Lecaimmeria tibetica* is the sister species to *L. lygaea*. They are similar in chemistry, but differ in its orange-brown thallus and dark orange brown apothecia.

Specimens examined (KUN). CHINA. Tibet: Changdu City, Mangkang Co., Luoni Village, 4099 m elev., 29°56'N, 98°33'E, on rock, 2020, Li-Song Wang et al. 20–69054, 4131 m elev., Li-Song Wang et al. 20–69070, 4127 m elev., Li-Song Wang et al. 20–69072, 4095 m elev., Li-Song Wang et al. 20–69053.

***Lecaimmeria mehadiana* (Calatayud & Rambold) C.M. Xie, comb. nov.**

MycoBank No: 839748

Basionyms. *Immersaria mehadiana* Calat & Rambold, Lichenologist 30(3): 233 (1998).**Type.** ROMANIA. Caras-Severin Comitat, Mehadia, Strajot Mtn., on rock, 1994, Rambold, G.W., M0101781 (M-holotype!), M0101780, M0101782, M0101783 (M-isotype!). High-resolution photographs seen.**Description.** Calatayud and Rambold (1998).**Notes.** As “*Immersaria*” *mehadiana* has lecanorine apothecia and resembles *L. lygaea* and *L. cupreoatra*, having a dark brown thallus, it is, therefore, transferred to *Lecaimmeria*. This species is characterised by its greyish prothallus, dark brown apothecia and the brown epihymenium. It is only known from Romania (Calatayud and Rambold 1998).***Lecaimmeria mongolica* C.M. Xie & Lu L. Zhang, sp. nov.**

MycoBank No: 839749

Figure 8a–d

Etymology. The name “mongolica” refers to the collection of the holotype within Inner Mongolia, an autonomous region of China.**Type.** CHINA. Inner Mongolia: Chifeng City, Balinyouqi, Han Mountain, 1445 m elev., 44°11'N, 118°44'E, on rock, 22 Jul 2019, Zun-Tian Zhao et al. SDNU20190354 (SDNU-holotype).**Description.** Thallus areolate, orange, continuous; areolae 0.4–0.8 mm across, epruinose, neatly arranged, irregular, tending to be rectangular, margin pruinose; prothallus black, not distinct. Upper cortex ca. 20.0 µm thick, brown; epinecral layer 5.0–8.0 µm thick; algal layer ca. 87.0 µm thick, cells 7.5–12.5 µm diam., round. Apothecia frequent, crowded, immersed or isolated from areolae, 0.2–0.8 mm in diam.; disc red-brown, flat, slightly convex, epruinose; margin pruinose. Hymenium 62.0–83.0 µm thick, colourless; paraphyses ca. 2.0 µm wide, unbranched, not anastomosing; epihymenium ca. 42.0 µm thick, orange; plectenchyma 5.0–10.0 µm thick; subhymenium 30.0–38.0 µm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 10.0–17.5 × 6.0–7.5 µm in diam., ellipsoid, halonate. Conidiomata immersed, oblate, rare ellipsoid, black, margin pruinose; conidia 5.0 × 1.0 µm, bacilliform.**Chemistry.** Thallus K–, C+. Medulla I+ violet. Gyrophoric acid.**Ecology and distribution.** In China, growing on granite at elevations of 1400–2000 m in steppe or mountains. This species is known from Inner Mongolia of China.**Notes.** This species was once reported as “*Immersaria*” *cupreoatra* from China (Zhang et al. 2015), but after comparing our collections with the type material, this was found to be a misclassification. Additionally, the phylogenetic results showed that these collections formed a well-supported lineage belonging to *Lecaimmeria*. Therefore, it is here treated as a new species, *Lecaimmeria mongolica*, characterised by its orange-brown thallus, red-brown apothecia with a distinct white margin and the thallus containing




Figure 8. *Lecaimmeria mongolica* (a–d SDNU20190354): a–b thallus c apothecial anatomy d ascospores. Scale bars: 1 mm (a–b); 20 μ m (c); 10 μ m (d).

gyrophoric acid. *Lecaimmeria cupreoatra* resembles *L. mongolica* by containing gyrophoric acid, but it differs in its dark black brown thallus and the black margin of its apothecia. *Lecaimmeria tibetica* is similar to *L. mongolica* in its orange thallus, but differs in its smaller, dark orange apothecia and that no substance can be detected by TLC.

Specimens examined (SDNU). CHINA. Inner Mongolia: Chifeng City, Balin Youqi, Mt. Qingyangcheng, 1445 m elev., 43°35'N, 117°30'E, on rock, 2019, Zun-Tian Zhao et al. SDNU20190350; Han Shan, 1563 m elev., 44°11'N, 118°44'E, on rock, Zun-Tian Zhao et al. SDNU20190354; A'ershan City, Mt. Jiguan, 1500 m elev., on rock, 2011, Yu-Liang Cheng SDNU20124912, 1400 m elev., Dai-Feng Jiang SDNU20124859; Ke Qi, Huanggangliang, 2000 m elev., on rock, Pan-Meng Wang SDNU20117613, Xing-Ran Kou SDNU20117399.

***Lecaimmeria orbicularis* C.M. Xie & Lu L. Zhang, sp. nov.**

Mycobank No: 839750

Figure 9a–e

Etymology. The name “orbicularis” refers to the round shape of the apothecia.

Type. CHINA. Sichuan Province: Rangtang Co., Gangmuda Village, 3800 m elev., 32°18'N, 101°3'E, on rock, 7 Sept 2020, Li-Song Wang et al. 20–66753 (KUN-holotype).

Description. Thallus areolate, red-brown, rarely orange-brown, continuous; areolae 0.2–1.0 mm across, flat, occasionally wrinkled, tending to rectangular, fissures between areolae often filled with black prothallus, margin pruinose; prothallus black, developed between areolae, also distinct in the margin. Upper cortex 42.0–58.0 μm thick, brown; epinecral layer 5.0–20.0 μm thick; algal layer 70.0–113.0 μm thick, cells 10.0–15.0 \times 7.5–10.0 μm in diam., ellipsoid to round. Apothecia frequent, scattered, immersed or isolated from areolae, 0.5–1.3 mm in diam.; disc red-brown, flat, round, epruinose; margin white, slightly raised. Hymenium 75.0–113.0 μm thick, colourless; paraphyses ca. 2.0 μm wide, simple, unbranched, not anastomosing; epihymenium 17.5–30.0 μm thick, orange; plectenchyma 5.0–15.0 μm thick; subhymenium 30.0–63.0 μm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 12.5–15.0 \times 5.0–6.0 μm , ellipsoid, halonate. Conidiomata not seen.

Chemistry. Thallus K–, C–. Medulla I+ violet. None.

Ecology and distribution. In China, growing on granite or sandstone at elevations of 3700–4200 m in the alpine zone. This species is known from Qinghai and Sichuan Provinces of China.

Notes. *Lecaimmeria orbicularis* is characterised by its orange brown thallus, neatly arranged areolae and round, flat apothecia. *Lecaimmeria botryoides* is similar to *L. orbicularis* (see notes for *L. botryoides*). *Lecaimmeria mongolica* might be confused with *L. orbicularis* due to its large apothecia with a white margin, but differs in its red-brown thallus and distribution in steppes. The red-brown thallus of *Lecaimmeria cupreoatra* resembles that of *L. orbicularis*, but differs in the black margin of its apothecia and its distinct black prothallus between areolae.

Specimens examined (KUN). CHINA. Qinghai Province: Jiuzhi Co., Nianbaoyuze, 4200 m elev., 33°14'N, 100°58'E, on rock, 2020, Li-Song Wang et al. 20–66811, 20–66829, 20–66801, 20–66826A, 20–66821, 20–66805, 20–66833, 20–66817, 20–66841; Banma Co., Nianbaoyuze, 3930 m elev., 32°40'N, 100°48'E, Li-Song Wang et al. 20–66909, 20–66908, 20–66896, 20–66886B, 20–66899, 20–66935, 20–66943; Zhiqingsongduo Town, 3712 m elev., 33°24'N, 101°25'E, Li-Song Wang et al. 20–66965; Suohurima Village, 4029 m elev., 33°23'N, 100°57'E, Li-Song Wang et al. 20–66979. Sichuan Province: Rangtang Co., Gangmuda Village, 3800 m elev., 32°18'N, 101°3'E, on rock, 2020, Li-Song Wang et al. 20–66753, 20–66750, 3793 m elev., Li-Song Wang et al. 20–66747, Shangrangtang Village, 3730 m elev., 32°16'N, 101°21'E, Li-Song Wang et al. 20–66743.

***Lecaimmeria qinghaiensis* C.M. Xie & Li S. Wang, sp. nov.**

MycoBank No: 839751

Figure 9f–i

Etymology. The name “qinghaiensis” refers to the location in which the holotype was collected, in “Qinghai”, a province of China.




Figure 9. *Lecaimmeria orbicularis* (a–e KUN 20–66753): a–b thallus c apothecial anatomy d ascus (Lugol's iodine) e ascospores. *L. qinghaiensis* (f–i KUN 20–68696): f–g thallus h apothecial anatomy i ascus. Scale bars: 1 mm (a–b, f–g); 20 μm (c, h); 10 μm (d, e, i).

Type. CHINA. Qinghai Province: Yushu City, Zaduo County, Sahuteng Town, 4634 m elev., 32°55'N, 95°46'E, on rock, 20 Sept 2020, Li-Song Wang et al. 20–68698 (KUN-holotype).

Description. Thallus areolate, yellow-brown, rusty, continuous; areolae 0.5–1.5 mm across, flat, epruinose; margin pruinose, occasionally green pigmented; prothallus black, distinct at the margin. Upper cortex 27.0–38.0 µm thick, brown; epinecral layer 12.0–20.0 µm thick; algal layer 57.0–93.0 µm thick, cells 7.5–12.5 × 5.0–12.5 µm in diam., ellipsoid to round. Apothecia frequent, immersed or isolated from areolae, round or somewhat irregular while immature, ellipsoid and tending to be rectangular or occupying the whole areolae once mature, 0.2–1.3 mm in diam.; disc brown, dark red-brown, flat, occasionally with a fissure when mature, epruinose; margin white, slightly raised. Hymenium 52.0–63.0 µm thick, colourless; paraphyses 2.0–3.0 µm wide, unbranched, not anastomosing; epihymenium 25.0–30.0 µm thick, dark orange-brown; plectenchyma 7.0–18.0 µm thick; subhymenium 50.0–63.0 µm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 8.0–15.0 × 5.0–7.5 µm in diam., ellipsoid, not distinctly halonate. Conidiomata rare, immersed, flat, slightly convex, liner, stellate, graphidoid once mature, black, margin pruinose; conidia not seen.

Chemistry. Thallus K–, C–. Medulla I+ violet. None.

Ecology and distribution. In China, growing on rock at elevations of 4600–4900 m in the alpine zone. This species is known from Qinghai Province of China.

Notes. *Lecaimmeria qinghaiensis* is characterised by the yellow-brown, rusty thallus, the red-brown apothecia often occupying the whole areolae at maturity and the dark orange-brown epihymenium. The phylogenetic results showed that *Lecaimmeria tuberculosa* is the sister species to *L. qinghaiensis* which is similar in the appearance of the thallus, but differs in the brown, never rusty thallus, the red-brown apothecia and the green, tuberculiform conidiomata. The red-brown apothecia of *Lecaimmeria qinghaiensis* resembles *L. iranica*, but differs in the rusty thallus and the white margin of the apothecia.

Specimens examined (KUN). CHINA. Qinghai Province: Zaduo Co., Sahuteng Town, 4634 m elev., 32°55'N, 95°46'E, on rock, 2020, Li-Song Wang et al. 20–68698, 4637 m elev., Li-Song Wang et al. 20–68687, 4622 m elev., Li-Song Wang et al. 20–68696; 4790 m elev., 33°31'N, 95°8'E, Xin-Yu Wang et al. 20–3115, 4791 m elev., Xin-Yu Wang et al. 20–3127; Zaqing Village, 4815 m elev., Xin-Yu Wang et al. 20–849.

***Lecaimmeria tibetica* C.M. Xie & Xin Y. Wang, sp. nov.**

MycoBank No: 839752

Figure 10a–e

Etymology. The name “tibetica” refers to the location from which the holotype was collected: “Tibet”, an autonomous region of China.

Type. CHINA. Tibet: Gongga Co., Jiangtang Town, 4557 m elev., 29°12'N, 90°38'E, on rock, 9 Sept 2019, Xin-Yu Wang et al. XY19–1291 (KUN-holotype).

Description. Thallus areolate, orange-brown, epruinose; areolae 0.3–0.5 mm across, irregular, upper surface uneven, margin lacking, pruinose; prothallus black, distinct at the margin. Upper cortex 17.0–33.0 μm thick, brown; epinecral layer 10.0–20.0 μm thick; algal layer ca. 75.0 μm thick, cells 7.0–10.0 μm diam., round. Apothecia rare, immersed or isolated from areolae, 0.2–0.5 mm in diam.; disc dark orange-brown, epruinose, flat, slightly convex; margin pruinose. Hymenium 105.0–138.0 μm thick, colourless; paraphyses ca. 2.0 μm wide, unbranched, not anastomosing; epihymenium ca. 25.0 μm thick, orange; plectenchyma ca. 12.0 μm thick; subhymenium almost absent, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 12.5–15.0 \times 5.0–6.0 μm , ellipsoid, halonate. Conidiomata immersed, oblate, black, margin pruinose; conidia 5.0 \times 1.5–2.0 μm in diam., bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. None.

Ecology and distribution. In China, growing on quartz sandstone at elevations of 4300–4600 m in the alpine zone. This species is known from Tibet, China.

Notes. *Lecaimmeria tibetica* is characterised by the orange-brown thallus, the black pigmentation of the areolae margin and the dark orange-brown and small size of the apothecia. *Lecaimmeria tibetica* is similar to *L. mongolica* (see notes for *L. mongolica*). The red-brown apothecia of *Lecaimmeria cupreoatra* resembles *L. tibetica*, but that species differs in its dark red-brown thallus and the presence of gyrophoric acid.

Specimens examined (KUN). CHINA. Tibet: Gongga Co., Jiangtang Town, 4583 m elev., 29°12'N, 90°38'E, on rock, 2019, Xin-Yu Wang et al. XY19–1288, 4557 m elev., XY19–1291, 4560 m elev., XY19–1280; Dingri Co., Zhaguozechong, 4310 m elev., 28°35'N, 86°53'E, Li-Song Wang et al. 19–64071.

***Lecaimmeria tuberculosa* C.M. Xie & Xin Y. Wang, sp. nov.**

Mycobank No: 839754

Figure 10f–i

Etymology. The name “tuberculosa” refers to the tuberculiform conidiomata.

Type. CHINA. Gansu Province: Zhangye City, Sunan Co., Along the way from Sunan to Qilian, 3928 m elev., 38°37'N, 99°28'E, on rock, 30 May 2018, Li-Song Wang et al. 18–58865 (KUN-holotype).

Description. Thallus areolate, red-brown, continuous; areolae 0.5–1.3 mm across, slightly convex, epruinose; margin pruinose, often jade-green pigmented; prothallus not distinct. Upper cortex ca. 27.0 μm thick, orange; epinecral layer up to 28.0 μm thick, uneven, sometimes absent; algal layer ca. 50.0 μm thick, cells 6.0–10.0 μm diam., round. Apothecia frequent, scattered, immersed, 0.3–0.6 mm in diam.; disc red-brown, concave, epruinose; margin absent. Hymenium 55.0–83.0 μm thick, colourless; paraphyses ca. 2.0 μm wide, unbranched, not anastomosing; epihymenium 15.0–30.0 μm thick, orange; plectenchyma ca. 5.0 μm thick, discontinuous; subhymenium ca. 38.0 μm thick, colourless; hypothecium brown. Asci *Porpidia*-type, cylindrical, eight-spored; ascospores 6.0–12.5 \times 3.0–5.0 μm in diam., ellipsoid, halonate.




Figure 10. *Lecaimmeria tibetica* (a–e KUN XY19–1288): a–b thallus c apothecial anatomy d ascus e ascospore. *L. tuberculosa* (f–i KUN 18–58864): f thallus g conidiomata h apothecial anatomy i ascospores. Scale bars: 1 mm (a–b, f–g); 20 μ m (c); 50 μ m (h); 10 μ m (d, e, i).

Conidiomata stellate, strongly convex, rarely immersed, forming tuberculiform, black, margin pruinose, jade-green pigmented; conidia $3.0\text{--}4.5 \times 1.0 \mu\text{m}$ in diam., oblong to bacilliform.

Chemistry. Thallus K–, C–. Medulla I+ violet. No substances by TLC.

Ecology and distribution. In China, growing on the Qilian jade or sandstone at elevations of 3900–4200 m in the alpine zone. This species is known from Qinghai Province and Gansu Province of China.

Notes. *Lecaimmeria tuberculosa* is characterised by its red-brown thallus, the jade-green pruinose colour at the margin of its areolae, its red-brown, concave apothecia without a proper margin and tuberculiform conidiomata. *Lecaimmeria qinghaiensis* is similar to *L. tuberculosa* (see notes for *L. qinghaiensis*). *Lecaimmeria tuberculosa* usually grows on jade and, interestingly, the margin of the conidiomata and areolae of the species are heavily jade-green pigmented. *Lecaimmeria iranica* resembles *L. tuberculosa* by the absence of an apothecial margin, but differs in its immersed conidiomata and the white margin of its areolae.

Specimens examined (KUN). CHINA. Qinghai Province: Gande Co., Qingzhen Village, 4124 m elev., $34^{\circ}11'N$, $100^{\circ}12'E$, on rock, 2020, Li-Song Wang et al. 20–68077, 4145 m elev., Li-Song Wang et al. 20–68055. Gansu Province: Zhangye City, Sunan Co., along the way from Sunan to Qilian, 3928 m elev., $38^{\circ}37'N$, $99^{\circ}28'E$, on rock, 2018, Li-Song Wang et al. 18–58856, 18–58857, 18–58865, 18–59835.

Key to species of *Immersaria* in China

- 1 Thallus greyish-brown; apothecia crowded *I. ferruginea*
- Thallus reddish-brown; apothecia rarely crowded 2
- 2 Thallus orange; areolae irregular *I. aurantia*
- Thallus not orange; areolae irregular, polygonal or rectangular 3
- 3 Thallus large, 6–10 cm across; areolae aggregated by several smaller areolae and black prothallus *I. shangrilaensis*
- Thallus smaller, 2–5 cm across; areolae not aggregated 4
- 4 Thallus areolae convex, not rusty, not cracked; ascospores over $15 \mu\text{m}$ long ..
..... *I. athroocarpa*
- Thallus areolae flat, often rusty, cracked; ascospores never over $15 \mu\text{m}$ long ..
..... *I. venusta*

Key to species of *Lecaimmeria* in China

- 1 Prothallus distinct and filling the fissures between areolae *L. lygaea*
- Prothallus only distinct at the margin 2
- 2 Thallus orange 3
- Thallus reddish-brown 4
- 3 Apothecia red-brown; containing gyrophoric acid *L. mongolica*
- Apothecia dark orange; no substance detected by TLC *L. tibetica*

4	Apothecia margin absent.....	5
–	Apothecia margin present	6
5	Areolae margin white; epihymenium brown; pycindia immersed, linear or stellate.....	<i>L. iranica</i>
–	Areolae margin green; epihymenium orange; pycindia convex, tuberculi-form	<i>L. tuberculosa</i>
6	Thallus rusty; apothecia occupy the whole areolae.....	<i>L. qinghaiensis</i>
–	Thallus not rusty; apothecia do not occupy the areolae	7
7	Apothecia irregular, crowded while immature, aggregate when mature.....	<i>L. botryooides</i>
–	Apothecia round, rarely crowded, not aggregate.....	<i>L. orbicularis</i>

Acknowledgements

We are very grateful to Dr. Saara Velmala and Dr. Leena Myllys from the University of Helsinki (H) and Dr. Water Till from the Universität Wien (WU) for the loan of type materials, to Dr. Fiona Ruth Worthy, a post-doctoral researcher from Kunming Institute of Botany, Centre for Mountain Ecosystem Studies, for carefully correcting our English and to Dr. Einar Timdal (Natural History Museum, Norway) and Dr. Alan Fryday (Michigan State University) for the very careful review and the useful comments on the manuscript. This study was supported by the National Natural Science Foundation of China (31750001, 31970022, 32170002), the Second Tibetan Plateau Scientific Expedition and Research Program (STEP) (2019QZKK0503), the Youth Innovation Promotion Association CAS (2020388) and the State Key Laboratory of Phytochemistry and Plant Resources in West China (P2020-KF08).

References

- Barbero M, Navarro-Rosinés P, Roux C (1990) *Immersaria usbekica* (Hertel) Barbero, Nav.-Ros. et Roux comb. nov.[= *Amygdalaria tellensis* Esnault et Roux] nove trovita en Europo. Bulletin de la Société Linéenne de Provence 41: 139–142.
- Buschbom J, Mueller G (2004) Resolving evolutionary relationships in the lichen-forming genus *Porpidia* and related allies (Porpidiaceae, Ascomycota). Molecular Phylogenetics and Evolution 32: 66–82. <https://doi.org/10.1016/j.ympev.2003.11.012>
- Calatayud V, Rambold G (1998) Two new species of the lichen genus *Immersaria* (Porpidiaceae). Lichenologist 30: 231–244. <https://doi.org/10.1006/lich.1997.0133>
- Clauzade G, Roux C (1984) Les genres *Aspicilia* Massal. et *Bellemerea* Hafellner & Roux. Bulletin de la Société Botanique du Centre-Ouest 15: 127–141.
- Fryday AM, Hertel H (2014) A contribution to the family Lecideaceae s. lat. (Lecanoromycetidae inc. sed., lichenized Ascomycota) in the southern subpolar region; including eight new species and some revised generic circumscriptions. Lichenologist 46(3): 389–412. <https://doi.org/10.1017/S0024282913000704>

- Fryday AM, Printzen C, Ekman S (2014) *Bryobilimbia*, a new generic name for *Lecidea hypnorum* and closely related species. *Lichenologist* 46(1): 25–37. <https://doi.org/10.1017/S0024282913000625>
- Hertel H, Zhao CF (1982) Lichens from Changbai Shan-some additions to the lichen flora of northeast China. *Lichenologist* 14: 139–152. <https://doi.org/10.1017/S0024282982000280>
- Hertel H (1977) Gesteinsbewohnende Arten der Sammelgattung *Lecidea* (Lichenes) aus Zentral-, Ost- und Sudasien. Khumbu Himal, Ergebnisse des Forschungsunternehmens Nepal Himalaya 6: 145–458.
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34(3): 772–773. <https://doi.org/10.1093/molbev/msw260>
- Larena I, Salazar O, Gonzalez V, Julian MC, Rubio V (1999) Design of a primer for ribosomal DNA internal transcribed spacer with enhanced specificity for ascomycetes. *Journal of Biotechnology* 75(2–3): 187–194. [https://doi.org/10.1016/S0168-1656\(99\)00154-6](https://doi.org/10.1016/S0168-1656(99)00154-6)
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Lücking R, Hodkinson BP, Leavitt SD (2017) The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota-approaching one thousand genera. *Bryologist* 119: 361–416. <https://doi.org/10.1639/0007-2745-119.4.361>
- Malíček J, Mayrhofer H (2017) Additions to the lichen diversity of Macedonia (FYROM) [Ergänzungen zur Flechtendiversität von Makedonien (FYROM)]. *Herzogia* 30(2): 431–444. <https://doi.org/10.13158/heia.30.2.2017.431>
- Matheny PB, Liu YJ, Ammirati JF, Hall BD (2002) Using RPB1 sequences to improve phylogenetic inference among mushrooms (Inocybe, Agaricales). *American Journal of Botany* 89: 688–698. <https://doi.org/10.3732/ajb.89.4.688>
- McCune B, Curtis MJ, Di Meglio J (2017) New taxa and a case of ephemeral spore production in Lecideaceae from western North America. *Bryologist* 120(2): 115–124. <https://doi.org/10.1639/0007-2745-120.2.115>
- Nylander W (1866) Lichenes Lapponiae orientalis. *Notiser ur Sällskapet pro Fauna et Flora Fennica Förhandlingar* 8: 101–192.
- Orange A, James PW, White FJ (2010) Microchemical methods for the identification of lichens (2nd edn.). British Lichen Society, London, 101 pp.
- Rambaut A (2012) FigTree, v. 1.4.0. Institute of Evolutionary Biology, University of Edinburgh.
- Rambold GA (1989) A monograph of the Saxicolous Lecideoid Lichens of Australia (excl. Tasmania). *Bibliotheca Lichenologica* 34: 239–240.
- Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* 98: 625–634. [https://doi.org/10.1016/S0953-7562\(09\)80409-7](https://doi.org/10.1016/S0953-7562(09)80409-7)
- Stiller JW, Hall BD (1997) The origin of red algae: Implications for plastid evolution. *PNAS* 94: 4520–4525. <https://doi.org/10.1073/pnas.94.9.4520>

- Valadbeigi T, Sipman H, Rambold G (2011) The genus *Immersaria* (Lecideaceae) in Iran, including *I. iranica* sp. nov. *Lichenologist* 43(3): 203–208. <https://doi.org/10.1017/S0024282911000077>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4239–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- White T, Bruns T, Lee S, Taylor F, White TJ, Lee SH, Taylor L, Shawetaylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis M, Gel-fand J, Sainsky J, et al. (Eds) *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Zhang LL, Hu L, Zhao XX, Zhao ZT (2015) New records of *Clauzadea* and *Immersaria* from China. *Mycotaxon*, Ithaca NY, 130 (3): 899–905. <https://doi.org/10.5248/130.899>
- Zoller S, Scheidegger C, Sperisen C (1999) PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. *Lichenologist* 31(5): 511–516. <https://doi.org/10.1006/lich.1999.0220>

Supplementary material I

Table S1

Authors: Cong-Miao Xie, Li-Song Wang, Zun-Tian Zhao, Yan-Yun Zhang, Xin-Yu Wang, Lu-Lu Zhang

Data type: docx. file

Explanation note: A list of specimens and GenBank accession number of sequences used in this study. The new sequences generated are in bold.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.87.72614.suppl1>