

Acremonium capsici* and *A. guizhouense*, two new members of *Acremonium* (Hypocreales, Sordariomycetes) isolated from the rhizosphere soil of *Capsicum annuum

Shuo-Qiu Tong¹, Lei Peng², Yong-Jun Wu¹

1 College of Life Sciences, Institute of Agro-bioengineering, Guizhou University, Guiyang 550025, China
2 Tea Academy, Guizhou University, Guiyang 550025, China

Corresponding author: Yong-Jun Wu (wyjbio@163.com)

Academic editor: Huzefa Raja | Received 2 November 2022 | Accepted 12 December 2022 | Published 4 January 2023

Citation: Tong S-Q, Peng L, Wu Y-J (2023) *Acremonium capsici* and *A. guizhouense*, two new members of *Acremonium* (Hypocreales, Sordariomycetes) isolated from the rhizosphere soil of *Capsicum annuum*. MycoKeys 95: 1–13. <https://doi.org/10.3897/mycokeys.95.97062>

Abstract

Two new species, *Acremonium capsici* and *A. guizhouense*, isolated from the rhizosphere soil of *Capsicum annuum*, are described and illustrated. Two-locus DNA sequences based on phylogeny, in combination with the morphology of the asexual morph, were used to characterize these species. In the phylogenetic tree, both new species clustered into a monophyletic clade with strong support, distinct from other previously known species of *Acremonium*. The new species differed from their allied species in their morphology.

Keywords

filamentous fungi, morphology, new species, phylogeny, taxonomy

Introduction

Capsicum annuum L. is a globally grown and consumed spice crop that is rich in vitamins. *C. annuum* originated from the tropical and subtropical regions of Central and South America. It was introduced into China at the end of the Ming Dynasty, and has a long history of cultivation in China. According to the Food and Agriculture Organization of the United Nations, global *C. annuum* production reached approximately 36.1 million ton in 2020, with China producing the most in the world.

Link (1809) erected the genus *Acremonium*, whose members are geographically widespread and involve many substrates (Yang et al. 2019). As described by Gams (1971), the main diagnostic criteria of the genus *Acremonium* are conidiophores simple or verticillate; phialides narrow, solitary, generally cylindrical and gradually tapered towards the tips; conidia unicellular, hyaline to light-pigmented, spherical to cylindrical, arranged in slimy heads or unconnected chains or both; chlamydospores and sclerotia present or absent. The genus *Acremonium* is similar to some genera – *Sarocladium* W. Gams & D. Hawksw., *Brunneomyces* Giraldo, Gené & Guarro, and *Chordomyces* Bilanenko, M.L. Georgieva & Grum-Grzhimaylo etc. (Giraldo et al. 2015, 2017), including some of the simplest morphologies of all filamentous anamorphic fungi (Summerbell et al. 2011), so the morphological delimitation between them is challenging (Yang et al. 2019). Recent phylogenetic studies have documented that the genus *Acremonium* is polyphyletic, including sexual and nomenclaturally complex asexual morphs (Summerbell et al. 2011; Giraldo et al. 2012). To date, *Acremonium* has 219 records in the Index Fungorum (<http://www.indexfungorum.org/Names/Names.asp>, retrieval on Dec. 2022). However, many *Acremonium* taxa have been reported, but there are no trustworthy classification systems and little sequence data are available in GenBank for multigene analyses (Park et al. 2017). In the future, the classification of *Acremonium* will become clearer with the increase of molecular data.

In this study, seven strains of *Acremonium* were isolated in the process of investigating the rhizosphere fungal diversity of cultivated *Capsicum annuum* in Guizhou Province, southwest China, based on a culturable method. Identification of these strains in combination with morphological characteristics and phylogenetic analysis showed that these strains belong to two previously undescribed species of *Acremonium*. The new species differed from their allied species in their morphology.

Materials and methods

Fungal isolation and morphology

Capsicum annuum plants were cultivated in farmlands located in Guiyang, Guizhou Province, China (26°45'75"N, 106°64'87"E). One composite rhizosphere soil sample was taken from five randomly selected *C. annuum* plants. The roots were shaken vigorously to separate soil that is not tightly attached to the roots, and the remaining soil attached to the region 2–3 mm from the plant root was collected as the rhizosphere soil sample (Smalla et al. 2001). Fungi were isolated and purified using a dilution plate method as follows: 2 g samples were weighed with glass beads in a conical flask containing 20 mL sterile water, mixed evenly using eddy shock for 10 min, diluted to 1:10,000, and cultured on Martin's medium supplemented with chloramphenicol and cycloheximide.

The purified isolates were transferred to potato dextrose agar (PDA), oatmeal agar (OA), malt extract agar (MEA), and corn meal agar (CMA) at 25 °C in darkness for 14 days to examine the macroscopic and morphological characteristics of the colonies. Photomicrographs of the diagnostic structures were obtained using an OLYMPUS

BX53 microscope equipped with differential interference contrast optics, an OLYMPUS DP73 high-definition color camera, and cellSens software v.1.18. Both dry and living cultures were deposited at the Institute of Agro-bioengineering, Guizhou University.

DNA extraction, PCR amplification, and sequencing

Total DNA was extracted from each of the new isolates using the BioTeke Fungus Genomic DNA Extraction kit (DP2032, BioTeke, Beijing, China) according to the manufacturer's instructions. According to Li et al. (2022), the internal transcribed spacers (ITS), the 28S nrRNA locus (LSU), translation elongation factor 1-alpha gene region (*TEF 1- α*), RNA polymerase II second largest subunit gene (*RPB2*), and small subunit rDNA (SSU) were amplified and sequenced using ITS1/ITS4 (White et al. 1990), LROR/LR7 (Vilgalys and Hester 1990), EF1-983F/EF1-2218R (Rehner and Buckley 2005), fRPB2-5f/fRPB2-7cR (Liu et al. 1999), and NS1/NS4 (White et al. 1990) primers, respectively. All new sequences were submitted to GenBank (Table 1).

Phylogenetic analyses

In this study, we utilized sequence data mainly from recent publications (Yang et al. 2019; Li et al. 2022) and the sequenced new isolates (Table 1). According to Li et al. (2022) and Yang et al. (2019), *Pestalotiopsis spathulata* (CBS 356.86) and *P. hawaiiensis* (CBS 114491) were chosen as the outgroup taxa. The sequences were aligned using MAFFT v7.037 (Katoh and Standley 2013) and adjusted using MEGA 6.06 (Tamura et al. 2013). The aligned sequences of LSU and ITS were concatenated using PhyloSuite v1.16 (Zhang et al. 2020).

The best-fit substitution model was selected using the corrected Akaike information criterion, in ModelFinder (Kalyaanamoorthy et al. 2017). The maximum likelihood (ML) and Bayesian inference (BI) methods were used in the analysis. The ML analysis was implemented in IQ-TREE v1.6.11 (Nguyen et al. 2015) with 10,000 bootstrap tests, using the ultrafast algorithm (Minh et al. 2013). For the BI, MrBayes v3.2 (Ronquist et al. 2012) was used and Markov chain Monte Carlo simulations were run for 5,000,000 generations with a sampling frequency of every 500 generations and a burn-in of 25%. The above analyses were carried out in PhyloSuite v1.16 (Zhang et al. 2020).

Results

Phylogenetic analyses

Ninety-five isolates (including the seven with new sequence data) were included in our dataset (Table 1), which comprised 976 positions (including gaps), of which 377 were phylogenetically informative (122 of LSU and 255 of ITS). For Maximum-likelihood analyses, IQ-TREE's ModelFinder under the corrected Akaike information

Table I. Strains included in the present study.

Species	Strains	LSU	ITS	SSU	TEF 1- α	RPB2
<i>Acremonium alternatum</i>	CBS 407.66 T	HQ231988	HE798150			
<i>Acremonium alternatum</i>	CBS 831.97	HQ231989				
<i>Acremonium arthrinii</i>	MFLU 18-1225 T	MN036334		MN036335	MN038169	
<i>Acremonium behniae</i>	CBS 146824 T	MW175400	MW175360			
<i>Acremonium biseptum</i>	CBS 750.69 T	HQ231998				
<i>Acremonium blocchii</i>	CBS 993.69	HQ232002	HE608636			
<i>Acremonium borodinense</i>	CBS 101148 T	HQ232003	HE608635			
<i>Acremonium brachyphenum</i>	CBS 866.73 T	HQ232004	AB540570			
<i>Acremonium camposporum</i>	CBS 756.69 T	HQ232008		HQ232186		
<i>Acremonium cavaeaeum</i>	CBS 101149 T	HF680202	HF680220			
<i>Acremonium cavaeaeum</i>	CBS 111656	HF680203	HF680221			
<i>Acremonium cavaeaeum</i>	CBS 758.69	HQ232012	HF680222			
<i>Acremonium cerealis</i>	CBS 207.65	HQ232013				
<i>Acremonium cerealis</i>	CBS 215.69	HQ232014				
<i>Acremonium chiangaiense</i>	MFLUCC 14-0397 T	MN648329	MN648324			
<i>Acremonium chrysogenum</i>	CBS 144.62 T	HQ232017		HQ232187		
<i>Acremonium chrysogenum</i>	CBS 401.65	MH870276	MH858636			
<i>Acremonium citrinum</i>	CBS 384.96 T	HF680217	HF680236			
<i>Acremonium curvum</i>	CGMCC 3.20954 T	ON041050	ON041034	ON876754	ON494579	ON494583
<i>Acremonium dimorphosporum</i>	CBS 139050 T	LN810506	LN810515			
<i>Acremonium exiguum</i>	CBS 587.73 T	HQ232035				
<i>Acremonium exuviarum</i>	UAMH 9995 T	HQ232036	AY882946			
<i>Acremonium felinum</i>	CBS 147.81 T	AB540488	AB540562			
<i>Acremonium flavum</i>	CBS 596.70 T	HQ232037		HQ232191		
<i>Acremonium flavum</i>	CBS 316.72	MH872204	MH860487			
<i>Acremonium fuci</i>	CBS 112868 T		AY632653			
<i>Acremonium fuci</i>	CBS 113889		AY632652			
<i>Acremonium fusidioides</i>	CBS 109069	HF680204	HF680223			
<i>Acremonium fusidioides</i>	CBS 991.69	HF680211	HF680230			
<i>Acremonium fusidioides</i>	CBS 840.68 T	HQ232039	FN706542			
<i>Acremonium globosiporum</i>	CGMCC 3.20955 T	ON041051	ON041035	ON876755	ON494580	ON494584
<i>Acremonium globosiporum</i>	GZUIFR 22.037	ON041052	ON041036	ON876756	ON494581	ON494585
<i>Acremonium globosiporum</i>	GZUIFR 22.038	ON041053	ON041037	ON876757	ON494582	ON494586
<i>Acremonium hansfordii</i>	CBS 390.73	HQ232043	AB540578			
<i>Acremonium hennebertii</i>	CBS 768.69 T	HQ232044	HF680238			
<i>Acremonium inflatum</i>	CBS 212.69 T	HQ232050				
<i>Acremonium mali</i>	ACCC 39305 T	MF993114	MF987658			
<i>Acremonium moniliforme</i>	CBS 139051 T	LN810507	LN810516			
<i>Acremonium moniliforme</i>	FMR 10363	LN810508	LN810517			
<i>Acremonium parvum</i>	CBS 381.70A	HQ231986	HF680219			
<i>Acremonium persicinum</i>	CBS 310.59 T	HQ232077				
<i>Acremonium persicinum</i>	CBS 101694	HQ232085				
<i>Acremonium pinkertoniae</i>	CBS 157.70 T	HQ232089		HQ232202		
<i>Acremonium polychroma</i>	CBS 181.27 T	HQ232091	AB540567			
<i>Acremonium potronii</i>	CBS 189.70	HQ232094				
<i>Acremonium pseudozeylanicum</i>	CBS 560.73 T	HQ232101				
<i>Acremonium pteridii</i>	CBS 782.69 T	HQ232102				
<i>Acremonium pteridii</i>	CBS 784.69	HQ232103				
<i>Acremonium sclerotigenum</i>	CBS 124.42 T	HQ232126	FN706552	HQ232209		
<i>Acremonium sclerotigenum</i>	A101	KC987215	KC987139	KC987177	KC998961	
<i>Acremonium sclerotigenum</i>	A130	KC987242	KC987166	KC987204	KC998988	
<i>Acremonium</i> sp.	E102	KC987248	KC987172	KC987210	KC998994	KC999030
<i>Acremonium spinosum</i>	CBS 136.33 T	HQ232137	HE608637	HQ232210		

Species	Strains	LSU	ITS	SSU	TEF 1- α	RPB2
<i>Acremonium stroudii</i>	CBS 138820 T		KM225291			
<i>Acremonium tumulicola</i>	CBS 127532 T	AB540478	AB540552			
<i>Acremonium variecolor</i>	CBS 130360 T	HE608651	HE608647			
<i>Acremonium variecolor</i>	CBS 130361	HE608652	HE608648			
<i>Acremonium verruculosum</i>	CBS 989.69 T	HQ232150				
<i>Acremonium capsici</i>	SQT01 T	OP740978	OP703286	OP750190	OP757287	OP730522
<i>Acremonium capsici</i>	SQT02	OP740979	OP703287	OP750191	OP757288	OP730523
<i>Acremonium capsici</i>	SQT03	OP740980	OP703288	OP750192	OP757289	OP730524
<i>Acremonium guizhouense</i>	SQT04 T	OP740981	OP703289	OP750193	OP757290	OP730525
<i>Acremonium guizhouense</i>	SQT05	OP740982	OP703290	OP750194	OP757291	OP730526
<i>Acremonium guizhouense</i>	SQT06	OP740983	OP703291	OP750195	OP757292	OP730527
<i>Acremonium guizhouense</i>	SQT07	OP740984	OP703292	OP750196	OP757293	OP730528
<i>Bryocentria bronniartii</i>	M139	EU940105		EU940052		
<i>Bryocentria bronniartii</i>	M190	EU940125		EU940052		
<i>Bryocentria metzgeriae</i>	M140	EU940106				
<i>Bulbithecium hyalosporum</i>	CBS 318.91 T	AF096187	HE608634			
<i>Cephalosporium purpurascens</i>	CBS 149.62 T	HQ232071				
<i>Cosmospora lavitskiae</i>	CBS 530.68 T	HQ231997				
<i>Emericellopsis alkalina</i>	CBS 127350 T	KC987247	KC987171	KC987209	KC998993	KC999029
<i>Emericellopsis terricola</i>	CBS 120.40 T	U57082	U57676	U44112		
<i>Gliomastix roseogrisea</i>	CBS 134.56 T	HQ232121				
<i>Hapsidospora irregularis</i>	ATCC 22087 T	AF096192		AF096177		
<i>Kiflmonium curvulum</i>	CBS 430.66 T	HQ232026	HE608638	HQ232188		
<i>Lanatonectria flavolanata</i>	CBS 230.31	HQ232157				
<i>Leucosphaerina arxii</i>	CBS 737.84 T	HE608662	HE608640			
<i>Nigrosabulum globosum</i>	ATCC 22102 T	AF096195				
<i>Paracremonium contagium</i>	CBS 110348 T	HQ232118	KM231831		KM231966	
<i>Parasarocladium breve</i>	CBS 150.62 T	HQ232005				
<i>Parasarocladium radiatum</i>	CBS 142.62 T	HQ232104		HQ232205		
<i>Pestalotiopsis hawaiiensis</i>	CBS 114491 T	KM116239	KM199339		KM199514	
<i>Pestalotiopsis spathulata</i>	CBS 356.86 T	KM116236	KM199338		KM199513	
<i>Pseudoacremonium sacchari</i>	CBS 137990 T	KJ869201	KJ869144			
<i>Sarcopodium vanillae</i>	CBS 100582	HQ232174	KM231780		KM231911	
<i>Sarocladium bacillisporum</i>	CBS 425.67 T	HQ231992	HE608639	HQ232179		
<i>Sarocladium bactrocephalum</i>	CBS 749.69 T	HQ231994	HG965006	HQ232180		
<i>Sarocladium strictum</i>	CBS 346.70 T	HQ232141	AY214439	HQ232211		
<i>Sarocladium terricola</i>	CBS 243.59 T	HQ232046		HQ232196		
<i>Selinia pulchra</i>	AR 2812	GQ505992	HM484859		HM484841	
<i>Trichothecium crotocinigenum</i>	CBS 129.64 T	HQ232018	AJ621773			
<i>Trichothecium indicum</i>	CBS 123.78T	AF096194		AF096179		
<i>Trichothecium roseum</i>	DAOM 208997	U69891		U69892		
<i>Trichothecium sympodiale</i>	ATCC 36477	U69889		U69890		

Notes: "T" stands for Ex-type strains. New isolates are in bold and blue.

criterion (AICc) proposed a TN+F+I+G4 for LSU, GTR+F+I+G4 for ITS. For Bayesian analysis, IQ-TREE's ModelFinder under the AICc proposed a GTR+F+G4 for LSU, GTR+F+I+G4 for ITS. The results show that the isolates SQT01, SQT02, and SQT03 clustered in a single clade with high support (ML BS 100/BI pp 1), and were closely related to *Acremonium variecolor* (Fig. 1). The isolates SQT04, SQT05, SQT06, and SQT07 also clustered in a single clade with high support (100/0.98), and were closely related to *A. persicinum* and *A. verruculosum* (Fig. 1).




Figure 1. Phylogram generated from maximum likelihood analysis based on LSU + ITS sequence data. Bootstrap support values of maximum likelihood higher than 75% and Bayesian posterior probabilities greater than 0.75 are given above each branch. The new collection is highlighted in blue bold. Clades are identified using clade nomenclature formally defined by Summerbell et al. (2011), and Yang et al. (2019). Ex-type strains are indicated by “T”.

Taxonomy

Acremonium capsici S.Q. Tong & Y.J. Wu, sp. nov.

MycoBank No: 846330

Fig. 2

Etymology. Referring to the type strain isolated from the rhizosphere soil of *Capsicum annuum*.




Figure 2. Morphology of *Acremonium capsici* sp. nov. **a-d** colony on PDA, MEA, OA, and CMA after 14 days at 25 °C (upper surface and lower surface) **e** phialides **f** conidia **g** phialides arising from ropes of hyphae **h** phialides arising from hyphal coils. Scale bars: 10 µm (**e-h**).

Type. Guiyang City, Guizhou Province, China 26°45'75"N, 106°64'87"E, isolated from the rhizosphere soil of *Capsicum annuum*, August 2022, Shuo-Qiu Tong (dried holotype culture SQT H-01, ex-holotype culture SQT01). GenBank: ITS = OP703286; LSU = OP740978; SSU = OP750190; TEF 1- α = OP757287; RPB2 = OP730522.

Description. Culture characteristics (14 days at 25 °C) – Colonies on PDA 20–21 mm diam, white, hairy, flat, radially striated, with a regular edge; reverse white. Colonies on MEA 18–19 mm in diameter, white, floccose, radially striated, with a regular edge; reverse white. Colonies on OA 18–19 mm in diameter, pale white, flat, with regular edge; reverse pale white. Colonies on CMA 18–19 mm in diameter, pale white, felty, with regular edge; reverse pale white. **Hypphae** hyaline, smooth, septate, branched, 1.0–2.5 μm wide. **Phialides** straight to flexuous, hyaline, smooth, arising from superficial hyphae, from hyphal strands or from hyphal coils, 20–42 μm (n = 50) long, 1–2 μm (n = 50) wide at the base. **Conidia** arranged in slimy heads, one-celled, ovoid to ellipsoidal, fusiform, 2.0–3.5 × 1.5–2.0 μm (n = 50), hyaline, smooth, or rough. **Chlamydospores** and teleomorph were not observed.

Additional specimens examined. Guiyang City, Guizhou Province, China 26°45'75"N, 106°64'87"E, isolated from the rhizosphere soil of *Capsicum annuum*, August 2022, Shuo-Qiu Tong, SQT02, *ibid.*, SQT03. GenBank: ITS = OP703287–OP703288; LSU = OP740979–OP740980; SSU = OP750191–OP750192; TEF 1- α = OP757288–OP757289; RPB2 = OP730523–OP730524.

Known distribution. Guiyang City, Guizhou Province, China.

Notes. In a phylogenetic tree based on LSU + ITS sequences, *Acremonium capsici* forms a separate clade sister to *A. variecolor* in *Acremonium sensu lato* (Bionectriaceae). In a comparison of LSU and ITS nucleotides, *A. capsici* (Type strain SQT01) has 93% and 83% similarity, in LSU (459/492 bp, one gap) and ITS (388/468 bp, 16 gaps), which is different from *A. variecolor* (CBS 130360). They are distinguished by the appearance of colonies on OA, MEA, and PDA: colonies of *A. capsici* grow slowly (less than 25 mm), and are white, while colonies of *A. variecolor* grow faster (more than 40 mm), and are white to yellowish (Giraldo et al. 2012). In addition, *A. capsici* bear simple phialides, while *conidiophores* of *A. variecolor* are mostly branched, bearing whorls of two to five phialides (Giraldo et al. 2012). *A. variecolor* produces sessile conidia, which is not seen in *A. capsici* (Giraldo et al. 2012).

Acremonium guizhouense S.Q. Tong & Y.J. Wu, sp. nov.

Mycobank No: 846331

Fig. 3

Etymology. Referring to the country where this fungus was first isolated.

Type. Guiyang City, Guizhou Province, China 26°45'75"N, 106°64'87"E, isolated from the rhizosphere soil of *Capsicum annuum*, August 2022, Shuo-Qiu Tong (dried holotype culture SQT H04, ex-holotype culture SQT04). GenBank: ITS = OP703289; LSU = OP740981; SSU = OP750193; TEF 1- α = OP757290; RPB2 = OP730525.

Description. Culture characteristics (14 days at 25 °C) – Colonies on PDA 16–19 mm in diameter, yellowish white to grayish yellow, flat, zonate, with regular edge; reverse brownish orange. Colonies on MEA 9–13 mm in diameter, yellowish white to white, compact, convex with papillate surface, margin dentate, aerial mycelia extremely sparse; reverse yellowish white to umber. Colonies on OA 14–16 mm in diameter, pale, felty, with regular edge; reverse pale white. Colonies on CMA 16–14 mm in diameter, pale white, felty, with regular edge; reverse pale white. **Hypphae** hyaline, smooth, septate,




Figure 3. Morphology of *Acremonium guizhouense* sp. nov. **a-d** colony on PDA, MEA, OA, and CMA after 14 days at 25 °C (upper surface and lower surface) **e, f** phialides and conidia **g, h** conidia are held together in slimy heads. Scale bars: 10 µm (**e-h**).

branched, 1.0–3.0 μm wide. ***Phialides*** straight to flexuous, hyaline, smooth, arising from hyphae, 15.5–33.5 μm ($n = 50$) long, 1.5–2.5 μm ($n = 50$) wide at the base. ***Conidia*** gathered in slimy heads, one-celled, ovoid to ellipsoidal, 2.5–3.0 \times 3.5–5.0 μm ($n = 50$), hyaline, smooth or rough. ***Chlamydospores*** and teleomorph not observed.

Additional specimens examined. Guiyang City, Guizhou Province, China 26°45'75"N, 106°64'87"E, isolated from the rhizosphere soil of *Capsicum annuum*, August 2022, Shuo-Qiu Tong, SQT05 = SQT06, *ibid.*, SQT07. GenBank: ITS = OP703290–OP703292; LSU = OP740982–OP740984; SSU = OP750194–OP750196; TEF 1- α = OP757291–OP757293; RPB2 = OP730526–OP730528.

Known distribution. Guiyang City, Guizhou Province, China.

Notes. Phylogenetic and morphological data (Figs 1, 3) support our isolates SQT04–SQT07 as new species of *Acremonium*. *A. guizhouense* is phylogenetically closely related to *A. verruculosum* and *A. persicinum*. However, they can be distinguished by their sequence similarity (97% similarity, 10 base pairs (bp) differences and two gaps in 497 bp of LSU in *A. verruculosum* CBS 989.69; 98% similarity, 12 base pairs (bp) differences, and four gaps in 809 bp of LSU in *A. persicinum* CBS310.59). Since *A. verruculosum* and *A. persicinum* lack ITS sequences, it was not possible to compare *A. guizhouense* with them. Morphologically, the conidia of *A. verruculosum* are long ellipsoidal to cylindrical, rather than ovoid to ellipsoidal in *A. guizhouense* (Gams 1971). *A. verruculosum*, on the other hand, has larger conidia than *A. guizhouense* (5.6–6.0 \times 2.3–2.5 μm vs. 2.5–3.0 \times 3.5–5.0 μm) (Gams 1971). Furthermore, conidia of *A. verruculosum* are catenate, fusiform, pyriform to ellipsoidal rather than arranged as slimy heads, ovoid to ellipsoidal in *A. guizhouense* (Gams 1971). The conidia of *A. guizhouense*, on the other hand, are smaller than that of *A. persicinum* (2.5–3.0 \times 3.5–5.0 μm vs. 3.2–4.8 \times 1.2–3.0 μm) (Gams 1971).

Discussion

Traditionally, a polyphasic approach based on morphology, physiology, biochemistry, or reactions to chemical tests, has been used to differentiate species (Senanayake et al. 2020). Currently, many new fungal taxa have been reported based on DNA sequences. Phylogenetic analysis is becoming increasingly important in reporting new taxa of fungi, and has gradually become a mandatory component. However, many previously published fungal taxa lack DNA molecular data, and even specimens have been lost (Zhang et al. 2022). Thus, there are still many undetermined, questionable, or misidentified taxa that warrant taxonomic investigations (Summerbell et al. 2018). Since most species of the genus *Acremonium* have only LSU and ITS sequences Li et al. (2022), we used only ribosomal sequences (LSU + ITS) for phylogenetic analysis, while the sequencing of other loci was aimed at establishing a database for future studies.

Members of the genus *Acremonium* are geographically widespread and ecologically diverse, and seem to colonize all types of substrates, including endophytes, epiphytes, saprophytes, human and plant pathogens, lichens, insects, or arthropods taxa (Yang et

al. 2019). In addition, *Acremonium* species have various functions, such as biological control (Shang et al. 2018), enhancing drought tolerance of grasses, and promoting nectar production of beans (Jaber and Vidal 2009), as well as improving plant resistance to plant pathogens (Kasselaki et al. 2006). In the present study, all the isolates were obtained from the rhizosphere soils of *Capsicum annuum*. Therefore, more studies are necessary to further confirm their relationship with their host plant *Capsicum annuum*.

In summary, seven isolates of *Acremonium* were obtained from the rhizosphere soils of *Capsicum annuum*. Morphological characteristics in combination with two-locus (LSU + ITS) phylogenetic analysis were used for delimitation. Therefore, two new species of *Acremonium capsici* (three isolates) and *Acremonium guizhouense* (four isolates) are introduced. This study contributes to our understanding of the rhizosphere microbial population of *Capsicum annuum* and also of *Acremonium* species.

Acknowledgements

This study was financially supported by Guizhou Provincial Science and Technology Projects (Qian Ke He Zhi Cheng-ZK[2022] General 172, and ZK[2021] General 262), and Guizhou Provincial Institutions of higher learning Engineering Center Projects (Qian Jiao He KY[2021]006). We appreciate Charlesworth for English-language editing of the whole manuscript.

References

- Gams W (1971) *Cephalosporium*-artige Schimmelpilze (Hyphomycetes). G. Fischer, Stuttgart.
- Giraldo A, Gené J, Cano J, de Hoog S, Guarro J (2012) Two new species of *Acremonium* from Spanish soils. *Mycologia* 104(6): 1456–1465. <https://doi.org/10.3852/11-402>
- Giraldo A, Gené J, Sutton DA, Madrid H, de Hoog GS, Cano J, Decock C, Crous PW, Guarro J (2015) Phylogeny of *Sarocladium* (Hypocreales). *Persoonia* 34(1): 10–24. <https://doi.org/10.3767/003158515X685364>
- Giraldo A, Gené J, Sutton DA, Wiederhold N, Guarro J (2017) New acremonium-like species in the Bionectriaceae and Plectosphaerellaceae. *Mycological Progress* 16(4): 349–368. <https://doi.org/10.1007/s11557-017-1271-7>
- Jaber LR, Vidal S (2009) Interactions between an endophytic fungus, aphids, and extrafloral nectaries: Do endophytes induce extrafloral-mediated defences in *Vicia faba*? *Functional Ecology* 23(4): 707–714. <https://doi.org/10.1111/j.1365-2435.2009.01554.x>
- Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Kasselaki AM, Shaw MW, Malathrakis NE, Haralambous J (2006) Control of *Leveillula taurica* in tomato by *Acremonium alternatum* is by induction of resistance, not hyperparasitism. *European Journal of Plant Pathology* 115(2): 263–267. <https://doi.org/10.1007/s10658-006-9010-y>

- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Li X, Zhang ZY, Ren YL, Chen WH, Liang JD, Pan JM, Huang JZ, Liang ZQ, Han YF (2022) Morphological characteristics and phylogenetic evidence reveal two new species of *Acremonium* (Hypocreales, Sordariomycetes). *MycoKeys* 91: 85–96. <https://doi.org/10.3897/mycokeys.91.86257>
- Link HF (1809) *Observationes in ordines plantarum naturales: Dissertatio I.* Magazin. der Gesellschaft Naturforschenden Freunde Berlin 3: 3–42.
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among Ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Minh Q, Nguyen M, von Haeseler AA (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Park SW, Nguyen TTT, Lee HB (2017) Characterization of two species of *Acremonium* (unrecorded in Korea) from soil samples: *A. variecolor* and *A. persicinum*. *Mycobiology* 45(4): 353–361. <https://doi.org/10.5941/MYCO.2017.45.4.353>
- Rehner SA, Buckley E (2005) A *Beauveria* phylogeny inferred from nuclear ITS and EF1- α sequences: Evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* 97(1): 84–98. <https://doi.org/10.3852/mycologia.97.1.84>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere: Journal of Fungal Biology* 11(1): 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Shang SQ, Chen YN, Bai YL (2018) The pathogenicity of entomopathogenic fungus *Acremonium hansfordii* to two-spotted spider mite, *Tetranychus urticae* and predatory mite *Neoseiulus barkeri*. *Systematic and Applied Acarology* 23(11): 2173–2183. <https://doi.org/10.111158/saa.23.11.10>
- Smalla K, Wieland G, Buchner A, Zock A, Parzy J, Kaiser S, Roskot N, Heuer H, Berg G (2001) Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: Plant-dependent enrichment and seasonal shifts revealed. *Applied and Environmental Microbiology* 67(10): 4742–4751. <https://doi.org/10.1128/AEM.67.10.4742-4751.2001>

- Summerbell RC, Gueidan C, Schroers H-J, de Hoog GS, Starink M, Rosete YA, Guarro J, Scott JA (2011) *Acremonium* phylogenetic overview and revision of *Gliomastix*, *Sarocladium*, and *Trichothecium*. Studies in Mycology 68: 139–162. <https://doi.org/10.3114/sim.2011.68.06>
- Summerbell RC, Gueidan C, Guarro J, Eskalen A, Crous PW, Gupta AK, Gené J, Cano-Lira JF, van Iperen A, Starink M, Scott JA (2018) The protean *Acremonium*. *A. sclerotigenum/egyptiacum*: Revision, food contaminant, and human disease. Microorganisms 6(3): 88. <https://doi.org/10.3390/microorganisms6030088>
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. Molecular Biology and Evolution 30(12): 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: a guide to methods and applications, Academic Press, San Diego, California, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Yang CL, Xu XL, Jeewon R, Boonmee S, Liu YG, Hyde KD (2019) *Acremonium arthrinii* sp. nov., a mycopathogenic fungus on *Arthrinium yunnanum*. Phytotaxa 420: 283–299. <https://doi.org/10.11646/phytotaxa.420.4.4>
- Zhang D, Gao FL, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20(1): 348–355. <https://doi.org/10.1111/1755-0998.13096>
- Zhang ZY, Ren YL, Chen WH, Liang JD, Han YF, Liang ZQ (2022) New taxonomic framework for Arthrodermataceae: A comprehensive analysis based on their phylogenetic reconstruction, divergence time estimation, phylogenetic split network, and phylogeography. Antonie van Leeuwenhoek 115(11): 1319–1333. <https://doi.org/10.1007/s10482-022-01774-0>